



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 3697 Type: **Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)**

(G*) Covariant Approach in Bethe-Heitler process for the calculation of electron-proton Scattering

Monday 19 June 2023 17:00 (15 minutes)

In order to search for the physics beyond the Standard Model at the precision frontier, it is sometimes essential to account for Next-to-Next-Leading Order (NNLO) corrections theoretical calculations. Using the covariant approach, we calculated the QED type leptonic tensor up to quadratic (one loop squared) NNLO (alpha cube) order, which can be used for the processes like (electron-proton) and (muon-proton) scattering relevant to MOLLER (background studies) and MUSE experiments, respectively. Recently we have used this approach for a hard photon bremsstrahlung process called "Bethe-Heitler". This is a 2->3 process where an electron scatters with a proton with the emission of a hard photon and is an important example in Quantum Electrodynamics (QED).

In this presentation, I will quickly review covariant approach and provide our latest results for quadratic QED electron-proton scattering along with the Bethe-Heitler process.

Keyword-1

electron-proton Scattering

Keyword-2

Covariant Approach

Keyword-3

Bethe-Heitler process

Authors: Dr ALEKSEJEVES, Aleksandrs (Memorial University of Newfoundland); Ms GHAFFAR, Mahumm (Memorial University of Newfoundland)

Presenter: Ms GHAFFAR, Mahumm (Memorial University of Newfoundland)

Session Classification: (DNP) M3-4 Testing the Standard Model at low and intermediate energies | Tester le modèle standard à des énergies faibles et intermédiaires (DPN)

Track Classification: Technical Sessions / Sessions techniques: Nuclear Physics / Physique nucléaire (DNP-DPN)