

Contribution ID: 3817

Canadian Association of Physicists

Association canadienne des physiciens et physiciennes

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Seeing the strongly-correlated zero-bias anomaly in double quantum dot measurements

Wednesday 21 June 2023 11:00 (15 minutes)

Experiments in doped transition metal oxides often show suppression in the single-particle density of states at the Fermi level, but disorder-induced zero-bias anomalies in strongly-correlated systems remain poorly understood. Numerical studies of the Anderson-Hubbard model have identified a zero-bias anomaly that is unique to strongly correlated materials, with a width proportional to the inter-site hopping amplitude t.[PRL **101**, 086401 (2008)] In ensembles of two-site systems, a zero-bias anomaly with the same parameter dependence also occurs, suggesting a similar physical origin.[PRB **82**, 073107 (2010)] We describe how this kinetic-energy-driven zero-bias anomaly in ensembles of two-site systems may be seen in a mesoscopic realization based on double quantum dots. Moreover, the double-quantum-dot measurements provide access not only to the ensemble-average density of states but also to the details of the transitions which give rise to the zero-bias anomaly.

Keyword-1

double quantum dot

Keyword-2

strong correlations

Keyword-3

zero-bias anomaly

Author: WORTIS, Rachel

Co-authors: FOLKS, Joshua (U. of British Columbia); LUESCHER, Silvia (U. of British Columbia); LUYBEN, Sylvia (U. of Guelph)

Presenter: WORTIS, Rachel

Session Classification: (DCMMP) W1-7 Condensed Matter Theory I | Théorie de la matière condensée I (DPMCM)

Track Classification: Technical Sessions / Sessions techniques: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)