



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 4429

Type: **Invited Speaker / Conférencier(ère) invité(e)**

## Twisting high intensity lasers to produce extreme magnetic fields

*Tuesday 28 May 2024 11:00 (30 minutes)*

Recently, orbital angular momentum (OAM) beams have demonstrated at relativistic intensities at several high-power laser facilities around the world using off-axis spiral phase mirrors. The additional angular momentum carried by OAM beams, even when linearly polarized, introduces a new control parameter in laser plasma interactions and has shown promise to introduce new and exciting phenomena not possible with a standard Gaussian beam.

Of particular interest is the relativistic inverse Faraday effect where laser angular momentum is absorbed by a plasma generating large axial magnetic fields colinear with the laser  $k$  vector. Our recent work has demonstrated that magnetic fields on the order of 100's of Tesla, extending 100's of microns, and lasting on the order of 10 picoseconds can be generated with laser powers less than 5 terawatts. In this work we will explore this phenomenon through theory, simulations, and present results from a recent campaign at the COMET laser at Lawrence Livermore National Laboratory in which we used a linearly polarized Laguerre Gaussian laser to drive magnetic fields for the first time in the laboratory. Experimental results will be compared and validated against theory and simulations.

### Keyword-1

Orbital Angular Momentum Laser

### Keyword-2

Kilo Tesla Magnetic Fields

### Keyword-3

Inverse Faraday Effect

**Author:** LONGMAN, Andrew (Lawrence Livermore National Laboratory)

**Presenter:** LONGMAN, Andrew (Lawrence Livermore National Laboratory)

**Session Classification:** (DPP) T1-6 Laser Plasma Interaction | Interaction laser-plasma (DPP)

**Track Classification:** Technical Sessions / Sessions techniques: Plasma Physics / Physique des plasmas (DPP)