

Contribution ID: 4431 Type: Poster Competition (Graduate Student) / Compétition affiches (Étudiant(e) 2e ou 3e cycle)

(G*) (POS-73) Diffuse emission from black hole remnants

Tuesday 28 May 2024 18:12 (2 minutes)

At the end of its evaporation, a black hole may leave a remnant where a large amount of information is stored. We argue that the existence of an area gap as predicted by Loop Quantum Gravity removes a main objection to this scenario. Remnants should radiate in the low-frequency spectrum. We model this emission and derive properties of the diffuse radiation emitted by a population of such objects. We show that the frequency and energy density of this radiation, which are measurable in principle, suffice to estimate the mass of the parent holes and the remnant density, if the age of the population is known.

Keyword-1

Black Holes

Keyword-2

Quantum Gravity

Keyword-3

Cosmology

Authors: ROVELLI, Carlo; Prof. VIDOTTO, Francesca (The University of Western Ontario); PASCUAL, Mateo; KAZEMIAN, Sina (University of Western Ontario)

Presenter: PASCUAL, Mateo

Session Classification: DTP Poster Session & Student Poster Competition (11) | Session d'affiches DPT et concours d'affiches étudiantes (11)

Track Classification: Technical Sessions / Sessions techniques: Theoretical Physics / Physique théorique (DTP-DPT)