



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 19 Type: Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)

## Modelling electroweak processes in light nuclei: Nuclear Fermi decays and $V_{ud}$

*Tuesday 10 June 2025 15:00 (15 minutes)*

Currently, the most precise extraction of the up-down quark mixing element  $V_{ud}$  of the Cabibbo-Kobayashi-Maskawa (CKM) matrix comes from a handful of  $ft$ -value measurements for nuclear Fermi decays in light- and medium-mass nuclei. However, a complete extraction of  $V_{ud}$  from hadronic decays requires challenging theory determinations of hadronic-structure-dependent electroweak radiative corrections (EWRC) to said decays. In fact, a novel evaluation of the free-hadron part of the dominant correction to nuclear Fermi decays, i.e., the free-hadron  $\gamma W$ -box diagram, has led to tension with the Standard Model expectation of CKM unitarity. Moreover, to reach the current precision goals for the CKM unitarity test via extraction of  $V_{ud}$  from nuclear Fermi decays, a consistent treatment of the hadronic-structure-dependent EWRCs in the nuclear medium is critical. Confirmation of this tension by way of increasingly precise nuclear theory and experiment would point towards a deficiency in the Standard Model (SM) weak sector.

Ultimately, this amounts to requiring a modern evaluation of the nuclear  $\gamma W$ -box diagram utilizing the ever-advancing set of tools available in nuclear many-body theory. Targeting an evaluation of the  $\gamma W$ -box diagram for the  $^{10}\text{C} \rightarrow ^{10}\text{B}$  and  $^{14}\text{O} \rightarrow ^{14}\text{N}$  Fermi transitions, we apply the no-core shell model (NCSM). The NCSM is a non-relativistic quantum many-body approach for modelling the low-lying bound states of light nuclei starting solely from inter-nucleonic interactions. Augmented by the Lanczos strengths method, the NCSM can be further utilized to target features of the entire many-body spectrum, a capability without which calculations of this kind would not be possible. The approach detailed represents one of the first utilized to compute these corrections in ab initio nuclear theory.

### Keyword-1

Precision electroweak tests

### Keyword-2

Nuclear many-body theory

### Keyword-3

Fundamental symmetries

**Author:** Mr GENNARI, Michael

**Co-authors:** SENG, Chien Yeah (University of Washington); DRISSI, Mehdi (TRIUMF); GORSHTEYN, Misha (Mainz University); NAVRATIL, Petr

**Presenter:** Mr GENNARI, Michael

**Session Classification:** (DNP) T2-6 Nuclear Theory | Théorie nucléaire (DPN)

**Track Classification:** Technical Sessions / Sessions techniques: Nuclear Physics / Physique nucléaire (DNP-DPN)