

# Dark Matter Searches at Super-Kamiokande



**Piotr Mijakowski**




National Centre For Nuclear Research, Warsaw, Poland


(c) Kamioka Observatory, ICRR(Institute for Cosmic Ray Research), The University of Tokyo

# OUTLINE

Indirect searches for dark matter induced neutrinos at Super Kamiokande:

1. Galactic Center & Halo 2017
2. Earth 2017
3. Sun 2015
4. Boosted dark matter 2017





# Super-Kamiokande

@ Kamioka Observatory (ICRR, University of Tokyo), Japan

located 1km underground

40m

40m




photomultipliers (PMTs) detect Cherenkov light



PMT



~11k ID  
~1.8k OD  
PMTs

- 50 kton of pure water (22.5 kton FV)
- inner (ID) & outer/veto (OD) detection regions
- SK runs from 1996
- measures solar, atmospheric, cosmic & accelerator neutrinos
- Far detector of **T2K**




## Detected Cherenkov light allows for reconstruction of:



- lepton momentum (neutrino energy)
- lepton direction
- lepton flavor (e-like vs.  $\mu$ -like, good separation possible)



# Atmospheric neutrinos: main background in DM-induced $\nu$ searches




## atmospheric neutrinos at SK



- $\sim 10$  events/day
- data period: 1996-2016
- $\sim 50\,000$  events in total

# Dark matter searches at Super-Kamiokande



- Search for excess of neutrinos from **Earth/Sun/Milky Way**
- **FIT**: for each tested WIMP mass, find configuration of **ATM  $\nu$**  + **DM** signal that would match **DATA** the best



Galactic WIMP search

Earth WIMP search

Solar WIMP search  
point-like source



- +— SK DATA
- ATM MC (BKG) with oscillations
- WIMP signal enhanced for illustration

- In these coordinate systems signal is easy to distinguish from atmospheric neutrino background

# Signal simulation


Simulate DM signal before detection  $\rightarrow$  DarkSUSY & WimpSim

P. Gondolo et al., JCAP 07, 008 (2004)

M. Blennow et al., arXiv: 0709.3898 (2008)



## EXAMPLE: Galactic WIMP search

differential  $\nu_\mu \bar{\nu}_\mu$  energy spectra per DM annihilation for  $M_\chi = 100$  GeV (oscillated throughout Galaxy)



## EXAMPLE: Earth WIMP search

muon neutrino flux produced in WIMP annihilation in the Earth's core




# Galactic WIMP search

- diffuse signal from entire Galaxy, peaked from Galactic Center
- GC visibility with SK:  
~71% with UPMU, 100% FC/PC
- search constrains DM self-annihilation cross section  $\langle\sigma V\rangle$



Detector




Expected signal intensity strongly depends on halo model  
NFW is considered as a benchmark model in this analysis



# Galactic WIMP search: data

- FIT based on lepton mom. &  $\cos\theta_{GC}$  distributions, 5326-5629 live-days, 1996-2016
- NFW halo model assumed
- Fit results are consistent with null WIMP contribution
- 90% CL upper limit on DM self-annihilation cross section  $\langle\sigma_A V\rangle$

example: 5GeV WIMPs  $b\bar{b}$  ann. channel







proportions of the signal in various samples are reflected

# Galactic WIMP search: fitted number of DM-induced $V$ 's

- FIT based on lepton mom. &  $\cos\theta_{GC}$  distributions, 5326-5629 live-days, 1996-2016
- NFW halo model assumed
- Fit results are consistent with null WIMP contribution
- 90% CL upper limit on DM self-annihilation cross section  $\langle\sigma_A V\rangle$

SK preliminary




~150 systematic uncertainty terms included in the fit

p-values in backup

# Galactic WIMP search: DM self-annihilation cross section


- FIT based on lepton mom. &  $\cos\theta_{GC}$  distributions, 5326-5629 live-days, 1996-2016
- NFW halo model assumed
- Fit results are consistent with null WIMP contribution
- 90% CL upper limit on DM self-annihilation cross section  $\langle\sigma_A V\rangle$



$$\frac{d\phi_{\Delta\Omega}}{dE} = \frac{\langle\sigma_A \cdot V\rangle}{2} J_{\Delta\Omega} \frac{R_{sc} \rho_{sc}^2}{4\pi \cdot M_\chi^2} \frac{dN}{dE}$$

# Galactic WIMP search: DM self-annihilation cross section



- FIT based on lepton mom. &  $\cos\theta_{GC}$  distributions, 5326-5629 live-days, 1996-2016
- NFW halo model assumed
- Fit results are consistent with null WIMP contribution
- 90% CL upper limit on DM self-annihilation cross section  $\langle\sigma_A V\rangle$



$$\frac{d\phi_{\Delta\Omega}}{dE} = \frac{\langle\sigma_A \cdot V\rangle}{2} J_{\Delta\Omega} \frac{R_{sc} \rho_{sc}^2}{4\pi \cdot M_\chi^2} \frac{dN}{dE}$$

# Earth WIMP search

- Spin-independent interactions dominate in the capturing process → scalar interaction in which WIMPs couple to the nucleus mass
- If the mass of DM matches given heavy element, the capture rate increases considerably




The peaks correspond to **resonant capture** on the most abundant elements  $^{16}\text{O}$ ,  $^{24}\text{Mg}$ ,  $^{28}\text{Si}$  and  $^{56}\text{Fe}$  and their isotopes

**WIMP-nucleon SI scattering cross section  $\sigma_{\chi n}$**  can be constrained and compared with results from direct DM detection.

# Earth WIMP search: data

example: 25GeV WIMPs  $\tau^+\tau^-$  ann. channel

- FIT based on lepton mom. &  $\cos\theta_{\text{zenith}}$  distributions, 5326-5629 live-days, 1996-2016
- Fit results are consistent with null WIMP contribution
- 90 % upper limits on SI WIMP-nucleon scattering cross section  $\sigma_{\chi-n}$





proportions of the signal in various samples are reflected

# Earth WIMP search: fitted number of DM-induced $\nu$ s

- FIT based on lepton mom. &  $\cos\theta_{\text{zenith}}$  distributions, 5326-5629 live-days, 1996-2016
- Fit results are consistent with null WIMP contribution
- 90 % upper limits on SI WIMP-nucleon scattering cross section  $\sigma_{\chi\text{-n}}$


SK preliminary



# Earth WIMP search: WIMP-nucleon SI cross-section limit

- FIT based on lepton mom. &  $\cos\theta_{\text{zenith}}$  distributions, 5326-5629 live-days, 1996-2016
- Fit results are consistent with null WIMP contribution
- 90 % upper limits on SI WIMP-nucleon scattering cross section  $\sigma_{\chi\text{-}n}$


SK preliminary





# Solar WIMP search

- DM particles passing through the Sun can **elastically scatter with nuclei** and lose energy
- WIMP density increases in core, leading to DM annihilation until equilibrium is achieved:  
***capture rate = annihilation rate***



- Scattering cross section  $\sigma_{\chi n}$  can be constrained and compared with results from direct DM detection







more: G.Wikström, J.Edsjö JCAP  
04, 009 (2009)

Published analysis: K.Choi et al.,  
Phys. Rev. Lett. 114, 141301 (2015)

# Solar WIMP search

example for: 200 GeV WIMPs,  $\tau^+\tau^-$  ann. channel

- FIT based on lepton mom. &  $\cos\theta_{\text{SUN}}$  distributions, 3903 days of SK data (1996-2012)
- No excess of  $\nu$ 's from the SUN as compared to atm bkg
- 90% CL upper limit on WIMP-nucleon scattering cross section  $\sigma_{\text{Xn}}$  for  $\tau^+\tau^-$ ,  $b\bar{b}$  and  $W^+W^-$  channels



+ DATA  
SK1-4, 1996-2012

— ATM MC



■ WIMP  
before fit

# Solar WIMP search: WIMP-nucleon SI & SD cross section limit

90% CL upper limit


spin dependent interactions

spin independent interactions




published: K.Choi et al., Phys. Rev. Lett. 114, 141301 (2015)


# 'Boosted' dark matter search




arXiv:1405.7370



Cone search: 8 cones from  $5^\circ$  to  $40^\circ$  around GC  
 $\rightarrow$  no clusters visible



limit for  $m_\nu = 20$  MeV



# Summary

- DM induced neutrinos has not been observed at Super-Kamiokande so far
- Galactic WIMP search (2017)
  - upper limits on  $\langle\sigma_A V\rangle$  for wide range of WIMPs masses (1 GeV to 10 TeV)
  - strongest limits  $< 20\text{-}100\text{GeV}$  among  $\nu$  experiments
- Earth WIMP search (2017)
  - upper limits on spin-independent WIMP-nucleon cross-section
  - high sensitivity to resonant capture region  $\rightarrow$  currently the strongest limits from  $\nu$  experiments  $< 100\text{ GeV}$
- Solar WIMP search (2015)
  - strongest limits  $< 20\text{-}100\text{GeV}$  among  $\nu$  experiments
- Boosted dark matter search
  - alternative DM models can also be tested with SK detector

# Thank you!



... we keep looking

supplementary  
slides

# Super-Kamiokande Collaboration






# Super-K data samples

## Fully-contained


### FC



- »  $\nu$  energy reconstruction
- »  $\nu$  direction info
- »  $e/\mu$  identification possible

## Partially-contained


### PC



- » partial  $E_\nu$  info (lepton leaves detector)
- »  $\nu$  direction info

## Upward-going muons

### UPMU




- » no  $E_\nu$  info
- » excellent  $\nu$  direction info
- » downward-going muons are neglected (mainly cosmic ray  $\mu$ )


# Galactic WIMP search: ON-/OFF-source

**Different approach:** search for large-scale anisotropy due to DM-induced  $\nu$ 's from Milky Way

$$\Delta N \approx N_{on}^{sig} - N_{off}^{sig} = \Delta N^{sig} \propto \langle \sigma_{AV} \rangle$$



expectation for DM-induced neutrinos




- Analysis uses ON-/OFF-source concept to estimate background directly from data
- Independent on MC simulations and related systematic uncertainties

# ON- & OFF-source results





$$A = \frac{N_{\text{ON}} - N_{\text{OFF}}}{N_{\text{ON}} + N_{\text{OFF}}}$$

SK 1-4, 1996-2016


*UpThruShower\_mu*  
*UpThruNonShower\_mu*  
*UpStop\_mu*  
*PCThru*  
*PCStop*  
*MultiRingOther*  
*MultiRing\_mulike*  
*MultiRing\_elike\_nuebar*  
*MultiRing\_elike\_nue*  
*MultiGeV\_mulike*  
*MultiGeV\_elike\_nuebar*  
*MultiGeV\_elike\_nue*  
*SubGeV\_pi0like*  
*SubGeV\_mulike\_2dcy*  
*SubGeV\_mulike\_1dcy*  
*SubGeV\_mulike\_0dcy*  
*SubGeV\_SingleRing\_pi0like*  
*SubGeV\_elike\_1dcy*  
*SubGeV\_elike\_0dcy*




# Galactic WIMP search: p-value's



# Galactic WIMP search: signal illustration 10GeV bb-bar




# Galactic WIMP search: signal illustration 100GeV bb-bar




$\cos\theta_{GC}$   
 $M_\chi = 100.0 \text{ GeV}/c$

# Galactic WIMP search: signal illustration 1000GeV bb-bar



# Galactic WIMP search: residuals for 5GeV bb-bar best fit



$\cos \theta_{GC}$

$M_\chi = 5.0 \text{ GeV}/c$

points: data set

red line: only ATM MC (with pulls)    color line: best fitted WIMP + ATM MC (all with pulls)

$\chi^2_{\text{total}} = \chi^2_{\text{data}} + \chi^2_{\text{syst}} \quad 604.0 = 566.9 + 37.0$


$601.6 = 564.9 + 36.7$

$\Delta\chi^2 = 2.4 = 2.0 + 0.4$




# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 3 GeV




# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 6 GeV




# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 10 GeV




# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 25 GeV



# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 50 GeV



# Signal illustration for Earth WIMP search

$\tau^+\tau^-$  ann. channel  
WIMP mass = 1 TeV

