

# Flavour Physics Problem Sheet

Warwick Week Graduate Lectures

Matthew Kenzie

April 2023

## Lecture 1

1. Estimate the ratio of the  $p + p \rightarrow d + \pi^+$  and  $p + n \rightarrow d + \pi^0$  cross sections
  - a) Note the deuteron is a bound state of a proton,  $p$ , and a neutron,  $n$  so think about all the possible isospin configurations of two nucleons.
  - b) Identify which isospin state is that of the deuteron (using the observation that bound states of two protons and two neutrons are not found in nature)
  - c) Now consider the total isospin in the initial and final states of the two reactions above
  - d) Now you can approximate the relative cross section
2. What is the quark content of the ground state spin-0 mesons ( $K^\pm$ ,  $\bar{K}^0$ ,  $\pi^\pm$ ,  $\pi^0$ ,  $\eta$ ,  $\eta'$ ) and the ground state spin-1/2 baryons ( $n$ ,  $p$ ,  $\Sigma^\pm$ ,  $\Sigma^0$ ,  $\Lambda^0$ ,  $\Xi^0$ ,  $\Xi^-$ )
3. What are the equivalent excited spin-1 and spin-3/2 states?
4. Show that the CKM matrix has  $N(N-1)/2$  real parameters and  $(N-1)(N-2)/2$  phases for the case of  $N$  fermion generations.
5. Derive the Wolfenstein parameterisation
  - a) First define expansion parameter  $\lambda := V_{us}$  and derive  $c_{13} \approx 1$
  - b) Next write down the standard parameterisation of the CKM-matrix expressing  $s_{12}$  and  $c_{12}$  in terms of  $\lambda$  (include terms up to  $\mathcal{O}(\lambda^3)$ )
  - c) The experimental data suggested that  $1 \approx V_{ud} > V_{us} > V_{cb} > V_{ub}$ . Apply the ansatz  $V_{cb} := A\lambda^2$  and  $V_{ub} := A\lambda^3(\rho - i\eta)$  and express the CKM matrix in terms of  $\lambda$  up to  $\mathcal{O}(\lambda^3)$
6. Show that CKM matrix is unitary if written in the Wolfenstein parameterisation including terms up to  $\mathcal{O}(\lambda^4)$

7. Why is it that down-type neutral mesons contain the anti-quark species but the up-type species contain the quark? In other words why is the state  $(\bar{s}, d)$  (with an anti- $s$ -quark) labelled as  $K^0$  and the state  $(s, \bar{d})$  (with an  $s$ -quark) labelled as  $\bar{K}^0$  whereas the state  $(c, \bar{u})$  is labelled as  $D^0$  and the state  $(\bar{c}, u)$  labelled as  $\bar{D}^0$ ?

## Lecture 2

- Why is it easier for theorists to make predictions for so-called “inclusive” processes, such as  $b \rightarrow c\bar{c}s$ ? Why is it easier for experimentalists to measure so-called “exclusive” modes in which all final states hadrons are identified, such as  $B^0 \rightarrow D^+D^-$ ? How could an experimentalist make an “inclusive” measurement? How could a theorist make an “exclusive” mode prediction?
- Draw the leading order Feynman diagrams for the following decays
  - $B^+ \rightarrow \tau^+ \nu_\tau$
  - $B^+ \rightarrow \bar{D}^0 \mu^+ \nu_\mu$
  - $B^- \rightarrow D^0 K^-$  (via both the favoured  $V_{cb}$  and the suppressed  $V_{ub}$ )
  - $B_s^0 \rightarrow J/\psi \phi$  and  $\bar{B}_s^0 \rightarrow J/\psi \phi$  (note that the  $J/\psi$  is a  $c\bar{c}$  state and the  $\phi$  is a  $s\bar{s}$  state)
  - $B^0 \rightarrow J/\psi K_S^0$  and  $\bar{B}^0 \rightarrow J/\psi K_S^0$
- Compare the Feynman diagrams for  $\bar{B}^0 \rightarrow D^+ \pi^-$  and  $\bar{B}^0 \rightarrow \pi^+ \pi^-$  and make a naive prediction of their ratio of branching fractions

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^+ \pi^-)}{\mathcal{B}(\bar{B}^0 \rightarrow \pi^+ \pi^-)} = ? \quad (1)$$

- Derive the equations for the time-evolution of a pure flavour state of a neutral meson at time  $t = 0$ 
  - First show that when diagonalising the neutral meson mixing matrix from the flavour basis into the mass basis that the following relations hold

$$\Delta M^2 - \frac{1}{4} \Delta \Gamma^2 = 4|M_{12}|^2 - |\Gamma_{12}|^2 \quad (2)$$

$$\Delta M \cdot \Delta \Gamma = -4\Re(M_{12}\Gamma_{12}^*) \quad (3)$$

$$\frac{q}{p} = -\frac{\Delta M + \frac{i}{2}\Delta \Gamma}{2M_{12} - \Gamma_{12}} \quad (4)$$

b) Solve these for the mass and decay rate difference to give

$$2\Delta M^2 = \sqrt{(4|M_{12}|^2 - |\Gamma_{12}|^2)^2 + 16|M_{12}|^2|\Gamma_{12}|^2 \cos^2 \phi_{12} + 4|M_{12}|^2 - |\Gamma_{12}|^2} \quad (5)$$

$$\frac{1}{2}\Delta\Gamma^2 = \sqrt{(4|M_{12}|^2 - |\Gamma_{12}|^2)^2 + 16|M_{12}|^2|\Gamma_{12}|^2 \cos^2 \phi_{12} - 4|M_{12}|^2 + |\Gamma_{12}|^2} \quad (6)$$

with  $\phi_{12} = \arg(-M_{12}/\Gamma_{12})$

c) Now use the time-dependent Schrodinger equation to show how the flavour states evolve with time, *i.e.* that

$$\begin{aligned} |B^0(t)\rangle &= g_+(t)|B^0\rangle + \frac{q}{p}g_-(t)|\bar{B}^0\rangle \\ |\bar{B}^0(t)\rangle &= g_+(t)|\bar{B}^0\rangle + \frac{p}{q}g_-(t)|B^0\rangle \end{aligned} \quad (7)$$

with

$$\begin{aligned} g_+(t) &= e^{-iMt}e^{-\Gamma t/2} \left[ \cosh\left(\frac{\Delta\Gamma t}{4}\right) \cos\left(\frac{\Delta mt}{2}\right) - i \sinh\left(\frac{\Delta\Gamma t}{4}\right) \sin\left(\frac{\Delta mt}{2}\right) \right] \\ g_-(t) &= e^{-iMt}e^{-\Gamma t/2} \left[ -\sinh\left(\frac{\Delta\Gamma t}{4}\right) \cos\left(\frac{\Delta mt}{2}\right) + i \cosh\left(\frac{\Delta\Gamma t}{4}\right) \sin\left(\frac{\Delta mt}{2}\right) \right] \end{aligned} \quad (8)$$

where  $M = (M_L + M_H)/2$  and  $\Gamma = (\Gamma_L + \Gamma_H)/2$ .

d) An alternative (perhaps easier) approach is to write the flavour states in terms of the mass states

$$|B^0\rangle = \frac{1}{2p} (|B_H^0\rangle + |B_L^0\rangle) \quad (9)$$

$$|\bar{B}^0\rangle = \frac{1}{2q} (|B_H^0\rangle - |B_L^0\rangle) \quad (10)$$

and then propagate the flavour states in time given the straightforward propagation of the mass states

$$|B_H^0(t)\rangle = e^{-iM_H t} e^{-\Gamma_H t/2} |B_H^0\rangle \quad (11)$$

$$|B_L^0(t)\rangle = e^{-iM_L t} e^{-\Gamma_L t/2} |B_L^0\rangle \quad (12)$$

5. Show that the probability to oscillate (+) or not (-) is given by

$$|g_{\pm}(t)|^2 = \frac{e^{-\Gamma t}}{2} \left[ \cosh\left(\frac{\Delta\Gamma t}{2}\right) \pm \cos(\Delta mt) \right] \quad (13)$$

6. Write down an expression for the oscillation length of a neutral meson and estimate its value for  $K^0$ ,  $D^0$ ,  $B^0$ ,  $B_s^0$  at LHCb

7. Explain the experimental vs theoretical mixing plot
8. Derive the equation for the time-integrated  $CP$  asymmetry for two amplitudes with different strong and weak phases
9. Derive the “master equations” for neutral meson decay rates