

Flavour Physics (of quarks)

Part 3: Measuring the CKM parameters

Matthew Kenzie

Warwick Week Graduate Lectures

April 2022

Overview

Lecture 1: Flavour in the SM

- ▶ Flavour in the SM
- ▶ Quark Model History
- ▶ The CKM matrix

Lecture 2: Mixing and CP violation

- ▶ Neutral Meson Mixing (no CPV)
- ▶ B -meson production and experiments
- ▶ CP violation

Lecture 3: Measuring the CKM parameters (Today)

- ▶ Measuring CKM elements and phases
- ▶ Global CKM fits
- ▶ CPT and T -reversal
- ▶ Dipole moments

Lecture 4: Flavour Changing Neutral Currents

- ▶ Effective Theories
- ▶ New Physics in B mixing
- ▶ New Physics in rare $b \rightarrow s$ processes
- ▶ Lepton Flavour Violation

Checkpoint Reached

1. Recap

Recap

- ▶ Last time we discussed neutral meson mixing and all three types of CPV
- ▶ Saw the “master” equations for neutral meson decays which are characterised by

$$\lambda_f = \frac{q}{p} \frac{\bar{A}_f}{A_f}$$

- ▶ CPV in decay (the only type possible for a charged initial state) $[\lvert \bar{A}_{\bar{f}}/A_f \rvert \neq 0]$
- ▶ CPV in mixing $[\lvert q/p \rvert \neq 1]$
- ▶ CPV in the interference between mixing and decay $[\arg(\lambda_f) \neq 0]$
- ▶ We got two important expressions which we will see again today
 1. The direct (time-integrated) CP asymmetry arising when we have two amplitudes with different strong (δ) and weak (ϕ) phases and magnitude ratio (r):

$$\mathcal{A}_{CP} = \frac{2r \sin(\delta) \sin(\phi)}{1 + r^2 + 2r \cos(\delta) \cos(\phi)} \quad (1)$$

2. The general time-dependent CP asymmetry for a neutral meson to a CP -eigenstate

$$\mathcal{A}_{CP}(t) = \frac{C_f \cos(\Delta mt) - S_f \sin(\Delta mt)}{\cosh(\frac{1}{2}\Delta\Gamma t) + D_f \sinh(\frac{1}{2}\Delta\Gamma t)} \quad (2)$$

where $C_f = \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}$, $D_f = \frac{2\mathcal{R}e(\lambda_f)}{1 + |\lambda_f|^2}$, $S_f = \frac{2\mathcal{I}m(\lambda_f)}{1 + |\lambda_f|^2}$ (3)

Recap

- ▶ Recall the CKM matrix which governs quark weak transitions

CKM exhibits a clear hierarchy

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \sim \begin{pmatrix} 1 & 0.2 & 0.004 \\ 0.2 & 1 & 0.04 \\ 0.008 & 0.04 & 1 \end{pmatrix}$$

experimentally
determined values

Commonly represented in the Wolfenstein parametrisation

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$

4 $\mathcal{O}(1)$ real parameters (A, λ, ρ, η)

Recap

- ▶ Wolfenstein parameterisation ensures that

$$\bar{\rho} + i\bar{\eta} = -(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*) \quad (4)$$

is phase convention independent and CKM matrix written in $(A, \lambda, \bar{\rho}, \bar{\eta})$ is unitary to all orders in λ

$$\bar{\rho} = \rho(1 - \lambda^2/2 + \dots) \quad \text{and} \quad \bar{\eta} = \eta(1 - \lambda^2/2 + \dots) \quad (5)$$


- ▶ The amount of CP violation in the SM is equivalent to asking how big is η relative to ρ .
- ▶ There are many experimental observables (9 element magnitudes and 4 phases) we can measure to over-constrain the CKM picture.

CKM Unitarity Triangles

- Unitarity gives 6 constraints for off-diagonals represented as triangles in $(\bar{\rho}, \bar{\eta})$ space
 - See [CKMfitter](#) or [UTfit](#) for the latest updates [1, 2]


The (B^0) Unitarity Triangle

$$\bar{\rho}_{(db)} + i\bar{\eta}_{(db)} = -(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)$$



The B_s^0 Unitarity Triangle

$$\bar{\rho}_{sb} + i\bar{\eta}_{sb} = -(V_{us}V_{ub}^*)/(V_{cs}V_{cb}^*)$$




CKM Unitarity Triangles

- Unitarity gives 6 constraints for off-diagonals represented as triangles in $(\bar{\rho}, \bar{\eta})$ space
 - See [CKMfitter](#) or [UTfit](#) for the latest updates [1, 2]


The K Unitarity Triangle

$$\bar{\rho}_{(ds)} + i\bar{\eta}_{(ds)} = -(V_{ud}V_{us}^*)/(V_{cd}V_{cs}^*)$$



The tc Unitarity Triangle

$$\bar{\rho}_{tc} + i\bar{\eta}_{tc} = -(V_{td}V_{cd}^*)/(V_{ts}V_{cs}^*)$$




CKM Unitarity Triangles

- Unitarity gives 6 constraints for off-diagonals represented as triangles in $(\bar{\rho}, \bar{\eta})$ space
 - See [CKMfitter](#) or [UTfit](#) for the latest updates [1, 2]


The $t u$ Unitarity Triangle

$$\bar{\rho}_{(tu)} + i\bar{\eta}_{(tu)} = -(V_{td}V_{ud}^*)/(V_{ts}V_{us}^*)$$



The D Unitarity Triangle

$$\bar{\rho}_{cu} + i\bar{\eta}_{cu} = -(V_{cd}V_{ud}^*)/(V_{cs}V_{us}^*)$$





2. Measuring CKM matrix element magnitudes

Measuring CKM matrix elements

Measuring V_{ud}

- ▶ Compare rates of neutron, n^0 , and muon, μ^- , decays
(see [PDG write-up](#)) [3]
- ▶ The ratio is proportional to $|V_{ud}|^2$
- ▶ $|V_{ud}| = 0.947417 \pm 0.000021$
- ▶ $|V_{ud}| \approx 1$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$




$$\frac{d\Gamma(n \rightarrow pe^- \bar{\nu}_e)}{dx_p} = \frac{G_F^2 m_n^2}{192\pi^2} |V_{ud}|^2 f(q^2)^2 \left(x_p^2 - 4 \frac{m_p^2}{m_n^2} \right)^{3/2}, \quad \text{where } x_p = \frac{2E_p}{m_n}$$

Measuring CKM matrix elements

Measuring V_{us}

- ▶ Compare rates of kaon, K^- , and muon, μ^- , decays (see [PDG write-up](#)) [4]
- ▶ The ratio is proportional to $|V_{us}|^2$
- ▶ $|V_{us}| = 0.2248 \pm 0.0006$
- ▶ $|V_{us}| \approx \sin(\theta_C) \approx \lambda$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$




$$\frac{d\Gamma(\bar{K}^0 \rightarrow \pi^+ e^- \bar{\nu}_e)}{dx_\pi} = \frac{G_F^2 m_K^2}{192\pi^2} |V_{us}|^2 f(q^2)^2 \left(x_\pi^2 - 4 \frac{m_\pi^2}{m_K^2} \right)^{3/2}, \quad \text{where } x_\pi = \frac{2E_\pi}{m_K}$$

Measuring CKM matrix elements

Measuring V_{cd} and V_{cs}

- ▶ Early measurements used neutrino DIS
- ▶ Now use semi-leptonic charm decays, $D^0 \rightarrow \pi^- \ell^+ \nu_\ell$ (V_{cd}) and $D^0 \rightarrow K^- \ell^+ \nu_\ell$ (V_{cs}) (see [PDG write-up](#)) [4]
- ▶ $|V_{cd}| = 0.220 \pm 0.005$
- ▶ $|V_{cs}| = 0.995 \pm 0.016$
- ▶ $|V_{cd}| \approx \sin(\theta_C) \approx \lambda$
- ▶ $|V_{cs}| \approx 1$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$




$$\frac{d\Gamma(\bar{D}^0 \rightarrow \pi^0 e^- \bar{\nu}_e)}{dx_\pi} = \frac{G_F^2 m_D^2}{192\pi^2} |V_{cd}|^2 f(q^2)^2 \left(x_\pi^2 - 4 \frac{m_\pi^2}{m_D^2} \right)^{3/2}, \quad \text{where} \quad x_\pi = \frac{2E_\pi}{m_D}$$

Measuring CKM matrix elements

Measuring V_{cb}

- ▶ Compare rates of $B^0 \rightarrow D^{*-} \ell^+ \bar{\nu}_\ell$ and muon decays (see [PDG write-up](#)) [4]
- ▶ Ratio is proportional to $|V_{cb}|^2$
- ▶ $|V_{cb}| = 0.0405 \pm 0.0013$
- ▶ $|V_{cd}| \approx \sin^2(\theta_C) \approx \lambda^2$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$




$$\frac{d\Gamma(b \rightarrow u_\alpha \ell^- \bar{\nu}_\ell)}{dx} = \frac{G_F^2 m_b^5}{192\pi^2} |V_{\alpha b}|^2 \left(2x^2 \left(\frac{1-x-\xi}{1-x} \right)^2 \left(3 - 2x + \xi + \frac{2\xi}{1-x} \right) \right)$$

$$\text{where } \alpha = u, c, \quad \xi = \frac{m_\alpha^2}{m_b^2}, \quad x = \frac{2E_l}{m_b}$$

Measuring CKM matrix elements

- ▶ The sides of the (B^0) unitarity triangle are constrained by
 - ▶ The ratio V_{ub}/V_{cb} for the left side (known sometimes as R_u)
 - ▶ The ratio $\Delta m_d/\Delta m_s$ for the right side (known sometimes as R_t)
- ▶ Sometimes called “UT constraints from CP -conserving quantities



- ▶ Notice that using the CKM paradigm we can observe CPV without CP -violating quantities

Measurements of V_{ub}

- ▶ There are three ways to determine V_{ub}
 1. "Inclusive" decays of $b \rightarrow u\ell^-\bar{\nu}_\ell$
 - ▶ Of course there are no bare quarks so we are really looking at a sum of exclusive decays of the form $B_{(s)}^{0(-)} \rightarrow \pi^{0(-)}\ell^-\bar{\nu}_\ell X$
 2. "Exclusive" decays e.g. $\bar{B}^0 \rightarrow \pi^+\ell^-\bar{\nu}_\ell$
 3. Leptonic "annihilation" decays e.g. $B^+ \rightarrow \ell^+\nu_\ell$
- ▶ These each come with various requirements on theory (form factors) and the results have historically been rather inconsistent
 - ▶ This is typical in flavour physics
 - ▶ Is the discrepancy a theory issue, an experimental issue or New Physics (or some combination)?

$$\begin{pmatrix} V_{ud} & V_{us} & \textcolor{red}{V_{ub}} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

Inclusive V_{ub}

- ▶ Measure the sum of all processes containing $b \rightarrow u\ell^-\bar{\nu}_\ell$

- ▶ Just think about what this means and how hard this is to achieve


From [HFLAV 2021 write-up \[5\]](#)

- ▶ Experimentally this is incredibly challenging due to backgrounds from the dominant $b \rightarrow c$ semileptonic decays
- ▶ These backgrounds are reduced by either

- ▶ Cutting on the mass of the X_u system or
 - ▶ Cutting on the lepton energy (use the end-point to reject X_c)

- ▶ Essential to have a hermetic detector (need to resolve the neutral) so can only be done at Belle and BaBar

- ▶ It is the mass or end-point cuts which then introduce large theory uncertainties
 - ▶ Need to estimate how much of the X_u phase space is being removed by these cuts




Exclusive V_{ub}

- ▶ Determined by fitting the decay rate seen by BaBar and Belle in e.g. $B^0 \rightarrow \pi^- \ell^+ \nu_\ell$

$$\frac{d\Gamma}{dq^2} = |V_{ub}|^2 \frac{G_F^2}{192\pi^3 m_B^3} \lambda(m_B, m_\pi, q^2)^{3/2} |f_+(q^2)|^2$$

- ▶ Much more straightforward experimentally but more challenging for the theory
 - ▶ Have a dependence on form-factors, $f_+(q^2)$, for the $B \rightarrow \pi$ transition
 - ▶ Use Lattice QCD calculations

[arXiv:1909.12524]




Measurements of V_{ub} and V_{cb}

- ▶ LHCb has also pioneered an approach with the $\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu$ decay
- ▶ Take the ratio with $\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu$ to get $|V_{ub}|/|V_{cb}|$
- ▶ Requires the form factor ratio, R_{FF} , from the Lattice

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)} = \frac{|V_{ub}|^2}{|V_{cb}|^2} R_{FF}$$


- ▶ The global average exhibits a considerable tension between inclusive and exclusive



Measurements of V_{td} and V_{ts}

- There is no top decay but can obtain indirect measurements from the loops which appear in B^0 and B_s^0 mixing
- $|V_{ts}| = 0.0082 \pm 0.0006$
- $|V_{td}| = 0.0400 \pm 0.0027$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ \textcolor{red}{V_{td}} & \textcolor{red}{V_{ts}} & V_{tb} \end{pmatrix}$$




- Ratio of frequencies for B^0 and B_s^0 :

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{B_s^0}}{m_{B^0}} \frac{f_{B_s^0}^2}{f_{B^0}^2} \frac{B_{B_s^0}^2}{B_{B^0}^2} \frac{|V_{ts}|^2}{|V_{td}|^2} = \frac{m_{B_s^0}}{m_{B^0}} \xi^2 \frac{|V_{ts}|^2}{|V_{td}|^2} \quad (6)$$


Measurements of the R_t side

- ▶ B^0 and B_s^0 oscillation frequencies (which we use to get constraints on V_{td} and V_{ts}) measured at LEP, Tevatron, B -factories and LHCb
- ▶ Most precise measurements now come from LHCb

Δm_d from $B^0 \rightarrow D^{(*)-} \mu^+ \nu_\mu X$
[arXiv:1604.03475]



Δm_s from $B_s^0 \rightarrow D_s^- \pi^+$
[arXiv:2104.04421]




Measuring CKM matrix elements

Measuring V_{tb}

- ▶ Use single top production at the Tevatron
- ▶ Ratio is proportional to $|V_{tb}|^2$
- ▶ $|V_{tb}| = 1.009 \pm 0.0031$

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \color{red}V_{tb} \end{pmatrix}$$



$$R = \frac{\mathcal{B}(t \rightarrow Wb)}{BR(t \rightarrow Wq)} = \frac{|V_{tb}|^2}{\sum_q |V_{tq}|^2}$$

Measuring CKM matrix elements

- ▶ These measurements have all been for the **magnitudes** of the CKM elements
 - ▶ Developed over a long period of time using several experiments

$$\begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| \\ |V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| & |V_{ts}| & |V_{tb}| \end{pmatrix} = \begin{pmatrix} 1 & \lambda & \lambda^3 \\ \lambda & 1 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix}$$


$$\lambda \approx \sin(\theta_C) = \sin(\theta_{12}) \approx 0.22$$

- ▶ These give no information on the phase(s)
 - ▶ Let's now consider measurements of this imaginary part
 - ▶ To find the imaginary part we need *CPV*

3. Measuring CKM matrix angles

Measuring CKM matrix phases

Amplitude	Rel. magnitude	phase
$b \rightarrow c$	Dominant	0
$b \rightarrow u$	Suppressed	γ
$t \rightarrow d$	Time-dependent	2β
$t \rightarrow s$	Time-dependent	$-2\beta_s$



- ▶ γ in interference between $b \rightarrow u$ and $b \rightarrow c$ transitions
- ▶ β in interference between B^0 mixing and decay
- ▶ $\beta_s \approx \phi_s$ in interference between B_s^0 mixing and decay
- ▶ α arises in the interference between different $b \rightarrow u$ transitions

$$V_{CKM} = \begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| e^{-i\beta} & -|V_{ts}| e^{-i\beta_s} & |V_{tb}| \end{pmatrix} + \mathcal{O}(\lambda^5)$$

CKM angle β

- ▶ Arises in the interference between $B^0 \rightarrow f_{CP}$ and $B^0 \rightarrow \bar{B}^0 \rightarrow f_{CP}$
- ▶ The golden mode is $B^0 \rightarrow J/\psi K_S^0$ because the master equations (see Lecture 2) simplify considerably
 1. For a B^0 we have no (or at least negligible) CPV in mixing


$$\left| \frac{q}{p} \right| \approx 1$$

2. For the $J/\psi K_S^0$ we have a CP -even final state so $f = \bar{f}$ therefore

$$\lambda_f \equiv \frac{q}{p} \frac{\bar{A}_f}{A_f} = \frac{q}{p} \frac{\bar{A}_{\bar{f}}}{A_{\bar{f}}} \equiv \lambda_{\bar{f}}$$

3. The B^0 and \bar{B}^0 amplitudes to f are (almost) identical (can you think what makes them unequal?)

$$|A_f| = |\bar{A}_f|$$



CKM angle β

- ▶ Recall from the master equations (Lecture 2) that

$$C_f = \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, \quad D_f = \frac{2\Re(\lambda_f)}{1 + |\lambda_f|^2}, \quad S_f = \frac{2\Im(\lambda_f)}{1 + |\lambda_f|^2}$$

- ▶ Giving a time-dependent asymmetry of

$$\mathcal{A}_{CP}(t) = \frac{\Gamma_{X^0 \rightarrow f}(t) - \Gamma_{\bar{X}^0 \rightarrow f}(t)}{\Gamma_{X^0 \rightarrow f}(t) + \Gamma_{\bar{X}^0 \rightarrow f}(t)} = \boxed{\frac{C_f \cos(\Delta mt) - S_f \sin(\Delta mt)}{\cosh(\frac{1}{2}\Delta\Gamma t) + D_f \sinh(\frac{1}{2}\Delta\Gamma t)}} \quad (7)$$

- ▶ In the case of $B^0 \rightarrow J/\psi K_S^0$ this hugely simplifies as $|\lambda_f| = 1$ and $\Delta\Gamma = 0$ so that

$$\mathcal{A}_{CP}(t) = -\Im(\lambda_f) \sin(\Delta mt) \quad (8)$$

CKM angle β

- ▶ Looking into more detail at what λ_f is in the case of $B^0 \rightarrow J/\psi K_S^0$

$$\lambda_{J/\psi K_S^0} = \left(\frac{q}{p}\right)_{B^0} \frac{\bar{A}_{J/\psi K_S^0}}{A_{J/\psi K_S^0}} = \left(\frac{q}{p}\right)_{B^0} \frac{\bar{A}_{J/\psi K^0}}{A_{J/\psi K^0}} \left(\frac{p}{q}\right)_{K^0} \quad (9)$$

$$= - \underbrace{\left(\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*}\right)}_{B^0 \text{ mixing}} \underbrace{\left(\frac{V_{cb} V_{cs}^*}{V_{cb}^* V_{cs}}\right)}_{B^0 \rightarrow J/\psi K^0} \underbrace{\left(\frac{V_{cs} V_{cd}^*}{V_{cs}^* V_{cd}}\right)}_{K^0 \text{ mixing}} \quad (10)$$

$$= -e^{-2i\beta} \quad (11)$$

it's a useful exercise to show this using the equations from Lecture 2

- ▶ So that the time-dependent asymmetry is



$$\mathcal{A}_{CP}(t) = \pm \sin(2\beta) \sin(\Delta m t) \quad (12)$$

the \pm is for CP -even (e.g. $J/\psi K_L^0$) or CP -odd (e.g. $J/\psi K_S^0$) final states


- ▶ A theoretically and experimentally clean signature
- ▶ Also has a relatively large branching fraction, $O(10^{-4})$

CKM angle β


[arXiv:0902.1708]



[arXiv:1201.4643]





[arXiv:1709.03944]



CKM angle β

$$\sin(2\beta) \equiv \sin(2\phi_1) \quad \text{HFLAV}$$

Moriond 2018
PRELIMINARY



$$\sin(2\beta) = 0.699 \pm 0.017$$

$$\beta = (22.2 \pm 0.7)^\circ$$

CKM angle β_s

- ▶ The B_s^0 analogue of β (recall the squeezed B_s^0 unitarity triangle)
- ▶ Use $B_s^0 \rightarrow J/\psi\phi$ which is a spectator quark $d \leftrightarrow s$ switch for $B^0 \rightarrow J/\psi K_S^0$
 - ▶ There are four main differences:

	$B^0 \rightarrow J/\psi K_S^0$	$B_s^0 \rightarrow J/\psi\phi$
1. CKM element	V_{td}	V_{ts}
2. $\Delta\Gamma$	~ 0	~ 0.1
3. Final state (spin)	$K^0 : s = 0$	$\phi : s = 1$
4. Final state (K)	K^0 mixing	-

- ▶ Recall from the master equations the time-dependent CP asymmetry

$$\mathcal{A}_{CP}(t) = \frac{\Gamma_{X^0 \rightarrow f}(t) - \Gamma_{\bar{X}^0 \rightarrow f}(t)}{\Gamma_{X^0 \rightarrow f}(t) + \Gamma_{\bar{X}^0 \rightarrow f}(t)} = \frac{2C_f \cos(\Delta mt) - 2S_f \sin(\Delta mt)}{2 \cosh(\frac{1}{2}\Delta\Gamma t) + 2 \sinh(\frac{1}{2}\Delta\Gamma t)} \quad (13)$$

- ▶ We still have one dominant amplitude so $A_f \approx A_{\bar{f}} \implies |\lambda_f| \approx 1 \implies C_f \approx 0$ so

$$\mathcal{A}_{CP}(t) = \frac{-\mathcal{I}m(\lambda_{J/\psi\phi}) \sin(\Delta mt)}{\cosh(\frac{1}{2}\Delta\Gamma t) + \mathcal{R}e(\lambda_{J/\psi\phi}) \sinh(\frac{1}{2}\Delta\Gamma t)} \quad (14)$$

CKM angle β_s

- ▶ Looking into more detail at what λ_f is in the case of $B_s^0 \rightarrow J/\psi\phi$

$$\lambda_{J/\psi\phi} = \left(\frac{q}{p}\right)_{B_s^0} \left(\eta_{J/\psi\phi} \frac{\bar{A}_{J/\psi\phi}}{A_{J/\psi\phi}} \right) \quad (15)$$

$$= (-1)^l \left(\frac{V_{tb}^* V_{ts}}{V_{tb} V_{ts}^*} \right) \left(\frac{V_{cb} V_{cs}^*}{V_{cb}^* V_{cs}} \right) \quad (16)$$

$$= (-1)^l e^{-2i\beta_s} \quad (17)$$

η represents the CP -eigenvalue

- ▶ Because we have two vectors in the final state there are three amplitudes to consider (as opposed to the one amplitude for $B^0 \rightarrow J/\psi K_S^0$)

$A_{||}$ ($\uparrow\uparrow$) $l = 2$

A_{\perp} ($\uparrow\rightarrow$) $l = 1$


A_0 ($\uparrow\downarrow$) $l = 0$

- ▶ Thus the time-dependent asymmetry becomes

$$\mathcal{A}_{CP}(t) = \frac{-\eta \sin(2\beta_s) \sin(\Delta mt)}{\cosh(\frac{1}{2}\Delta\Gamma t) + \eta \cos(2\beta_s) \sinh(\frac{1}{2}\Delta\Gamma t)} \quad (18)$$

at least it does for each polarisation amplitude independently (the interference between the amplitudes is slightly more complicated)

CKM angle β_s



- ▶ But watch out for polluting penguin contributions!

$$\phi_s = -2\beta_s + \delta\phi^{\text{SM}} + \phi^{\text{NP}}$$

(19)


CKM angle α

- ▶ Following a similar logic to that of $B^0 \rightarrow J/\psi K_S^0$ for β one finds that α arises in the time-dependent asymmetry for modes containing a $b \rightarrow u\bar{u}d$ transition
 - ▶ For example $B^0 \rightarrow \pi^+ \pi^-$ or $B^0 \rightarrow \rho^+ \rho^-$
- ▶ Recalling the master equations with $\Delta\Gamma = 0$
- ▶ Nominally we should have $C_f = 0$ and $S_f = \sin(2\alpha)$ to give

$$\mathcal{A}_{CP}(t) = \pm \sin(2\alpha) \sin(\Delta mt) \quad (20)$$

exactly equivalent to the extraction of β

- ▶ However, in this case there is a **non-negligible contribution from penguin decays of $b \rightarrow d\bar{u}u$**
 - ▶ Similar in magnitude to the $b \rightarrow u\bar{u}q$ transition but has a different weak phase
 - ▶ Therefore $C \neq 0$ and $S \neq \pm \sin(2\alpha)$
 - ▶ How do we deal with the penguin contamination?



CKM angle α


- ▶ The contributions from the penguin amplitudes can be accounted for using an “isospin analysis”

- ▶ Relate the amplitudes for isospin partners

$$A^{+-} \text{ for } B^0 \rightarrow \pi^+ \pi^-, \quad A^{+0} \text{ for } B^+ \rightarrow \pi^+ \pi^0, \quad A^{00} \text{ for } B^0 \rightarrow \pi^0 \pi^0, \quad (21)$$

- ▶ There is no penguin contribution to A^{+0} and \bar{A}^{-0} because $\pi^\pm \pi^0$ is a pure isospin-2 state and the QCD-penguin ($\Delta I = 1/2$) only contributes to the isospin-0 final states
- ▶ Obtain isospin triangle relations


$$A^{+0} = \frac{1}{\sqrt{2}} A^{+-} + A^{00}, \quad \text{and} \quad \bar{A}^{-0} = \frac{1}{\sqrt{2}} \bar{A}^{+-} + \bar{A}^{00} \quad (22)$$



CKM angle α

Add in the related $B \rightarrow \rho\rho$ modes

- ▶ These are vectors (not scalars like the π s) so do not have a fixed CP -eigenvalue
- ▶ However it is found that these decays are almost entirely longitudinally polarised (so approximately CP -even)
- ▶ Much easier to reconstruct, have a much higher branching fraction and have much smaller penguin contributions (triangles are flattened) so have better sensitivity and reduced ambiguities





Add the $B \rightarrow \rho\pi$ system

- ▶ Get a pentagonal (rather than triangular) isospin relation
- ▶ The relative amplitudes of $\rho^+\pi^-$, $\rho^-\pi^+$ and $\rho^0\pi^0$ can all be determined from Dalitz analysis of $B^0 \rightarrow \pi^+\pi^-\pi^0$

CKM angle γ

- ▶ γ is the phase between $V_{ub}^* V_{ud}$ and $V_{cb}^* V_{cd}$
 - ▶ Require interference between $b \rightarrow cW$ and $b \rightarrow uW$ to access it
 - ▶ No dependence on CKM elements involving the top
 - ▶ Can be measured using tree level B decays
- ▶ The “textbook” case is $B^\pm \rightarrow \bar{D}^0 K^\pm$:
 - ▶ Transitions themselves have different final states (D^0 and \bar{D}^0)
 - ▶ Interference occurs when D^0 and \bar{D}^0 decay to the same final state f

Reconstruct the D^0/\bar{D}^0 in a final state accessible to both to achieve interference



- ▶ The crucial feature of these (and similar) decays is that the D^0 can be reconstructed in several different final states [all have same weak phase γ]

Measuring γ

Categorise decays sensitive to γ depending on the $D \xrightarrow{\leftrightarrow} f$ final state

Optimal sensitivity is only achieved when combining them all together

► GLW

- CP eigenstates e.g. $D \rightarrow KK$, $D \rightarrow \pi\pi$
- [Phys. Lett. B253 (1991) 483]
- [Phys. Lett. B265 (1991) 172]

► ADS

- CF or DCS decays e.g. $D \rightarrow K\pi$
- [Phys. Rev. D63 (2001) 036005]
- [Phys. Rev. Lett. 78 (1997) 3257]

► BPGGSZ


- 3-body final states e.g. $D \rightarrow K_S^0 \pi\pi$
- [Phys. Rev. D68 (2003) 054018]

► TD (Time-dependent)

- Interference between mixing and decay e.g. $B_s^0 \rightarrow D_s^- K^+$ [phase is $(\gamma - 2\beta_s)$]
- Penguin free measurement of ϕ_s ?


► Dalitz

- Look at 3-body B decays with D^0 or \bar{D}^0 in the final state, e.g. $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
- [Phys. Rev. D79 (2009) 051301]



γ with CP eigenstates (GLW)

- ▶ Use the $B^\pm \rightarrow \bar{D}^0 K^\pm$ case as an example:
 - ▶ Consider only D decays to CP eigenstates, f_{CP}
 - ▶ Favoured: $b \rightarrow c$ with strong phase δ_F and weak phase ϕ_F
 - ▶ Suppressed: $b \rightarrow u$ with strong phase δ_S and weak phase ϕ_S



Subsequent amplitude to final state f_{CP} is:

$$B^- : A_f = |F|e^{i(\delta_F - \phi_F)} + |S|e^{i(\delta_S - \phi_S)} \quad (23)$$

$$B^+ : \bar{A}_f = |F|e^{i(\delta_F + \phi_F)} + |S|e^{i(\delta_S + \phi_S)} \quad (24)$$

because strong phases (δ) don't change sign under CP while weak phases (ϕ) do

γ with CP eigenstates (GLW)

- ▶ Can define the sum and difference of rates with B^+ and B^-

Rate difference and sum

$$|\bar{A}_{\bar{f}}|^2 - |A_f|^2 = 2|F||S| \sin(\delta_F - \delta_S) \sin(\phi_F - \phi_S) \quad (25)$$

$$|\bar{A}_{\bar{f}}|^2 + |A_f|^2 = |F|^2 + |S|^2 + 2|F||S| \cos(\delta_F - \delta_S) \cos(\phi_F - \phi_S) \quad (26)$$

- ▶ Choose $r_B = \frac{|S|}{|F|}$ (so that $r < 1$) and use strong phase difference $\delta_B = \delta_F - \delta_S$
- ▶ γ is the weak phase difference $\phi_F - \phi_S$
- ▶ Subsequently have two **experimental observables** which are

GLW CP asymmetry

$$\mathcal{A}_{CP} = \frac{\pm 2r_B \sin(\delta_B) \sin(\gamma)}{1 + r_B^2 \pm 2r_B \cos(\delta_B) \cos(\gamma)}$$

GLW total rate



$$\mathcal{R}_{CP} = 1 + r_B^2 \pm 2r_B \cos(\delta_B) \cos(\gamma)$$

- ▶ The $+(-)$ sign corresponds to CP -even (-odd) final states
- ▶ Note that r_B and δ_B (ratio and strong phase difference of favoured and suppressed modes) are different for each B decay
- ▶ **The value of γ is shared by all such decays**

GLW observables

$$A_{CP} = \frac{\pm 2r_B(2F^+ + 1) \sin(\delta_B) \sin(\gamma)}{1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma)} \quad (25)$$



$$R_{CP} = 1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma) \quad (26)$$



GLW observables

$$A_{CP} = \frac{\pm 2r_B(2F^+ + 1) \sin(\delta_B) \sin(\gamma)}{1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma)} \quad (25)$$

$$R_{CP} = 1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma) \quad (26)$$





- LHCb has recently extracted GLW observables from partially reconstructed $B^- \rightarrow D^{*0} K^-$ in the same fit - [arXiv:2012.09903](https://arxiv.org/abs/2012.09903)

GLW observables

$$A_{CP} = \frac{\pm 2r_B(2F^+ + 1) \sin(\delta_B) \sin(\gamma)}{1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma)} \quad (25)$$

$$R_{CP} = 1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma) \quad (26)$$




- ▶ LHCb has recently extracted GLW observables from partially reconstructed $B^- \rightarrow D^{*0} K^-$ in the same fit - [\[arXiv:2012.09903\]](https://arxiv.org/abs/2012.09903)
- ▶ Can extend to quasi- CP -eigenstates ($D^0 \rightarrow K K \pi^0$) if fraction of CP content, F^+ , is known

GLW observables

$$A_{CP} = \frac{\pm 2r_B(2F^+ + 1) \sin(\delta_B) \sin(\gamma)}{1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma)} \quad (25)$$

$$R_{CP} = 1 + r_B^2 \pm 2r_B(2F^+ + 1) \cos(\delta_B) \cos(\gamma) \quad (26)$$



- ▶ Multiple (**but very narrow**) solutions
- ▶ Require knowledge of F^+ from charm friends

γ with CF and DCS decays (ADS)

- ▶ A 2-body D decay to final state f accessible to both D^0 and \bar{D}^0 can be
 - ▶ Cabibbo-favoured (CF) - $D^0 \rightarrow \pi^- K^+$
 - ▶ Doubly-Cabibbo-suppressed (DCS) - $\bar{D}^0 \rightarrow \pi^- K^+$
- ▶ Introduces 2 new hadronic parameters:
 - ▶ r_D - ratio of magnitudes for D^0 and \bar{D}^0 decay to f
 - ▶ δ_D - relative phase for D^0 and \bar{D}^0 decay to f
- ▶ Gives a modified asymmetry and rate defintion

ADS asymmetry

$$\mathcal{A}_{ADS} = \frac{2r_D r_B \sin(\delta_B + \delta_D) \sin(\gamma)}{r_D^2 + r_B^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma)}$$

ADS ratio

$$\mathcal{R}_{ADS} = \frac{|\bar{A}_{\bar{f}}|^2 + |A_f|^2}{|\bar{A}_f|^2 + |\bar{A}_{\bar{f}}|^2} = r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma)$$



- ▶ Hadronic parameters r_D and δ_D can be de independently determined (using CLEO data and HFAG averages)

ADS Method

ADS observables

$$A_{ADS} = \frac{2r_B r_D \sin(\delta_B + \delta_D) \sin(\gamma)}{r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma)} \quad (25)$$

$$R_{ADS} = r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma) \quad (26)$$





- ▶ Much harder to extract partially reconstructed observables because of $B_s^0 \rightarrow D^{(*)0} K^+ \pi^-$ backgrounds.

ADS Method

ADS observables

$$A_{ADS} = \frac{2r_B r_D \sin(\delta_B + \delta_D) \sin(\gamma)}{r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma)} \quad (25)$$

$$R_{ADS} = r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma) \quad (26)$$




- ▶ Much harder to extract partially reconstructed observables because of $B_s^0 \rightarrow D^{(*)0} K^+ \pi^-$ backgrounds.
- ▶ Can extend to multibody-DCS-decays ($D^0 \rightarrow K\pi\pi^0$) if dilution from interference, κ_D , is known

ADS Method

ADS observables

$$A_{ADS} = \frac{2r_B r_D \sin(\delta_B + \delta_D) \sin(\gamma)}{r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma)} \quad (25)$$

$$R_{ADS} = r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos(\gamma) \quad (26)$$




- ▶ A single (yet broader) solution
- ▶ Require knowledge of r_D , δ_D , κ_D from charm friends

γ with 3-body self-conjugate states (BPGGSZ)

- ▶ Now get additional sensitivity over the 3-body phase space
- ▶ Idea is to perform a GLW/ADS type analysis across the D decay phase space
- ▶ For example $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ has contributions from
 - ▶ Singly-Cabibbo-suppressed decay $D^0 \rightarrow K_S^0 \rho^0$
 - ▶ Doubly-Cabibbo-suppressed decay $D^0 \rightarrow K^{*+} \pi^-$
 - ▶ Interference between them enhances sensitivity and resolves ambiguities in γ

BPGGSZ observables (partial rate as function of Dalitz position)

$$d\Gamma_{B^\pm}(x) = A_{(\pm, \mp)}^2 + r_B^2 A_{(\mp, \pm)}^2 + 2A_{(\pm, \mp)} A_{(\mp, \pm)} \\ [r_B \cos(\delta_B \pm \gamma) \cos(\delta_{D(\pm, \mp)}) + r_B \sin(\delta_B \pm \gamma) \sin(\delta_{D(\pm, \mp)})] \quad (27)$$




[arXiv:2010.08483]

BPGGSZ Method

BPGGSZ observables (partial rate as function of Dalitz position)

$$d\Gamma_{B^\pm}(\mathbf{x}) = A_{(\pm, \mp)}^2 + r_B^2 A_{(\mp, \pm)}^2 + 2A_{(\pm, \mp)} A_{(\mp, \pm)} \\ \left[\underbrace{r_B \cos(\delta_B \pm \gamma) \cos(\delta_D(\pm, \mp))}_{x_\pm} + \underbrace{r_B \sin(\delta_B \pm \gamma) \sin(\delta_D(\pm, \mp))}_{c_i} \right] \quad (28)$$
$$\underbrace{_{y_\pm} + \underbrace{r_B \sin(\delta_B \pm \gamma) \sin(\delta_D(\pm, \mp))}_{s_i}}$$




$$\sigma_\gamma \sim \frac{1}{r_B}$$

- ▶ $x_\pm + iy_\pm = r_B e^{i(\delta_B \pm \gamma)}$
- ▶ Uncertainty on γ is inversely proportional to central value of hadronic unknown!!
- ▶ Fluctuation in nuisance parameter = fluctuation in error on parameter of interest!

Model-independent BPGGSZ Analysis

- ▶ Consider both $D \rightarrow K_S^0 \pi\pi$ and $D \rightarrow K_S^0 KK$ decays
- ▶ Divide up the Dalitz space into $2N$ symmetric bins chosen to optimise sensitivity to γ

[arXiv:2010.08483]



Decay amplitude is a superposition of suppressed and favoured contributions

$$A_B(m_-^2, m_+^2) \propto A_D(m_-^2, m_+^2) + r_B e^{i(\delta_B - \gamma)} A_{\bar{D}}(m_-^2, m_+^2)$$

Expected number of B^+ (B^-) events in bin i

$$N_{\pm i}^+ = h_{B^+} \left[F_{\mp i} + (x_+^2 + y_+^2) F_{\pm i} + 2\sqrt{F_i F_{-i}} (x_+ c_{\pm i} - y_+ s_{\pm i}) \right]$$


$$N_{\pm i}^- = h_{B^-} \left[F_{\pm i} + (x_-^2 + y_-^2) F_{\mp i} + 2\sqrt{F_i F_{-i}} (x_- c_{\pm i} - y_- s_{\pm i}) \right]$$

- ▶ $N_{\pm i}^\pm$ - events in each bin
- ▶ c_i, s_i - from CLEO-c (QC $D^0 \bar{D}^0$) measurements
- ▶ $F_{\pm i}$ - from $B \rightarrow D^{*\pm} \mu^\mp \nu_\mu X$
- ▶ h_{B^\pm} - overall normalisation

BPGGSZ Method

Expected number of B^+ (B^-) events in bin i


$$N_{\pm i}^+ = h_{B^+} \left[F_{\mp i} + (x_+^2 + y_+^2) F_{\pm i} + 2\sqrt{F_i F_{-i}} (x_+ c_{\pm i} - y_+ s_{\pm i}) \right]$$
$$N_{\pm i}^- = h_{B^-} \left[F_{\pm i} + (x_-^2 + y_-^2) F_{\mp i} + 2\sqrt{F_i F_{-i}} (x_- c_{\pm i} - y_- s_{\pm i}) \right]$$



- ▶ A single and narrow solution
- ▶ Require knowledge of $c_{\pm i}$ and $s_{\pm i}$ from charm friends


A comment on BPGGSZ systematics

- ▶ Sensitivity to γ starts to degrade due to dependence on input from charm sector
- ▶ Measurements from BES-III (Beijing) will be vital to achieve ultimate precision on γ



The time-dependent method with $B_s^0 \rightarrow D_s^\mp K^\pm$

- ▶ B_s^0 and \bar{B}_s^0 can both decay to same final state $D_s^\mp K^\pm$ (one via $b \rightarrow cW$, the other via $b \rightarrow uW$)
- ▶ Interference achieved by neutral B_s^0 mixing (requires knowledge of $-2\beta_s \equiv \phi_s$)
 - ▶ Weak phase difference is $(\gamma - 2\beta_s)$



- ▶ Requires tagging the initial B_s^0 flavour
- ▶ Requires a time-dependent analysis to observe the meson oscillations
- ▶ Fit the decay-time-dependent decay rates
- ▶ Also requires knowledge of Γ_s , $\Delta\Gamma_s$, Δm_s

The time-dependent method with $B_s^0 \rightarrow D_s^\mp K^\pm$

- Recall the master equations (and equivalents for CP conjugate final state \bar{f})

Time-dependent decay rate for initial B_s^0 or \bar{B}_s^0 at $t = 0$

$$\frac{d\Gamma_{B_s^0 \rightarrow f}(t)}{dt} \propto e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + D_f \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) + C_f \cos(\Delta m_s t) - S_f \sin(\Delta m_s t) \right]$$



$$\frac{d\Gamma_{\bar{B}_s^0 \rightarrow f}(t)}{dt} \propto e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + D_f \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - C_f \cos(\Delta m_s t) + S_f \sin(\Delta m_s t) \right]$$

Time-dependent rate asymmetry

$$\mathcal{A}_{CP}(t) = \frac{\Gamma_{\bar{B}_s^0 \rightarrow f}(t) - \Gamma_{B_s^0 \rightarrow f}(t)}{\Gamma_{\bar{B}_s^0 \rightarrow f}(t) + \Gamma_{B_s^0 \rightarrow f}(t)} = \frac{S_f \sin(\Delta m_s t) - C_f \cos(\Delta m_s t)}{\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + D_f \sinh\left(\frac{\Delta\Gamma_s t}{2}\right)}$$

The time-dependent method with $B_s^0 \rightarrow D_s^\mp K^\pm$

- Fit for decay-time-dependent asymmetry




Variable definitions

$$C_f = -C_{\bar{f}} = \frac{1 - r_B^2}{1 + r_B^2}$$




$$D_{f(\bar{f})} = \frac{-2r_B \cos(\gamma - 2\beta_s \mp \delta_B)}{1 + r_B^2}$$

$$S_{f(\bar{f})} = \frac{\pm 2r_B \sin(\gamma - 2\beta_s \mp \delta_B)}{1 + r_B^2}$$



Dalitz methods


- ▶ Study Dalitz structure of 3-body *B* decays with $B^0 \rightarrow DK^+\pi^-$
 - ▶ In principle has excellent sensitivity to γ
 - ▶ “GW method”? (Gershon-Williams - [\[arXiv:0909.1495\]](#))
- ▶ Get multiple interfering resonances which increase sensitivity to γ
 - ▶ $D^*_0(2400)^-, D^*_2(2460)^-, K^*(892)^0, K^*(1410)^0, K^*_{-2}(1430)^0$
- ▶ Fit *B* decay Dalitz Plot for cartesian parameters (similar to BPGGSZ except for the *B* not the *D*)
 - ▶ $D \rightarrow K^+K^-, D \rightarrow \pi^+\pi^-$ - GLW-Dalitz (done by LHCb - [\[arXiv:1602.03455\]](#))
 - ▶ $D \rightarrow K^\pm\pi^\mp$ - ADS-Dalitz (problematic backgrounds from $B_s^0 \rightarrow DK^\pm\pi^\mp$)
 - ▶ $D \rightarrow K_S^0\pi^+\pi^-$ - BPGGSZ-Dalitz!)



Building up sensitivity

Different B decays


$$\begin{array}{cccc} B^\pm \rightarrow DK^\pm & B^\pm \rightarrow D^*K^\pm & B^\pm \rightarrow DK^{*\pm} & B^0 \rightarrow DK^{*0} \\ r_B^{DK}, \delta_B^{DK} & r_B^{D^*K}, \delta_B^{D^*K} & r_B^{DK^*}, \delta_B^{DK^*} & r_B^{DK^{*0}}, \delta_B^{DK^{*0}} \end{array}$$



Building up sensitivity

Different B decays

$$B^\pm \rightarrow DK^\pm \quad r_B^{DK}, \delta_B^{DK}$$
$$B^\pm \rightarrow D^*K^\pm \quad r_B^{D^*K}, \delta_B^{D^*K}$$
$$B^\pm \rightarrow DK^{*\pm} \quad r_B^{DK^*}, \delta_B^{DK^*}$$
$$B^0 \rightarrow DK^{*0} \quad r_B^{DK^{*0}}, \delta_B^{DK^{*0}}$$



$$D \rightarrow hh$$

$$D \rightarrow hh\pi^0 \quad F^+$$

$$D \rightarrow hhh \quad F^+$$

$$D \rightarrow hh' \quad r_D, \delta_D$$

$$D \rightarrow hh'\pi^0 \quad r_D, \delta_D, \kappa_D$$

$$D \rightarrow hh'h \quad r_D, \delta_D, \kappa_D$$

$$D \rightarrow K_S hh \quad c_i, s_i$$

Different D decays

MANY NUISANCE PARAMETERS

LHCb Input Status

		Highest Statistics	Poorer sensitivity		High potential (Dalitz structure of B)		Low stats (multibody B)	
Method	B Decay D Decay	$B^- \rightarrow D^0 K^-$	$B^- \rightarrow D^0 K^{*-} [K^{*-} \rightarrow K_s^0 \pi^-]$	$B^- \rightarrow D^{*0} K^- [D^{*0} \rightarrow D^0 \pi^0], [D^{*0} \rightarrow D^0 \gamma]$	$B^0 \rightarrow D^0 K^+ \pi^-$	$B^- \rightarrow D^0 K^- \pi^+ \pi^-$		
		part-rec	full-rec	K^{*0} res	Dalitz			
GLW	(+)	$D^0 \rightarrow K^+ K^-$	5 fb^{-1}	5 fb^{-1}	5 fb^{-1}	•	$3 \text{ fb}^{-1}(\bullet)$	3 fb^{-1}
		$D^0 \rightarrow \pi^+ \pi^-$	5 fb^{-1}	5 fb^{-1}	5 fb^{-1}	•	$3 \text{ fb}^{-1}(\bullet)$	3 fb^{-1}
		$D^0 \rightarrow K^+ K^- \pi^0$	$3 \text{ fb}^{-1}(\bullet)$	-	-	-	-	-
		$D^0 \rightarrow \pi^+ \pi^- \pi^0$	3 fb^{-1}	-	-	-	-	-
		$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$	•	-	-	-	-	-
		$D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	$3 \text{ fb}^{-1}(\bullet)$	5 fb^{-1}	•	•	-	-
ADS	(-)	$D^0 \rightarrow K_s^0 \pi^0$	•	-	-	-	-	-
		$D^0 \rightarrow K^+ \pi^-$	$3 \text{ fb}^{-1}(\bullet)$	5 fb^{-1}	•	•	$3 \text{ fb}^{-1}(\bullet)$	•
		$D^0 \rightarrow K^+ \pi^- \pi^0$	3 fb^{-1}	-	-	-	-	-
GGSZ		$D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$	$3 \text{ fb}^{-1}(\bullet)$	5 fb^{-1}	•	•	-	-
		$D^0 \rightarrow K_s^0 \pi^+ \pi^-$	5 fb^{-1}	•	-	•	$3 \text{ fb}^{-1}(\bullet)$	•
		$D^0 \rightarrow K_s^0 K^+ K^-$	5 fb^{-1}	•	-	•	$3 \text{ fb}^{-1}(\bullet)$	•
		$D^0 \rightarrow K_s^0 \pi^+ \pi^- \pi^0$	•	-	-	-	-	-
		$D^0 \rightarrow K_s^0 K^+ K^- \pi^0$	•	-	-	-	-	-

KEY: •: (update) in progress

•: requires input from Charm sector (r_D, δ_D, κ_D)

NOTE: TD result with $B_s^0 \rightarrow D_s^- K^+ 3 \text{ fb}^{-1}(\bullet)$

TD result with $B^0 \rightarrow D^- \pi^+ 3 \text{ fb}^{-1}$

GLS result from $B^- \rightarrow D^0 K^-$ with $D^0 \rightarrow K_s^0 K^\pm \pi^\mp 3 \text{ fb}^{-1}(\bullet)$

Working on $B^- \rightarrow D^0 K^{*-}$ with $K^{*-} \rightarrow K^- \pi^0$ •

LHCb Input Status

		Highest Statistics	Poorer sensitivity		High potential (Dalitz structure of B)	Low stats (multibody B)
Method		B Decay	D Decay			
GLW	(+) D	$B^- \rightarrow D^0 K^-$				
			$B^- \rightarrow D^0 K^{*-}$ [$K^{*-} \rightarrow K_S^0 \pi^-$]	$B^- \rightarrow D^{*0} K^-$ [$D^{*0} \rightarrow D^0 \pi^0$], [$D^{*0} \rightarrow D^0 \gamma$]	$B^0 \rightarrow D^0 K^+ \pi^-$	$B^- \rightarrow D^0 K^- \pi^+ \pi^-$
		$D^0 \rightarrow K^+ K^-$	GLW	5 fb ⁻¹		
		$D^0 \rightarrow \pi^+ \pi^-$		5 fb ⁻¹		
		$D^0 \rightarrow K^+ K^- \pi^0$		3 fb ⁻¹ (●)		
	(-) D	$D^0 \rightarrow \pi^+ \pi^- \pi^0$		3 fb ⁻¹		
		$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$		•		
		$D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$		3 fb ⁻¹ (●)		
		$D^0 \rightarrow K_S^0 \pi^0$		5 fb ⁻¹		
ADS	D	$D^0 \rightarrow K^+ \pi^-$	GGSZ	3 fb ⁻¹ (●)	5 fb ⁻¹	
		$D^0 \rightarrow K^+ \pi^- \pi^0$			•	
		$D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$		•		
	S	$D^0 \rightarrow K_S^0 \pi^+ \pi^-$		5 fb ⁻¹		
GGSZ	S	$D^0 \rightarrow K_S^0 K^+ K^-$		5 fb ⁻¹		
		$D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$		•		
		$D^0 \rightarrow K_S^0 K^+ K^- \pi^0$		•		
		$D^0 \rightarrow K_S^0 K^+ K^- \pi^0$		-		

KEY: ●: (update) in progress


●: requires input from Charm sector (r_D, δ_D, κ_D)

NOTE: TD result with $B_s^0 \rightarrow D_s^- K^+ 3 \text{ fb}^{-1}$ (●)

TD result with $B^0 \rightarrow D^- \pi^+ 3 \text{ fb}^{-1}$

GLS result from $B^- \rightarrow D^0 K^-$ with $D^0 \rightarrow K_S^0 K^\pm \pi^\mp 3 \text{ fb}^{-1}$ (●)



Working on $B^- \rightarrow D^0 K^{*-}$ with $K^{*-} \rightarrow K^- \pi^0$ ●



Combined constraints on γ

World Average (HFLAV) - [PDG 2021]

$$\gamma = (66.2^{+3.4}_{-3.6})^\circ$$



Indirect constraints are: $\gamma = (65.3^{+1.0}_{-2.5})^\circ$

4. CKM constraints from kaon decays

CPV in the kaon sector

- ▶ *CPV* first observed in 2π decays of K_L^0 mesons
 - ▶ Is this just mixing induced or is it direct *CPV* also (*i.e.* *CPV* in decay)?
 - ▶ For the *CPV* in kaon mixing we introduce the complex parameter ϵ such that

$$|K_S^0\rangle = \frac{1}{\sqrt{1+|\epsilon|^2}} (|K_1\rangle + \epsilon|K_2\rangle) \quad \text{and} \quad |K_L^0\rangle = \frac{1}{\sqrt{1+|\epsilon|^2}} (|K_2\rangle + \epsilon|K_1\rangle) \quad (29)$$

- ▶ If *CPV* is only mixing induced then we expect $K_L^0 : K_S^0$ amplitude ratios to be equivalent for neutral and charged final states (*i.e.* $\eta_{00} = \eta_{+-}$) where

$$\eta_{00} = \frac{\mathcal{A}(K_L^0 \rightarrow \pi^0 \pi^0)}{\mathcal{A}(K_S^0 \rightarrow \pi^0 \pi^0)}, \quad \eta_{+-} = \frac{\mathcal{A}(K_L^0 \rightarrow \pi^+ \pi^-)}{\mathcal{A}(K_S^0 \rightarrow \pi^+ \pi^-)}. \quad (30)$$

- ▶ But we also see evidence for *CPV* in kaon decay (via semileptonic decays)

$$\delta \equiv \mathcal{A}_{CP}(K_L^0 \rightarrow \ell^+ \nu_\ell \pi^-) \quad (31)$$

- ▶ Can then summarise *CPV* in the kaon system using two parameters, (ϵ, ϵ') where

$$\eta_{00} = \epsilon - 2\epsilon' \quad (32)$$

$$\eta_{11} = \epsilon + \epsilon' \quad (33)$$


$$\delta = \frac{2\mathcal{R}e(\epsilon)}{1+|\epsilon|^2} \quad (34)$$

NA48 experiment

- ▶ Established that $\mathcal{R}e(\epsilon'/\epsilon) \neq 0$ by NA48 at CERN and KTEV in Japan
- ▶ NA48 is a fixed target experiment in CERN's North Area
- ▶ Measure the double ratio of $\pi^0\pi^0$ and $\pi^+\pi^-$ decays from K_L^0 and K_S^0

$$R = \frac{|\eta_{00}|^2}{|\eta_{+-}|^2} \approx 1 - 6\mathcal{R}e\left(\frac{\epsilon'}{\epsilon}\right) \quad (35)$$
$$= (13.7 \pm 2.5 \pm 1.8) \times 10^{-4}$$

- ▶ Now replaced by NA62 an even more sensitive kaon physics experiment looking for very rare kaon decays



5. Status of CKM matrix global fits

Putting all the constraints together


- ▶ All of these separate measurements can be put together to over-constrain the CKM picture
- ▶ This is incredibly powerful because we can attack the (ρ, η) vertex of the unitarity triangle in several ways

World Averages are performed by several groups

- ▶ CKMfitter (frequentist)
 - ▶ <http://ckmfitter.in2p3.fr/>
- ▶ UTfit (Bayesian)
 - ▶ <http://www.utfit.org/UTfit/>
- ▶ Heavy Flavour Averaging Group (HFLAV)
 - ▶ <https://hflav.web.cern.ch/>
- ▶ Particle Data Group (PDG)
 - ▶ <http://pdg.lbl.gov/>

The CKM fit


1995




- ▶ Before the B -factories and LHC the CKM picture was not even established

The CKM fit

1995






2005






- With data from the Tevatron and B -factories the CKM picture is verified
- When adding the LHC it now becomes a suite of precision physics measurements




The CKM fit



The CKM fit





The CKM fit





The CKM fit

Comparison between CP -conserving (lengths of sides) and CP -violating (angles)



Comparison between tree-level (γ, V_{ub}) and loop-level ($\alpha, \beta, \Delta m, \epsilon$)



6. CPT and T-reversal

CPT theorem

- ▶ It is not possible to write a quantum field theory that is Lorentz invariant, with a Hermitian Hamiltonian $H = H^\dagger$, that violates the product of *CPT*
 - ▶ i.e. one in which measurements are not invariant under position translations and Lorentz boosts of the system
- ▶ There are several important consequences that *CPT* invariance implies
 1. Mass and lifetime of particles and antiparticles are identical
 2. Quantum numbers of antiparticles are opposite those of particles
 3. Integer spin particles obey Bose-Einstein statistics and half-integer spin particle obey Fermi-Dirac statistics
- ▶ Time reversal symmetry translates $t \rightarrow -t$
 - ▶ Obviously we can't test this experimentally (cannot run an experiment backwards in time)
 - ▶ However if *CP* is violated and the product *CPT* is conserved then *T* must also be violated

T violation in the B system


- ▶ This can actually be tested in the B system
- ▶ A generalisation of the $\sin(2\beta)$ analysis
- ▶ Identify the flavour of the B by tagging the other B in the event and in addition separate the events by CP -odd ($J/\psi K_S^0$) and CP -even ($J/\psi K_L^0$) final states
- ▶ A T reversal violation would appear as a difference in the rates between

$$\bar{B}^0(t_1) \rightarrow B_-(t_2) \quad \text{and} \quad B_-(t_1) \rightarrow \bar{B}^0(t_2)$$

- ▶ T violation has been observed by BaBar ([\[arXiv:1207.5832\]](https://arxiv.org/abs/1207.5832))

$$\Delta S_T^+ = -1.37 \pm 0.15$$

$$\Delta S_T^- = 1.17 \pm 0.21$$



7. Dipole Moments

Magnetic dipole moments

- ▶ A “spinning” charge acts as a magnetic dipole with moment, μ , which gives an energy shift to an externally applied magnetic field

$$\Delta E = -\vec{\mu} \cdot \vec{B} \quad (36)$$

- ▶ The prediction of $g = 2$ (classically $g = 1$) was a big success of the Dirac equation
- ▶ In an external field A^μ


$$\left(\frac{1}{2m}(\vec{p} + e\vec{A}) + \frac{e}{2m}\vec{\sigma} \cdot \vec{B} - eA^0 \right) \psi = E\psi \quad (37)$$

- ▶ The magnetic dipole moment μ is given by

$$\vec{\mu} = -\frac{e}{2m}\vec{\sigma} = -g\frac{\mu_B}{\hbar}\vec{S} \quad (38)$$


- ▶ Receives corrections from higher order processes (e.g. at order α^2)

$$g = 2 + \frac{\alpha}{2\pi} + \mathcal{O}(\alpha^2)$$



Anomalous magnetic moment

Dirac




Charged,
spin $\frac{1}{2}$
Bparticle


12672 diagrams



Kinoshita




Up to 10th
Order QED



Electroweak


$$g_\mu = 2.002\ 331\ 841\ 78(126)$$

Schwinger



1st Order QED

Dominant uncertainty
in calculation



?

Slide from Becky Chislett (via Tom Blake)

Anaomalous magnetic moments

- ▶ $(g - 2)_e$ is a powerful precision test of QED

$$(g - 2)_e = (1159.652186 \pm 0.000004) \times 10^{-6}$$


- ▶ $(g - 2)_\mu$ receives important Weak and QCD contributions. The latest experimental value from Brookhaven E821 and Fermilab $g - 2$ experiments

$$(g - 2)_\mu = (116591810 \pm 43) \times 10^{-11} \text{ (Theory)}$$

$$(g - 2)_\mu = (116592061 \pm 41) \times 10^{-11} \text{ (Experiment)}$$



from [arXiv:2104.03281] is 4.2σ from the SM expectation [arXiv:2006.04822]

- ▶ Is this a hint of a NP contribution to $(g - 2)_\mu$ (review in [arXiv:0902.3360])?



The $g - 2$ experiment

- ▶ Experiment at Fermilab aiming for $\sim 0.1 - 0.2\text{ppm}$ precision
- ▶ The anomalous magnetic moment causes the spin to precess at a different rate to the momentum vector
- ▶ Can use this precession to precisely measure $g - 2$




Electric dipole moments

- ▶ Classically, EDMs are a measure of the spatial separation of positive and negative charges in a particle
 - ▶ A finite EDM can only exist if the charge centres do not coincide
- ▶ EDMs can also be measured for fundamental particles (electron, muon, neutron etc.)
 - ▶ Can interpret this as a measure of the “sphericity” of the particle
- ▶ This is tested using the Zeeman effect
 - ▶ Look for a shift in energy levels under an external electrical field (analogous to the magnetic moment)

$$\Delta E = -\vec{d} \cdot \vec{E} \quad (39)$$

- ▶ A non zero EDM would violate T and P symmetries
 - ▶ Under T reversal, the MDM would change direction but the EDM would remain unchanged
 - ▶ Under P , the EDM would change direction but the MDM remains unchanged
- ▶ Violation of P and T implies CP violation



Electric dipole moments

- ▶ Electron EDM:
 - ▶ $d_e < 8.7 \times 10^{-29}$ [[arXiv:1310.7534](#)]
- ▶ Muon EDM:
 - ▶ $d_e < 1.9 \times 10^{-19}$ [[arXiv:0811.1207](#)]
- ▶ Neutron EDM:
 - ▶ $d_e < 3.0 \times 10^{-26}$ [[arXiv:hep-ex/0602020](#)]
- ▶ Probing incredibly small charge separation distances!

Strong CP problem

- ▶ The complicated nature of the QCD vacuum should give rise to a term in the Lagrangian like

$$\mathcal{L}_\theta = \theta \frac{\alpha_s}{8\pi} F_\alpha^{\mu\nu} \tilde{F}_{\alpha,\mu\nu} \quad (40)$$

- ▶ This is both P and T -violating but C -conserving (hence CP -violating)
- ▶ This terms would also contribute to the neutron dipole moment, but experimentally we know this is very small

$$d_n \sim e \cdot \theta \cdot m_q / M_N^2 \implies \theta \leq 10^{-9} \quad (41)$$

- ▶ This incredibly small size of the θ parameter is (another) massive fine tuning problem (the so-called “[strong \$CP\$ problem](#)”)
- ▶ What mechanism forces θ to be so small?

Axion searches

- ▶ The Peccei-Quin solution to the strong CP problem is to introduce a $U(1)$ symmetry that removes the strong CP problem by dynamically making θ small
- ▶ Spontaneous breaking of this symmetry is associated with a pseudo-Nambu-Goldstone boson (in analogy with the Higgs mechanism), [the axion](#)
- ▶ The axion can be a light particle that couples very weakly to known SM particles
- ▶ There are a large number of searches for axions produced in particle colliders (direct searches)
- ▶ Can also be detected by the presence of axions converting into photons in the presence of a strong magnetic field (e.g. the CAST experiment at CERN)

8. Recap

Recap

In this lecture we have covered

- ▶ Recap of the CKM matrix and unitarity triangles
- ▶ Measurements of the CKM matrix element magnitudes
 - ▶ In particular the sides of the unitarity triangle
 - ▶ The tension between inclusive and exclusive measurements of V_{ub}
- ▶ Measurements of the CKM matrix angles
 - ▶ The angles α , β , γ and ϕ_s
- ▶ CP violation in the kaon system
- ▶ Global constraints on the CKM matrix and unitarity triangle(s)
- ▶ T violation and CPT
- ▶ Electric and magnetic dipole moments

Checkpoint Reached

End of Lecture 3

References I

- [1] CKMfitter group, J. Charles *et al.*, *Current status of the standard model CKM fit and constraints on $\Delta F = 2$ new physics*, Phys. Rev. D91 (2015) 073007, arXiv:1501.05013, updated results and plots available at <http://ckmfitter.in2p3.fr/>.
- [2] UTfit collaboration, M. Bona *et al.*, *The unitarity triangle fit in the standard model and hadronic parameters from lattice QCD: A reappraisal after the measurements of Δm_s and $BR(B \rightarrow \tau \nu_\tau)$* , JHEP 10 (2006) 081, arXiv:hep-ph/0606167, updated results and plots available at <http://www.utfit.org/>.
- [3] J. C. Hardy and I. S. Towner, *Superallowed $0^+ \rightarrow 0^+$ nuclear β decays: 2020 critical survey, with implications for V_{ud} and CKM unitarity*, Phys. Rev. C 102 (2020) 045501.
- [4] Particle Data Group, R. L. Workman *et al.*, *Review of Particle Physics*, PTEP 2022 (2022) 083C01.
- [5] HFLAV, Y. S. Amhis *et al.*, *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2021*, arXiv:2206.07501.