

High P_T physics at the LHC - Lecture I

(Introduction and LHC accelerator)

Juraj Bracinik, Andrew Chisholm
(University of Birmingham)

Warwick week, 21 March 2024

- ♦ Introduction
- ♦ LHC machine
- ♦ High P_T experiments - Atlas and CMS
- ♦ Standard Model physics and BSM searches
- ♦ Higgs data analysis



Introduction

What are these lectures going to be?

- ♦ Introduction to the topics
- ♦ An overview, the topic is too broad to go into details
- ♦ Different people need to know different details, but an overview can be useful to everyone...
- ♦ Not too much maths, most of it just to give us feeling for orders of magnitude
- ♦ (maybe) a discussion ???!

Four lectures ...






Lecture 1 : Introduction to LHC physics, the LHC accelerator

Lecture 2: General purpose experiments (ATLAS and CMS)

Lecture 3: SM physics and searches

Lecture 4: Higgs physics

Please allow me to introduce myself ...



I work for ATLAS trigger, so naturally biased ...

Lecture 1 - Introduction and the LHC

- ♦ Standard model of elementary particle physics and its Big Open Questions
- ♦ LHC machine:
 - General parameters
 - How are particles accelerated: RF
 - What keeps them running around?
 - Interaction points, that is where it all happens!
- ♦ The future of the LHC and beyond

Standard model and its (standard) troubles


Standard model and standard (model) troubles

Standard Model describes all observed phenomena in Elementary particle physics

- ◆ 2 x 6 fundamental fermions - "particles of matter"
- ◆ 4 fundamental, spin 1 bosons - "particles of interaction"

Language (mathematical) of SM:


- ➔ Local renormalizable Quantum Field Theory




Standard model and its problems

- ▶ Quantum field theory with massive spin-1 boson is not renormalisable
- ▶ W and Z are very massive!
- ▶ Several attempts to solve this problem ...

- ▶ The simplest model:
 - Postulate additional scalar field - Higgs particle
 - It has nonzero vacuum expectation value
 - W and Z acquire mass in interaction with Higgs
 - γ remains mass-less
- ▶ Build LHC to check if this is the case (or not)
- ▶ Looks like this simple model is correct (or close ...) !




Standard model and its problems





More details in Andy's lectures...

The Standard Model of particle physics

Years from concept to discovery



Standard model and its problems





Standard model and its problems

| Parameters of the Standard Model [hide] | | | |
|--|--------------------------------|--|-------------------------------|
| Symbol | Description | Renormalization scheme (point) | Value |
| m_e | Electron mass | | 511 keV |
| m_μ | Muon mass | | 105.7 MeV |
| m_τ | Tau mass | | 1.78 GeV |
| m_u | Up quark mass | μ $\overline{MS} = 2 \text{ GeV}$ | 1.9 MeV |
| m_d | Down quark mass | μ $\overline{MS} = 2 \text{ GeV}$ | 4.4 MeV |
| m_s | Strange quark mass | μ $\overline{MS} = 2 \text{ GeV}$ | 87 MeV |
| m_c | Charm quark mass | μ $\overline{MS} = m_c$ | 1.32 GeV |
| m_b | Bottom quark mass | μ $\overline{MS} = m_b$ | 4.24 GeV |
| m_t | Top quark mass | On shell scheme | 173.5 GeV |
| θ_{12} | CKM 12-mixing angle | | 13.1° |
| θ_{23} | CKM 23-mixing angle | | 2.4° |
| θ_{13} | CKM 13-mixing angle | | 0.2° |
| δ | CKM CP violation Phase | | 0.995 |
| g_1 or g' | U(1) gauge coupling | μ $\overline{MS} = m_Z$ | 0.357 |
| g_2 or g | SU(2) gauge coupling | μ $\overline{MS} = m_Z$ | 0.652 |
| g_3 or g_s | SU(3) gauge coupling | μ $\overline{MS} = m_Z$ | 1.221 |
| θ_{QCD} | QCD vacuum angle | | ~ 0 |
| v | Higgs vacuum expectation value | | 246 GeV |
| m_H | Higgs mass | | $125.09 \pm 0.24 \text{ GeV}$ |

→ Nineteen free parameters that need to be determined from the experiment!


Dark matter

Rotation curve of typical spiral galaxy:




- ◆ We know there is a physics beyond Standard model!
- ◆ Just need to look at the sky!

Dark matter




Basic kinematics of high energy hadron collisions

One accelerator with TeV energy, please...



Hadron colliders have their advantages when it comes to exploration of unknown territory (discovery physics)

Detailed look at high energy pp collision



- ◆ Protons at high energy behave as beams of pointlike particles - partons
- ◆ Proton beam offers wide range of (elementary) collision energies
- ◆ Variable x (Bjorken x) gives fraction of proton energy carried by a parton:



$$\hat{x} = \frac{P_{parton}}{P_{proton}}$$

- ◆ Energy of (elementary) collision is then

$$\sqrt{s_{elementary}} = \hat{x}_1 \hat{x}_2 \sqrt{s} \quad s = (p_1 + p_2)^2$$

Proton colliders offer wide range of available center-of-mass energy for elementary collisions :-)

Structure of the proton I



- ◆ Distribution of partons in the proton is well known!
- ◆ measured (mainly) in Deep Inelastic ep Scattering (DIS)
- ◆ DIS \Leftrightarrow elastic electron-quark scattering!
- ◆ Distribution of scattered electrons is very sensitive to distribution of partons in the proton

Luminosity

Energy is not enough, one needs luminosity, too...


$$N = \sigma \times L$$

N - number of events (we want)

σ - cross section (given by Nature)


L - luminosity (parameter of an accelerator)


- ◆ Higgs couples mainly to particles with high mass, its cross section in pp collisions is rather small
- ◆ Need a machine with high luminosity !!!



Introduction to the LHC









LHC nominal parameters

at collision energy


| | |
|----------------------------------|---|
| Particle type | p, Pb |
| Proton energy E_p at collision | 7000 GeV |
| Peak luminosity (ATLAS, CMS) | $1 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ |
| Circumference C | 26 658.9 m |
| Bending radius ρ | 2804.0 m |
| RF frequency f_{RF} | 400.8 MHz |
| # particles per bunch n_p | 1.15×10^{11} |
| # bunches n_b | 2808 |

Particle accelerators

Accelerator: accelerate and steer particles (and collide them):

$$\vec{F} = q(\vec{E} + \vec{v} \times \vec{B}) = \vec{F}_E + \vec{F}_B$$

- ◆ Both F_E and F_B cause deflection:
- ◆ when $v \sim c$, $1T \sim 3 \times 10^8$ V/m
- ◆ Achievable E field \sim few MV/m
- Magnetic field is used in accelerators when possible (beam steering)




$$\vec{F}_B \perp \vec{v}$$




Only electric field accelerates!

Accelerating particles

Linear (electrostatic) accelerator




Cyclotron






An important function of an accelerator is to accelerate ...

Cyclotron, betatron, synchrotron, oh my ...



- ◆ LHC is a synchrotron!
- ➔ (in fact most of high energy accelerators are synchrotrons, for example HERA, Tevatron, LEP, SPS, ...)
- ◆ Means that particles follow the same (circular) trajectories, steered by magnets
- ◆ When accelerating, changing magnetic field
- ◆ Acceleration done by RF cavities
- ◆ Changing also frequency of accelerating (RF) field

Accelerating particles - RF cavities





- ◆ In any closed metallic box it is possible to generate electromagnetic oscillations
- ◆ For example an ideal cylindrical cavity
- ◆ Many (infinite number) of solutions for E and B - oscillating modes
- ◆ The fundamental mode normally used for acceleration is named TM_{010}
- ◆ E_z is constant in space along the axis of acceleration, z, at any instant

$$\left. \begin{aligned} E_z &= J_0(kr) \\ H_\theta &= -\frac{j}{Z_0} J_1(kr) \end{aligned} \right\} e^{j\omega t}$$

$$k = \frac{2\pi}{\lambda} = \frac{\omega}{c} \quad \lambda = 2,62 a \quad Z_0 = 377 \Omega$$

RF cavities




- ◆ RF power is fed into the cavity from RF power generators (for example Klystrons)
- ◆ RF power oscillates at desired frequency
- ◆ Good to have cavity with superconducting walls to minimize losses...
- ◆ Particles oscillating in the accelerator pass through the cavity many times, to be accelerated, they need to come at fixed phase
 - ➔ Beam is composed of bunches with a large number of particles

Phase stability and bunches I


Assume the situation where energy increase is transferred into a velocity increase

Particles P_1 , P_2 have the synchronous phase.




M_1 & N_1 will move towards P_1 \Rightarrow stable
 M_2 & N_2 will go away from P_2 \Rightarrow unstable (and finally be lost)

Phase stability and bunches II



- ◆ Area of stability in phase space - bunches
- ◆ Energy and phase oscillate around nominal values - synchrotron oscillations
- ◆ For small amplitudes: Harmonic Oscillator
- ◆ Higher amplitudes: non-linearities


Phase stability and bunches III






- ▶ Around 2800 filled bunches per beam
- ▶ Bunch length ~ 10 cm
- ▶ Maximum transversal size (far from experimental collision points):
 - ▶ ~ 1 mm at injection energy
 - ▶ ~ 0.5 mm at full energy

LHC bunch structure - 2016

- 25 ns bunch spacing
- Nominal bunch intensity 1.15×10^{11} protons per bunch




Injection and Extraction



Extraction follows the same principle, but the beam travels in the **opposite direction**.

Accelerating particles - LHC cavities



- ◆ 400 MHz superconducting cavity system
- ◆ 8 single-cell cavities per ring
- ◆ 1 klystron per cavity
- ◆ 4 cells are in one cryostat (4.5° K)

Maximum field 5 MV/m

Energy gain/turn ~ 0.5 MeV

RF frequency varies from 400.789 MHz


(450 GeV) to 400.790 MHz (7 TeV)

Energy ramp

The LHC can accelerate protons and heavier ions.

In the past: runs with p^+ and Pb^{82+}

For the ramp of lead ions larger frequency swing



- ◆ Slow ramp ($> 15min$)
- ◆ Small energy gain/turn ($\sim 500keV$)

Accelerating particles - LHC cavities



Table 4.1: The Main Beam and RF Parameters.

| | Unit | Injection 450 GeV | Collision 7 TeV |
|---|---------------|----------------------|--------------------|
| Bunch area (2σ)* | eVs | 1.0 | 2.5 |
| Bunch length (4σ)* | ns | 1.71 | 1.06 |
| Energy spread (2σ)* | 10^{-3} | 0.88 | 0.22 |
| Intensity per bunch | 10^{11} p | 1.15 | 1.15 |
| Number of bunches | | 2808 | 2808 |
| Normalized rms transverse emittance V/H | μm | 3.75 | 3.75 |
| Intensity per beam | A | 0.582 | 0.582 |
| Synchrotron radiation loss/turn | keV | - | 7 |
| Longitudinal damping time | h | - | 13 |
| Intrabeam scattering growth time - H | h | 38 | 80 |
| - L | h | 30 | 61 |
| Frequency | MHz | 400.789 | 400.790 |
| Harmonic number | | 35640 | 35640 |
| RF voltage/beam | MV | 8 | 16 |
| Energy gain/turn (20 min. ramp) | keV | 485 | |
| RF power supplied during acceleration/ beam | kW | ~ 275 | |
| Synchrotron frequency | Hz | 63.7 | 23.0 |
| Bucket area | eVs | 1.43 | 7.91 |
| RF (400 MHz) component of beam current | A | 0.87 | 1.05 |


* The bunch parameters at 450 GeV are an upper limit for the situation after filamentation, ~ 100 ms after each batch injection. The bunch parameters at injection are described in the text.

Keeping particles on circle - dipoles I

- ◆ Circular accelerator - deflecting forces are needed
 - Usually done with pieces of circular trajectory
 - Straight sections used to accelerate particles (RF) and to collide them (detectors)
 - In circular arc section - bending by magnetic fields
- ◆ Dipole magnets:

$$\frac{1}{\rho} = \frac{eB}{p}$$

$$\frac{1}{\rho} [m^{-1}] = 0.3 \frac{B [T]}{p [GeV/c]}$$



Keeping particles on circle - dipoles II

Assuming:

$$B = 8.3 T$$

$$p = 7000 \frac{\text{GeV}}{c}$$

$$\frac{1}{\rho} = e \frac{8.3 \text{ Vs/m}^2}{7000 * 10^9 \text{ eV/c}} = \frac{8.3 \text{ s}^3 * 10^8 \text{ m/s}}{7000 * 10^9 \text{ m}^2}$$




$$\frac{1}{\rho} = 0.333 \frac{8.3}{7000} \text{ 1/m}$$

Gives:

$$\rho = 2.53 \text{ km} \longrightarrow 2\pi\rho = 17.6 \text{ km} \approx 66\%$$

- ◆ Need strong magnets to bend high energy beam!
- ◆ Most of LHC circumference used by dipole magnets!
- ◆ In fact this limits maximum energy of LHC beams!

LHC dipole magnets



- ◆ Edge of present technology
- ◆ NbTi superconductors used at 2° K
- ◆ Magnetic fields up to 8 T
- ◆ Two-in-one (twin bore) design for two beam in common cryostat
- ◆ 1232 dipole magnets, each 15m long

Keeping particles on circle - dipoles





Table 3.4: Main parameters of the dipole cold mass.

| | Value | Unit |
|--|---------|------|
| Injection field (0.45 TeV beam energy) | 0.54 | T |
| Current at injection field | 763 | A |
| Nominal field (7 TeV beam energy) | 8.33 | T |
| Current at nominal field | 11850 | A |
| Inductance at nominal field | 98.7 | mH |
| Stored energy (both apertures) at nominal field | 6.93 | MJ |
| Ultimate field | 9.00 | T |
| Current at ultimate field | 12840 | A |
| Stored energy (both apertures) at ultimate field | 8.11 | MJ |
| Maximum quench limit of the cold mass (from short samples) | 9.7 | T |
| Operating temperature | 1.9 | K |
| Magnetic length at 1.9 K and at nominal field | 14312 | mm |
| Distance between aperture axes at 1.9 K | 194.00 | mm |
| Cold mass sagitta at 293 K | 9.14 | mm |
| Bending radius at 1.9 K | 2803.98 | m |
| Inner coil diameter at 293 K | 56.00 | mm |
| Number of conductor blocks / pole | 6 | |
| Number of turns / pole, inner layer | 15 | |
| Number of turns / pole, outer layer | 25 | |
| Electromagnetic forces / coil quadrant at nominal field | | |
| Horizontal force component (inner and outer layer) | 1.8 | MN/m |
| Vertical force component (inner and outer layer) | 0.81 | MN/m |
| Electromagnetic forces / coil quadrant at ultimate field | | |
| Horizontal force component (inner and outer layer) | 2.1 | MN/m |
| Vertical force component (inner and outer layer) | 0.94 | MN/m |
| Axial electromagnetic force at each ends at nominal field | 0.40 | MN |
| Coil aperture at 293 K | 56.00 | mm |
| Cold tube inner diameter at 293 K | 50.00 | mm |
| Cold tube outer diameter at 293 K | 53.00 | mm |
| Cold mass length at 293 K (active part) | 15.18 | m |
| Cold mass diameter at 293 K | 570.0 | mm |
| Cold mass overall length with ancillaries | 16.5 | m |
| Total mass | ~ 27.5 | t |


Squeezing the beam - quadrupoles I

- ▶ Want to keep particles rotating on (around) reference trajectories
- ▶ Problem to keep the beam together
 - even small disturbances (for example gravity) may lead to lost particles
- ▶ restoring force of the type $F=-kx$, $F=-ky$ would keep the particles close to the ideal orbit!




Squeezing the beam - quadrupoles II

- ◆ Magnet surfaces shaped as hyperbolas give linear field!
- ◆ $B_x = -gy$
- ◆ $B_y = -gx$
- ◆ Quadrupole magnets!
- ◆ Unfortunately, forces are focusing in one plane and defocusing in the orthogonal plane
- ◆ $F_x = -qvgx$
- ◆ $F_y = qvgy$
- ◆ Opposite focusing/defocusing is achieved by rotating the magnet by 90°




Strong focusing and FODO lattice I



Analogy with optics

- ▶ Alternating focusing and defocusing lenses give together total focusing effect in both planes
 - Strong focusing, one of big ideas in accelerator physics
- ▶ Modern accelerators - using FODO (FocusingDefocusing) structures
- ▶ Particles oscillate around nominal trajectories - betatron oscillations

Strong focusing and FODO lattice II



The envelope around all the trajectories of the particles circulating in the machine is called β -function:

- ◆ Minimum at QD, maximum at QF
- ◆ Property of particular machine (beam optics)

Beam size:
$$\sigma_{x,y} = \sqrt{\epsilon\beta_{x,y}}$$

ϵ is the emittance of the beam:


- ◆ describes the quality of the beam

Strong focusing and FODO lattice III

Typically particles in accelerator have Gaussian particle distribution in position and angle.

$$\rho(x) = \frac{N}{\sqrt{2\pi}\sigma_x} \cdot e^{-\frac{x^2}{2\sigma_x^2}}$$

$$\sigma_x = \sqrt{\epsilon\beta_x}$$



LHC quadrupoles





Table 3.5: Parameter list for main quadrupole magnets at 7.0 TeV.


| | | |
|---|----------------------------|---------------|
| Integrated Gradient | 690 | T |
| Nominal Temperature | 1.9 | K |
| Nominal Gradient | 223 | T/m |
| Peak Field in Conductor | 6.85 | T |
| Temperature Margin | 2.19 | K |
| Working Point on Load Line | 80.3 | % |
| Nominal Current | 11870 | A |
| Magnetic Length | 3.10 | m |
| Beam Separation distance (cold) | 194.0 | mm |
| | | |
| Inner Coil Aperture Diameter (warm) | 56.0 | mm |
| Outer Coils Diameter | 118.44 | mm |
| Outer Yoke diameter | 452 | mm |
| Collar Material | Austenitic Steel | |
| Yoke Material | Low Carbon Steel | |
| Yoke Length including End Plates | 3250 | mm |
| | | |
| Cold Mass Length Between End Covers | 5345 | mm |
| Total Mass Including Correctors | 6500 | kg |
| | | |
| Number of turns per Coil (pole) | 24 | |
| Number of turns per coil inner layer (2 blocks) | 2+8 | |
| Number of turns per coil outer layer (2 blocks) | 7+7 | |
| Cable length per coil (pole) | 160 | m |
| Cable length per two-in-one quadrupole | 1280 | m |
| | | |
| Bare Cable | Same as dipole outer layer | |
| Insulation Thickness 1 st layer | 50 | μm |
| 2 nd layer | 37.5 | μm |
| 3 rd layer (adhesive) | 50+5 | μm |
| Self-inductance, one aperture | 5.6 | mH |
| Stored energy, one aperture | 395 | KJ |
| Electromagnetic forces: Resultant in x-dir | 537 | KN |
| Resultant in y-dir | -732 | KN |

LHC layout


- ◆ Circumference = 26658.9 m
- ◆ 8 arcs and 8 straight sections
 - Straight sections have either experiment or "utilities"
 - Four used for experiments
 - Arcs contain magnets (LHC lattice)
 - 23 FODO cells in each arc
 - A FODO cell consists of 2 quadrupoles, 6 dipoles and additional correction magnets




The LHC FODO cell



Each LHC arc consists of 23 FODO cells




Getting particles to collide - interaction point I



Special drift: Minibeta insertion

Minibeta insertion is a symmetric drift space with a beta waist in the center of the insertion




On each side of the symmetry point a quadrupole doublet or triplet are used to generate the waist.


They are not part of the regular lattice.

E.g. collider experiments are located in minibeta insertions: smallest beam size possible for the colliding beam to increase probability of collisions.

Minibeta insertion – Example LHC




Getting particles to collide - interaction point I



Bringing beams together for collisions:

- ◆ Bend them in dedicated dipole magnets (D1, D2 on the figure)
- ◆ Then squeeze them as much as possible just before the collision point
 - Low- β triplets

Getting particles to collide - interaction point III



- ◆ Low beta triplets :
 - Set of quadrupole magnets designed to squeeze beam before interaction point

$$L = f \frac{n_1 n_2}{4 \pi \sigma_x \sigma_y}$$





- ◆ Luminosity depends on:
- ◆ Number of particles per bunch (n_1, n_2)
- ◆ Bunch transverse size at the interaction point (σ_x, σ_y)
- ◆ Bunch collision rate f

And many others ...

- ♦ Vacuum system
- ♦ Beam injection system
- ♦ Beam dumping
- ♦ Pre-accelerators
- ♦ Cryogenic system
- ♦ Power distribution and protection
- ♦ Correction magnets
- ♦ Beam monitoring
- ♦ Control system
- ♦ ...



LHC in the near future (Run 3)

Current LHC schedule



Last update: April 2023

2024 LHC Plan



- Start of 2024 operation **2 weeks earlier** compared to 2023
- 1 week longer re-commissioning period for commissioning of reserve polarity optics (local optics change)
- **Pb ion running** days recalculated balancing unforeseen stops in 2022 and 2023: 5.5 weeks left, divided in both years 2024 / 2025
 - pp reference run attached to Pb ion run periods
 - Exact details of pp reference run schedule ongoing (preferred choice is to remove it for 2025)
- **Oxygen run moved to 2025**

Integrated luminosity expectations:

Protons:

- ATLAS/CMS collected $\sim 40+32 \text{ fb}^{-1}$ in 2022 and 2023
 - Initial wish to collect as close as possible to 300 fb^{-1} in Run 3
 - **Expectation for 2024 is $\sim 90 \text{ fb}^{-1}$ and for 2025 $\sim 100 \text{ fb}^{-1}$**
 \Rightarrow Expect now $\geq 260 \text{ fb}^{-1}$

Ions:

- 66% target reached in 2023 (1.75 nb^{-1})
- **Expectation for $\sim 5.35 \text{ nb}^{-1}$ for full Run 3** in ATLAS/CMS/ALICE
 - **1.9 nb^{-1} in 2024 (18d), 1.45 nb^{-1} (15 days) in 2025**
- pp reference targets: $250-350 \text{ pb}^{-1}$ ATLAS/CMS
- No time for pPb

LHC version 1.0


New since last week

2024 LHC Plan (under discussion)



Silvia Franchino, Place your bets - LHC in 2024, L1Calo Joint Meeting 29.02.24

Possible LHC schedule change under discussion



Ongoing discussions
Baseline: Rigid shift of the Ions Run and an extension of the pp Run


- LHC 5-week shift of the YETS, but 19 weeks length maintained
- The injectors schedule would shift by 6 weeks and be reduced by 4 weeks
- Additional electricity cost (mainly SPS) being evaluated
- Possible impact on various technical activities (machine and experiments) being evaluated

R. Steerenberg | SPSC#150 Accelerator Complex Status Report




pp integrated luminosity, 2023 predictions vs reality


what 2023 could have been



Chamonix 2024 Workshop: LHC Availability 2023 Michi Hostettler



Run3 vs Run2 faults



**long faults:
dominant for run 3**



**faults < 1 day:
stable ~16%**

thanks to the continuous effort of the equipment groups!



may: RF fingers @ 4L1

| | | | | | |
|---|---|---------------------------------|--|----------------------------------|-----------------------------------|
| Description & Root Cause | Localized heating (>500°C) of spring triggers localized plasticization with consequent loss of electrical contact. Heating due to small irregularities between finger gaps enhancing impedance and power dissipation. | | | | |
| Impact | ~4 days for repair & performance impact during recovery Thereafter, operation limited to intensity < 1.6x10¹¹ ppb | | | | |
| Wear-Out or Aging phenomenon? | Beam intensity related? | Random Hardware Failure? | Due to modifications or upgrades? | Inadequate specification? | Production non-conformity? |
| Possibly | Yes | No | No | Possibly * | Likely ** |
| Mitigation strategy & status | 71 ID212 modules to be replaced by new DRF (deformable RF bridges) design. 47 modules replaced this YETS; the remaining 24 modules in YETS 24/25. | | | | |
| Outlook | Similar problems can be expected until all modules are replaced. Other types of warm modules may also be affected, impedance studies ongoing (<i>details in the talk of C. Zannini</i>). | | | | |




* there was no clear specification in 2002-2003 apart the one done by the old vacuum group about the pumping speed and the design was for the LHC era.
 ** no non-conformities from mechanical or vacuum point of view, but with high intensity beam small deviations from the design could trigger impedance problems

input from G. Berglozzi, P. Krkotic, C. Antuono, C. Zannini

july: IT.L8 vacuum leak

| | | | | | |
|--|--|---------------------------------|--|-----------------------------------|-----------------------------------|
| Description & Root Cause | Due to electrical network glitch, the quench heaters of a few LHC magnets correctly triggered , including the ones of the IT.L8. As expected, the pressure inside the cold masses increased up to 18 bars but provoked a leak in an edge-welded bellow of the Q1-Q2 interconnection . The pressure in the vacuum vessel degraded, preventing the running of the LHC and demanding the bellow repair. It was found that a few bellow convolutions were partially blocked. | | | | |
| Impact | ~50 days total impact on physics runs | | | | |
| Wear-Out or Aging phenomenon? | Beam intensity related? | Random Hardware Failure? | Due to modifications or upgrades? | Inadequate specification ? | Production non-conformity? |
| <p style="color: orange; text-align: center;"> The Inner Triplet vacuum leak remains a concern – these will be replaced in LS3 but there is little that can be done before. The one that broke was known to be slightly worse than the others though... </p> | | | | | |
| Outlook | Similar events possible on triplets and mainly in Q1-Q2 interconnection until LS3. Inspection not possible without warm-up, which would pose a significant risk due to a thermal cycle of irradiated triplets. Consolidation of IP2 and IP8 proposed in LS3 (IP1 and IP5 replaced by HL upgrade). | | | | |



L8



repaired interconnect


input from S. Le Naour

september: TDIS8 vacuum leak


| | | | | | |
|---|---|---------------------------------|--|----------------------------------|-----------------------------------|
| Description & Root Cause | Two TDIS jaw actuator bellows developed vacuum leaks. Repaired by applying varnish and blocking movement. Resulted in degraded function, allowing ion run but preventing proton run at nominal intensity. Root cause was a misspecification of the bellow , causing wear-out after 2-3 years. | | | | |
| Impact | Intensity per injection severely limited. End of high-intensity proton physics. Ion run extended, p-p reference run moved to 2024. | | | | |
| Wear-Out or Aging phenomenon? | Beam intensity related? | Random Hardware Failure? | Due to modifications or upgrades? | Inadequate specification? | Production non-conformity? |
| Yes, due to inadequate spec. | No | No | Yes (LS2) | Yes | No * |
| Mitigation strategy & status | Both TDIS (points 2 and 8) replaced by existing (non-conform) spares during YETS 23/24. Expected cycle life covers the year of operation. Will be replaced again during YETS 24/25 by conform spares (based on refurbished TDIS with new bellows). Until then, movement should be limited (spares available only from summer 2024). New spare TDIS tanks will be built in parallel as back-up. | | | | |
| Outlook | Risk until YETS 24/25. New bellows should avoid similar problems until the end of the devices' lifetime (i.e. ~20 years). | | | | |

* Cannot exclude a non-conformity until the faulty bellows have been analysed in detail (to be done during the next months).

input from A. Perillo Marcone, C. Sharp



actuator bellows in TDIS



TDIS taken out of Pt 8

Conclusions ...

- ♦ LHC is a complicated and fascinating machine
 - Many interesting sub-systems and sub-components one could speak about for days!
- ♦ Run 3 is currently planned to last for two more years (2024 and 2025)
 - It should give us the best data set for analysis available so far
 - Keep your fingers crossed!
 - Machine is 15 years old by now!

Slides that were not good
enough to make it in to the
talk

The H boson is not just ... "yet another particle"

- ❑ Profoundly different from all elementary particles discovered so far
- ❑ Related to the most obscure sector of SM
- ❑ Linked to some of the deepest structural questions (flavour, naturalness, vacuum, ...)



Its discovery opens new paths of exploration, and a very broad and challenging experimental programme

Every problem of the SM originates from Higgs interactions

$$\mathcal{L} = \lambda H \psi \bar{\psi} + \mu^2 |H|^2 - \lambda |H|^4 - V_0$$

↑
↑
↑
↑
 flavour naturalness stability C.C.


G.F. Giudice

- ❑ Precision measurements of couplings (as many generations as possible, loops, ...)
- ❑ Forbidden and rare decays (e.g. $H \rightarrow \tau\mu$) \rightarrow flavour structure and source of fermion masses
- ❑ H potential (HH production, self-couplings):
 - \rightarrow EWSB mechanism (strong dynamics?)
 - \rightarrow EW phase transition \rightarrow baryogenesis?
- ❑ Exotic decays (e.g. $H \rightarrow E_T^{miss}$) \rightarrow new physics?
- ❑ Other H properties (width, CP, ...)
- ❑ Searches for additional H bosons
- ❑ ...

New Year CERN directorate meeting

13 January 2022, Agenda: <https://indico.cern.ch/event/1106493/>

Fabiola on long-term HL-LHC planning



13

HL-LHC, luminosity upgrade

Higher intensity

Increase bunch intensity

$$\mathcal{L} = \frac{N_1 N_2 f_{\text{rev}} k_B}{4\pi\beta^* \epsilon_{xy}} F$$

Increase F: shorter bunches, smaller crossing angle, crabs


$$\frac{1}{\sqrt{1 + \left(\frac{\sigma_s \phi}{\sigma_x} \frac{\phi}{2}\right)^2}}$$

Smaller β^*

Smaller beam size



Smaller emittance

New, wide-aperture quadrupoles




- ◆ Requires new, Nb_3Sn technology

Current LHC, operation with crossing angle



HL-LHC, crossing angle compensation using crab cavities



Crab Cavity

- Create an oscillating transverse electric field
- Kick head and tail of the bunch in opposite directions
- Serving to mitigate the effect of the crossing angle at the IP





Figure 4. Electric (left) and magnetic (right) field distributions inside the DQWCC.

Perspectives of Energy

Bringing together physics goals and collider energy



[J. R. Reuter]

Thresholds and cross sections set collider energy targets:

- 91.2 GeV** - The Z pole
- 160 GeV** - The WW threshold
- 250 GeV** - The ZH maximum
- 350 GeV** - The top threshold, VBF Higgs production
- 500 GeV** - ttH, ZHH
- 1+ TeV** - VBF double Higgs

- Precision electroweak, Flavour, QCD, ...
- Higgs properties & couplings
- Top properties, Top as probe
- Direct top Yukawa
- Higgs selfcoupling
- Search at the energy frontier


The International Linear Collider

e^+e^- Collider - Construction in Japan?

ILC in Japan: A linear e^+e^- collider at 250 GeV, upgradeable to 500 GeV

- perspectives to 1+ TeV

- Established technology - but no decision (yet)
- The main technology: Superconducting acceleration structure
- Well-proven technology by now: XFEL at DESY based on this, now in operation



FUTURE CIRCULAR COLLIDERS

International **FCC** collaboration (CERN as host lab) to study:

- **pp-collider (FCC-hh)** → main emphasis, defining infrastructure requirements
- **~100 km tunnel infrastructure** in Geneva area, site specific
- **e⁺e⁻ collider (FCC-ee)**, as potential first step
- **HE-LHC** with *FCC-hh* technology
- **p-e (FCC-he) option**, IP integration, e⁻ from ERL


CDRs published in **European Physical Journal C (Vol 1) and ST (Vol 2 – 4)**

Summary documents provided to EPPSU SG

- **FCC-integral, FCC-ee, FCC-hh, HE-LHC**
- Accessible on <http://fcc-cdr.web.cern.ch/>

Cost: ~28.6 BCHF


Power: ~580 MW (hh) ≤ 340 MW (ee)



| | LHC | HL-LHC | FCC-hh | |
|--|------|---------|-------------------------|----------|
| | | | Initial | Ultimate |
| c.m. Energy [TeV] | | 14 | | 100 |
| Peak luminosity [$10^{34} \text{ cm}^{-2} \text{ s}^{-1}$] | 1.0 | 5.0 | 5.0 | < 30.0 |
| Optimum integrated lumi / day [fb^{-1}] | 0.47 | 2.8 | 2.2 | 8 |
| Circumference [km] | | 26.7 | | 97.75 |
| Arc filling factor | | 0.79 | | 0.8 |
| Straight sections | | 8 × 528 | 6 × 1400 m + 2 × 2800 m | |
| Number of IPs | | 2 + 2 | 2 + 2 | |
| Injection energy [TeV] | | 0.45 | 3.3 | |

MOUNS COLLIDER SCHEME

Would be easy if the muons did not decay: lifetime is $\tau = \gamma \times 2.2 \mu\text{s}$



Short, intense proton bunch



Protons produce pions which decay into muons
muons are captured

Ionisation cooling of muon in matter

Acceleration to collision energy

Collision

Conclusions and outlook

- ▶ Elementary particle physics is in very exciting period indeed !
- LHC Runs 1 and 2 were very successful !!
- At the moment preparing for Run 3, hopefully early 2022
- Long-term future of particle physics at CERN is bright !!!!

LHC - 2015

- Target energy: 6.5 TeV
 - to be confirmed at end of powering tests!!!
- Bunch spacing: 25 ns
 - strongly favored by experiments (pile-up limit around 50)
- Beta* in ATLAS and CMS: 80 to 40 cm

Energy


- Lower quench margins
- Lower tolerance to beam loss
- Lower intensity set-up beams
- Hardware closer to maximum (beam dumps, power converters etc.)

25 ns

- Electron-cloud
- UFOs
- More long range collisions
- Larger crossing angle, higher beta*
- Higher total beam current
- Higher intensity per injection

How?

- **Beam from injectors**
 - High bunch population, low emittance, 25 ns beam
- **Lower beta* (~15 cm)**
 - New inner triplet magnets - wide aperture Nb₃Sn
 - Large aperture NbTi separator magnets
 - Novel optics solutions
- **Crossing angle compensation**
 - Crab cavities
- **Dealing with the regime**
 - Collision debris, high radiation
 - High machine availability
 - Beam stability, losses etc.




HL-LHC: key 25 ns parameters

| | |
|----------------------------|---|
| Protons per bunch | 2.2×10^{11} |
| Number of bunches | 2750 |
| Normalized emittance | 2.5 micron |
| Beta* | 15 cm |
| Crossing angle | 590 microrad |
| Geometric reduction factor | 0.305 |
| Virtual luminosity | $2.4 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ |
| Levelled luminosity | $5 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ |
| Levelled <pile-up> | 140 |

Future e^+e^- colliders

| \sqrt{s} (GeV) | Main physics goals |
|------------------|---|
| 90 | Z-pole precision EW measurements beyond LEP, SLC |
| 160 | WW precision physics (mass at threshold) |
| 250 | H precision physics (HZ) |
| ~350 | H (HZ, H $\nu\nu$) and top (mass, couplings) precision physics |
| 500-3000 | t \bar{t} H, HH (self-couplings), direct searches for new physics |



- Linear:
- Larger \sqrt{s} reach
 - Low repetition rate
 - L from nm size beams
 - large beamstrahlung
 - larger E-spread
 - Long. polarization easier

- Circular:
- \sqrt{s} limited by SR $\sim E_{\text{beam}}^4/R$
 - Large number of circulating bunches → high L (increases at lower \sqrt{s} as less SR → spare RF power used to accelerate more bunches). Note: need top-up injection ring to compensate fast L burn-off (lifetime $\sim 30'$)
 - Several interaction regions possible
 - Precise E-beam measurement from resonant depolarization

Searches for physics beyond SM

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: March 2016

ATLAS Preliminary

$\sqrt{s} = 7, 8, 13$ TeV

| Model | e, μ, τ, γ | Jets | E_T^{miss} | $\int \mathcal{L} dt [\text{fb}^{-1}]$ | Mass limit | Reference | | | | |
|--|--|--|---------------------|--|--------------------|------------------------|---|---|--|------------|
| | | | | | | $\sqrt{s} = 7, 8$ TeV | $\sqrt{s} = 13$ TeV | | | |
| Inclusive Searches | MSUGRA/CMSSM | 0-3 e, μ /1-2 τ | 2-10 jets/3 b | Yes | 20.3 | \tilde{g}, \tilde{g} | 1.85 TeV | $m(\tilde{g})=m(\tilde{g})$ | 1507.05525 | |
| | $\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ | 0 | 2-6 jets | Yes | 3.2 | \tilde{q} | 980 GeV | $m(\tilde{q})=0$ GeV, $m(1^{\text{st}} \text{ gen. } \tilde{q})=m(2^{\text{nd}} \text{ gen. } \tilde{q})$ | ATLAS-CONF-2015-062 | |
| | $\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ (compressed) | mono-jet | 1-3 jets | Yes | 3.2 | \tilde{q} | 610 GeV | $m(\tilde{q})=m(\tilde{q}_1^0)<5$ GeV | To appear | |
| | $\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\ell(\ell\ell/\nu\nu)/\tilde{\chi}_1^0$ | 2 e, μ (off-Z) | 2 jets | Yes | 20.3 | \tilde{q} | 820 GeV | $m(\tilde{q}_1^0)=0$ GeV | 1503.03290 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{\chi}_1^0$ | 0 | 2-6 jets | Yes | 3.2 | \tilde{g} | 1.52 TeV | $m(\tilde{q}_1^0)=0$ GeV | ATLAS-CONF-2015-062 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{\chi}_1^0 \rightarrow qgW^{\pm}\tilde{\chi}_1^0$ | 1 e, μ | 2-6 jets | Yes | 3.3 | \tilde{g} | 1.6 TeV | $m(\tilde{q}_1^0)<350$ GeV, $m(\tilde{\chi}^{\pm})=0.5(m(\tilde{q}_1^0)+m(\tilde{g}))$ | ATLAS-CONF-2015-076 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow qg(\ell\ell/\nu\nu)/\tilde{\chi}_1^0$ | 2 e, μ | 0-3 jets | - | 20 | \tilde{g} | 1.38 TeV | $m(\tilde{q}_1^0)=0$ GeV | 1501.03555 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow qgWZ\tilde{\chi}_1^0$ | 0 | 7-10 jets | Yes | 3.2 | \tilde{g} | 1.4 TeV | $m(\tilde{q}_1^0)=100$ GeV | 1602.06194 | |
| | GMSB ($\tilde{\ell}$ NLSP) | 1-2 τ + 0-1 ℓ | 0-2 jets | Yes | 20.3 | \tilde{g} | 1.63 TeV | $\tan\beta > 20$ | 1407.0603 | |
| | GGM (bino NLSP) | 2 γ | - | Yes | 20.3 | \tilde{g} | 1.34 TeV | $c\tau(\text{NLSP})<0.1$ mm | 1507.05493 | |
| | GGM (higgsino-bino NLSP) | γ | 1 b | Yes | 20.3 | \tilde{g} | 1.37 TeV | $m(\tilde{q}_1^0)<950$ GeV, $c\tau(\text{NLSP})<0.1$ mm, $\mu<0$ | 1507.05493 | |
| | GGM (higgsino-bino NLSP) | γ | 2 jets | Yes | 20.3 | \tilde{g} | 1.3 TeV | $m(\tilde{q}_1^0)>850$ GeV, $c\tau(\text{NLSP})<0.1$ mm, $\mu>0$ | 1507.05493 | |
| GGM (higgsino NLSP) | 2 e, μ (Z) | 2 jets | Yes | 20.3 | \tilde{g} | 900 GeV | $m(\text{NLSP})>430$ GeV | 1503.03290 | | |
| Gravitino LSP | 0 | mono-jet | Yes | 20.3 | $F^{1/2}$ scale | 865 GeV | $m(\tilde{g})>1.8 \times 10^{-4}$ eV, $m(\tilde{g})=m(\tilde{g})=1.5$ TeV | 1502.01518 | | |
| 3 rd gen. \tilde{g} med. | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$ | 0 | 3 b | Yes | 3.3 | \tilde{g} | 1.78 TeV | $m(\tilde{q}_1^0)>800$ GeV | ATLAS-CONF-2015-067 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$ | 0-1 e, μ | 3 b | Yes | 3.3 | \tilde{g} | 1.76 TeV | $m(\tilde{q}_1^0)=0$ GeV | To appear | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$ | 0-1 e, μ | 3 b | Yes | 20.1 | \tilde{g} | 1.37 TeV | $m(\tilde{q}_1^0)<300$ GeV | 1407.0600 | |
| 3 rd gen. squarks direct production | $\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$ | 0 | 2 b | Yes | 3.2 | \tilde{b}_1 | 840 GeV | $m(\tilde{q}_1^0)<100$ GeV | ATLAS-CONF-2015-066 | |
| | $\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^0$ | 2 e, μ (SS) | 0-3 b | Yes | 3.2 | \tilde{b}_1 | 325-540 GeV | $m(\tilde{q}_1^0)=50$ GeV, $m(\tilde{t}_1^0)=m(\tilde{t}_1^0)+100$ GeV | 1602.09059 | |
| | $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{\chi}_1^0$ | 1-2 e, μ | 1-2 b | Yes | 4.7/20.3 | \tilde{t}_1 | 117-170 GeV | $m(\tilde{q}_1^0)=2m(\tilde{q}_1^0), m(\tilde{q}_1^0)=55$ GeV | 1209.2102, 1407.0583 | |
| | $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^0$ or $t\tilde{\chi}_1^0$ | 0-2 e, μ | 0-2 jets/1-2 b | Yes | 20.3 | \tilde{t}_1 | 90-198 GeV | $m(\tilde{q}_1^0)=1$ GeV | 1506.08616, ATLAS-CONF-2016-007 | |
| | $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$ | 0 | mono-jet/c-tag | Yes | 20.3 | \tilde{t}_1 | 90-245 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0)<85$ GeV | 1407.0608 | |
| | $\tilde{t}_1\tilde{t}_1$ (natural GMSB) | 2 e, μ (Z) | 1 b | Yes | 20.3 | \tilde{t}_1 | 150-600 GeV | $m(\tilde{q}_1^0)>150$ GeV | 1403.5222 | |
| | $\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$ | 3 e, μ (Z) | 1 b | Yes | 20.3 | \tilde{t}_2 | 290-610 GeV | $m(\tilde{q}_1^0)<200$ GeV | 1403.5222 | |
| $\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + h$ | 1 e, μ | 6 jets + 2 b | Yes | 20.3 | \tilde{t}_2 | 320-620 GeV | $m(\tilde{q}_1^0)=0$ GeV | 1506.08616 | | |
| EW direct | $\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$ | 2 e, μ | 0 | Yes | 20.3 | $\tilde{\ell}$ | 90-335 GeV | $m(\tilde{q}_1^0)=0$ GeV | 1403.5294 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell\nu(\ell\nu)$ | 2 e, μ | 0 | Yes | 20.3 | $\tilde{\chi}_1^0$ | 140-475 GeV | $m(\tilde{q}_1^0)=0$ GeV, $m(\tilde{\ell}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{q}_1^0))$ | 1403.5294 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}\nu(\tilde{\tau}\nu)$ | 2 τ | - | Yes | 20.3 | $\tilde{\chi}_1^0$ | 355 GeV | $m(\tilde{q}_1^0)=0$ GeV, $m(\tilde{\tau}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{q}_1^0))$ | 1407.0350 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0 \rightarrow \tilde{\nu}_i\tilde{\nu}_j(\tilde{\nu}_i\tilde{\nu}_j), \ell\tilde{\nu}_i\ell(\tilde{\nu}_i\tilde{\nu}_j)$ | 3 e, μ | 0 | Yes | 20.3 | $\tilde{\chi}_1^0$ | 715 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0), m(\tilde{q}_1^0)=0, m(\tilde{\ell}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{q}_1^0))$ | 1402.7029 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\nu}_i$ | 2-3 e, μ | 0-2 jets | Yes | 20.3 | $\tilde{\chi}_1^0$ | 425 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0), m(\tilde{q}_1^0)=0, \text{ sleptons decoupled}$ | 1403.5294, 1402.7029 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0 \rightarrow W\tilde{\chi}_1^0 h, \tilde{h} \rightarrow b\tilde{b}/WV/\tau\tau/\gamma\gamma$ | e, μ, γ | 0-2 b | Yes | 20.3 | $\tilde{\chi}_1^0$ | 270 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0), m(\tilde{q}_1^0)=0, \text{ sleptons decoupled}$ | 1501.07110 | |
| | $\tilde{\chi}_{2,3}^0\tilde{\chi}_{2,3}^0 \rightarrow \tilde{\ell}_R\tilde{\ell}_R$ | 4 e, μ | 0 | Yes | 20.3 | $\tilde{\chi}_{2,3}^0$ | 635 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0), m(\tilde{q}_1^0)=0, m(\tilde{\ell}, \tilde{\nu})=0.5(m(\tilde{q}_1^0)+m(\tilde{q}_1^0))$ | 1405.5086 | |
| | GGM (wino NLSP) weak prod. | 1 $e, \mu + \gamma$ | - | Yes | 20.3 | \tilde{W} | 115-370 GeV | $c\tau < 1$ mm | 1507.05493 | |
| | Long-lived particles | Direct $\tilde{\chi}_1^0\tilde{\chi}_1^0$ prod., long-lived $\tilde{\chi}_1^0$ | Disapp. trk | 1 jet | Yes | 20.3 | $\tilde{\chi}_1^0$ | 270 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0)\sim 160$ MeV, $\tau(\tilde{\chi}_1^0)=0.2$ ns | 1310.3675 |
| | | Direct $\tilde{\chi}_1^0\tilde{\chi}_1^0$ prod., long-lived $\tilde{\chi}_1^0$ | dE/dx trk | - | Yes | 18.4 | $\tilde{\chi}_1^0$ | 495 GeV | $m(\tilde{q}_1^0)=m(\tilde{q}_1^0)\sim 160$ MeV, $\tau(\tilde{\chi}_1^0)<15$ ns | 1506.05332 |
| Stable, stopped \tilde{g} R-hadron | | 0 | 1-5 jets | Yes | 27.9 | \tilde{g} | 850 GeV | $m(\tilde{q}_1^0)=100$ GeV, $10 \mu\text{s}<\tau(\tilde{g})<1000$ s | 1310.6584 | |
| Metastable \tilde{g} R-hadron | | dE/dx trk | - | - | 3.2 | \tilde{g} | 1.54 TeV | $m(\tilde{q}_1^0)=100$ GeV, $\tau > 10$ ns | To appear | |
| GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{\ell}, \tilde{\mu}) + \tau(e, \mu)$ | | 1-2 μ | - | - | 19.1 | $\tilde{\chi}_1^0$ | 537 GeV | $10 < \tan\beta < 50$ | 1411.6795 | |
| GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$ | | 2 γ | - | Yes | 20.3 | $\tilde{\chi}_1^0$ | 440 GeV | $1 < \tau(\tilde{\chi}_1^0) < 3$ ns, SPS8 model | 1409.5542 | |
| $\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow \nu\tilde{\nu}/\mu\tilde{\mu}/\nu\tilde{\nu}$ | displ. ee/μμ/μμ | - | - | 20.3 | $\tilde{\chi}_1^0$ | 1.0 TeV | $7 < c\tau(\tilde{\chi}_1^0) < 740$ mm, $m(\tilde{g})=1.3$ TeV | 1504.05162 | | |
| GGM $\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow Z\tilde{G}$ | displ. vtx + jets | - | - | 20.3 | $\tilde{\chi}_1^0$ | 1.0 TeV | $6 < c\tau(\tilde{\chi}_1^0) < 480$ mm, $m(\tilde{g})=1.1$ TeV | 1504.05162 | | |
| RPV | LFV $pp \rightarrow \tilde{\nu}_i + X, \tilde{\nu}_i \rightarrow e\mu/\tau/\mu/\tau$ | $e\mu, \tau\mu, \mu\tau$ | - | - | 20.3 | $\tilde{\nu}_i$ | 1.7 TeV | $\lambda_{311}^e=0.11, \lambda_{132/133/233}=0.07$ | 1503.04430 | |
| | Bilinear RPV CMSSM | 2 e, μ (SS) | 0-3 b | Yes | 20.3 | \tilde{q}, \tilde{g} | 1.45 TeV | $m(\tilde{g})=m(\tilde{g}), c\tau_{\text{LSP}} < 1$ mm | 1404.2500 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow ee\nu_\mu, e\mu\tilde{\nu}_e$ | 4 e, μ | - | Yes | 20.3 | $\tilde{\chi}_1^0$ | 760 GeV | $m(\tilde{q}_1^0)>0.2 \times m(\tilde{q}_1^0), \lambda_{211} \neq 0$ | 1405.5086 | |
| | $\tilde{\chi}_1^0\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tau\tau\nu_e, e\tau\tilde{\nu}_e$ | 3 $e, \mu + \tau$ | - | Yes | 20.3 | $\tilde{\chi}_1^0$ | 450 GeV | $m(\tilde{q}_1^0)>0.2 \times m(\tilde{q}_1^0), \lambda_{133} \neq 0$ | 1405.5086 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{q}$ | 0 | 6-7 jets | - | 20.3 | \tilde{g} | 917 GeV | $\text{BR}(\tilde{g})-\text{BR}(\tilde{h})-\text{BR}(\tilde{c})=0\%$ | 1502.05686 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow qq\tilde{q}$ | 0 | 6-7 jets | - | 20.3 | \tilde{g} | 980 GeV | $m(\tilde{q}_1^0)=600$ GeV | 1502.05686 | |
| | $\tilde{g}\tilde{g}, \tilde{g} \rightarrow \tilde{\nu}_i\tilde{\nu}_j, \tilde{\nu}_i \rightarrow b\tilde{s}$ | 2 e, μ (SS) | 0-3 b | Yes | 20.3 | \tilde{g} | 880 GeV | - | 1404.2500 | |
| | $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{s}$ | 0 | 2 jets + 2 b | - | 20.3 | \tilde{t}_1 | 320 GeV | - | 1601.07453 | |
| | $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{\ell}$ | 2 e, μ | 2 b | - | 20.3 | \tilde{t}_1 | 0.4-1.0 TeV | $\text{BR}(\tilde{t}_1 \rightarrow b\ell/\mu) > 20\%$ | ATLAS-CONF-2015-015 | |
| | Other | Scalar charm, $\tilde{c} \rightarrow c\tilde{\chi}_1^0$ | 0 | 2 c | Yes | 20.3 | \tilde{c} | 510 GeV | $m(\tilde{q}_1^0)<200$ GeV | 1501.01325 |

*Only a selection of the available mass limits on new states or phenomena is shown.

Haven't found anything, but keep searching ...

Structure of the proton II


Inclusive DIS cross section:

$$\frac{d^2\sigma_{ep \rightarrow eX}}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} (y^2 xF_1(x, Q^2) + (1-y)F_2(x, Q^2))$$

- Q - virtuality of exchanged photon
- x - Bjorken x


In Leading Order:

$$F_2(x, Q^2) = 2xF_1(x, Q^2) = x \sum_q e_q^2 (q(x, Q^2) + \bar{q}(x, Q^2))$$



Parton distributions ($q(x, Q^2)$) are obtained from (QCD) fits to cross section of various processes (ep NC, CC, high P_T jet production, ...)

Kinematics of produced particles I



We are interested in momentum of produced particles:

- ◆ Could use p_x, p_y, p_z ...
- ◆ Geometry of collision is cylindrical, can use cylindrical coordinates
 - P, θ, ϕ
- ◆ Physics is symmetric in ϕ
- ◆ The fact that collisions are not collisions between pointlike particles complicates kinematic analysis
 - Total longitudinal momentum of elementary collision is not known
 - Transversal momentum (P_T) is conserved (and used very often)

Produced particles - kinematics II

→ Usually do not use P and θ , but rapidity:

$$y = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right)$$


♦ Rapidity interval Δy and P_T are invariant with respect to Lorentz boosts along beam direction!

$$m_T^2 = m^2 + p_T^2$$

$$E = m_T \cosh(y) \quad p_z = m_T \sinh(y)$$

♦ For zero mass particles (or high p) rapidity is equal to pseudorapidity η :

$$\eta = -\ln \left(\tan \left(\frac{\theta}{2} \right) \right)$$



Production of massive particles

- Two partons (with x_1, x_2) inside of two protons (with $p_{\text{proton A}}, p_{\text{proton B}}$) collide, create a heavy (new!) particle with mass M and rapidity y_M

$$M^2 = (x_1 p_{\text{proton A}} + x_2 p_{\text{proton B}})^2 \quad \longrightarrow \quad x_1 x_2 = \frac{M^2}{s}$$

- Higher x means higher M
- To produce mass of 100 GeV with accelerator running at 14 TeV requires $x=0.007$
- To produce mass of 5 TeV requires $x = 0.36$

$$p_{zM} = m_T sh(y_M) \rightarrow M sh(y_M) \quad x_1 = \left[\frac{M}{\sqrt{s}} \right] \exp(y_M)$$


$$p_{zM} = p_{z\text{parton 1}} - p_{z\text{parton 2}} = (x_1 - x_2) \frac{\sqrt{s}}{2} \quad \longrightarrow \quad x_2 = \left[\frac{M}{\sqrt{s}} \right] \exp(-y_M)$$

- To produce M at zero rapidity we need partons with same x , going to higher rapidities of particle M means one parton at higher x , the other one at smaller x

A consequence of phase stability

Longitudinal stability - particle that comes earlier gets accelerated less:

$$\frac{\partial V}{\partial t} > 0 \Rightarrow \frac{\partial E_z}{\partial z} < 0$$




$$\nabla \cdot \vec{E} = 0 \Rightarrow \frac{\partial E_x}{\partial x} + \frac{\partial E_z}{\partial z} = 0 \Rightarrow \frac{\partial E_x}{\partial x} > 0$$

- ◆ Maxwell equations show that this leads to de-focusing in transverse direction...
- ◆ Want to keep beam profile small, too ...
- ➔ Need some magnets ...

Squeeze in ATLAS

- Lower beta* implies larger beams in the triplet magnets
- Larger beams implies a larger crossing angle
- Aperture concerns dictate caution



Requirement: Lorentz force increases as a function of distance from design trajectory

E.g. in the horizontal plane

$$F(x) = q \cdot v \cdot B(x)$$

We want a magnetic field that

$$B_y = g \cdot x \quad B_x = g \cdot y$$


→ Quadrupole magnet

Gradient of quadrupole

$$g = \frac{2\mu_0 n I}{r^2} \left[\frac{T}{m} \right]$$

Normalized gradient, focusing strength

$$k = \frac{g}{p/q} [m^{-2}]$$



The red arrows show the direction of the force on the particle

The emittance at LHC injection energy 450 GeV: $\varepsilon = 7.3 \text{ nm}$

At 7 TeV: $\varepsilon = 0.5 \text{ nm}$

$$\varepsilon_{7\text{TeV}} = \varepsilon_{450\text{GeV}} \frac{\gamma_{450\text{GeV}}}{\gamma_{7\text{TeV}}}$$

Normalized emittance: $\varepsilon^* = 3.5 \text{ } \mu\text{m}$

Normalized emittance preserved during acceleration.

And for the beam sizes:


At the location with the maximum beta function ($\beta_{\text{max}} = 180 \text{ m}$):


$$\sigma_{450\text{GeV}} = 1.1 \text{ mm}$$

$$\sigma_{7\text{TeV}} = 300 \text{ } \mu\text{m}$$

Aperture requirement: $a > 10 \sigma$


Vertical plane: $19 \text{ mm} \sim 16 \sigma @ 450 \text{ GeV}$





High Energy LHC???

Higher \sqrt{s} in the LHC tunnel ?



Various options, with increasing amount of HW changes, technical challenges, cost, and physics reach

- 1) Pushing present dipoles to ultimate performance ($\rightarrow \sqrt{s} \sim 15.5$ TeV ?)
- 2) 1) + replacing 30% of present dipoles with higher-field magnets:
 - B=11 T $\rightarrow \sqrt{s} \sim 16.5$ TeV
 - B=14 T $\rightarrow \sqrt{s} \sim 18.5$ TeV
- HE-LHC: filling existing 16-20 T magnets $\rightarrow \sqrt{s} = 26-33$ TeV
 - strongly motivated if new physics discovered at the LHC/HL-LHC
 - demonstration of technology in view of future higher-E pp colliders
 - would capitalize on existing tunnel and infrastructure
 - magnets might be reused in a bigger tunnel ??

These options are being studied (physics case, technical feasibility, cost, time scale) in time for next round of European Strategy (~2018/2019)


F. Gianotti

Standard model and its problems


What did we accomplish so far in particle physics ?

With the discovery of the Higgs boson, we have completed the Standard Model (> 50 years of theoretical and experimental efforts !)

Note: fermions (c, b, t, τ) discovered at accelerators in the US, bosons (g, W, Z, H) in Europe ...



We have tested the Standard Model with very high precision (wealth of measurements since early '60s, in particular at accelerators)
→ it works BEAUTIFULLY (puzzling ...)
→ no significant deviations observed (but difficult to accommodate non-zero neutrino masses)





However: SM is not a complete theory of particle physics, as several outstanding questions remain (raised also by precise experimental observations) that cannot be explained within the SM.


These questions require NEW PHYSICS

But, can λ become negative? Yes, two main competing effects:

$$\mu \frac{d\lambda(\mu)}{d \log(\mu)} = (\# \lambda^2 + \dots - \# h_t^4 + \dots) + \dots$$



$$h_t(v) = \sqrt{2}M_t/v \quad \text{and} \quad \lambda(v) = M_h^2/(2v^2).$$



J.EM, J. Espinosa, G.F. Giudice, G. Isidori, A. Riotto, A. Strumia. [hep/1112.3022]

Circular colliders: the CERN FCC project






International conceptual design study for Future Circular Colliders in a ~100 km ring:

- goal: pp, $\sqrt{s} = 100 \text{ TeV}$ (FCC-hh), $L \sim 2 \times 10^{35}$; 4 IP
- possible intermediate step: e^+e^- , $\sqrt{s} = 90\text{-}350 \text{ GeV}$ (FCC-ee), $L = 2 \times 10^{36}\text{-}2 \times 10^{34}$, 2-4 IP
- option: ep, $\sqrt{s} = 3.5 \text{ TeV}$ (FCC-eh), $L \sim 10^{34}$

Goal of the study: CDR in ~2018

Machine studies are site-neutral.

However, FCC at CERN would greatly benefit from existing infrastructure (e.g. FCC-hh injector chain would be based on existing accelerator complex)



Preamble

- Many mysteries about the universe remain to be explored: **nature of dark matter, preponderance of matter over antimatter, origin and pattern of neutrino masses**
- Nature hides the secrets of the fundamental physical laws in the **tiniest nooks of space and time**
- Particle Physics develops technologies to probe ever smaller distance scales (higher energies)
- **The Higgs** (discovered at the LHC) is a **unique particle** that raises profound questions about the fundamental laws of nature
 - ✓ Higgs properties study is in itself a powerful experimental tool to look for answers
 - **electron-positron collider as Higgs factory**
 - ✓ Higgs boson pair-production study is key to understanding the fabric of the universe
 - **collider with significantly higher energies than Higgs factory**
- New realm of energies is expected to lead to new discoveries and provide answers to existing mysteries
- **The 2020 Strategy update aims to significantly extend knowledge beyond current limits, to drive innovative technological developments, to maintain Europe's leading role**

The European vision is thus to prepare a Higgs factory, followed by a future hadron collider with sensitivity to energy scales an order of magnitude higher than those of the LHC, while addressing the associated technical and environmental challenges.

The 2020 Strategy presents exciting and ambitious scientific goals that will drive technological and scientific exploration into new and uncharted territory for the benefit of the field and of society.

Guide through the statements

2 statements on **Major developments from the 2013 Strategy**

- a) Focus on successful completion of HL-LHC upgrade remains a priority
- b) Continued support for long-baseline experiments in Japan and US and the Neutrino Platform

3 statements on **General considerations for the 2020 update**

- a) Preserve the leading role of CERN for success of European PP community
- b) Strengthen the European PP ecosystem of research centres
- c) Acknowledge the global nature of PP research

2 statements on **High-priority future initiatives**

- a) Higgs factory as the highest-priority next collider and investigation of the technical and financial feasibility of a future hadron collider at CERN
- b) Vigorous R&D on innovative accelerator technologies

Letters for itemizing the statements are introduced for identification, do not imply prioritization

4 statements on **Other essential scientific activities**

- a) Support for high-impact, financially implementable, experimental initiatives world-wide
- b) Acknowledge the essential role of theory
- c) Support for instrumentation R&D
- d) Support for computing and software infrastructure

2 statements on **Synergies with neighbouring fields**

- a) Nuclear physics - cooperation with NuPECC
- b) Astroparticle - cooperation with APPEC

3 statements on **Organisational issues**


- a) Global collaboration on projects in and out of Europe
- b) Relations with European Commission
- c) Open science

4 statements on **Environmental and societal impact**

- a) Mitigate environmental impact of particle physics
- b) Investment in next generation of researchers
- c) Knowledge and technology transfer
- d) Cultural heritage: public engagement, education and communication

Electron cloud – One of the LHC Challenges

In high intensity accelerators with positively charged beams and closely spaced bunches electrons liberated from vacuum chamber surface can multiply and build up a **cloud of electrons**.



Electrons are generated through:


- Residual gas ionization
- Photo-electrons with synchrotron radiation
- Desorption from the losses on the wall

- 1) Seed electrons accelerated by beam
- 2) Produce secondary electrons when hitting chamber
- 3) Secondary electrons accelerated, producing more electrons on impact
- 4) May lead to exponential growth of electron density (multipacting)
- 5) Trailing bunches interact with cloud

Beam Dump – How to safely kill the LHC beam



Schematic of LHC beam dump system

LHC beam stores beam $\sim 360\text{MJ}$ energy.




Lecture on Accelerator Operation and Design Challenges by A. Lechner

Sweep of beam on beam dump window



Source graphics: <http://clipart-library.com>

LHC / HL-LHC Plan




Hard squeeze ...

| | 2016 | HL-LHC |
|-------------------------|---------------------|---------------------|
| β^* | 40 cm | 15 cm |
| Beam size at IP (sigma) | 17 μm | 7 μm |
| β at triplet | ~ 4.5 km | ~ 20 km |
| Beam size at triplet | 1.5 mm | 2.6 mm |
| | | |
| Crossing angle | 370 μrad | 590 μrad |

The reduction in beam size buys a factor of 1.6 in luminosity but:

- Bigger beams in inner triplets and so
- Larger crossing angle
- And thus larger aperture in inner triplets is required.

Luminosity leveling



~130 inelastic collisions/BC