



Contribution ID: 179

Type: **Parallel Talk**

Revisiting the Gamma-Ray Galactic Center Excess with Multi-Messenger Observations

Monday 8 August 2022 16:50 (20 minutes)

The Galactic Center Excess (GCE) remains one of the most intriguing discoveries from the Fermi Large Area Telescope (LAT) observations. I will revisit the characteristics of the GCE by using a new set of high-resolution galactic diffuse gamma-ray emission templates. This diffuse emission, which accounts for the bulk of the observed gamma rays, is ultimately due to cosmic-ray interactions with the interstellar medium. Using recent high-precision cosmic-ray observations, in addition to the continuing Fermi-LAT observations and observations from lower energy photons, we constrain the properties of the galactic diffuse emission. I will describe a large set of diffuse gamma-ray emission templates which account for a very wide range of initial assumptions on the physical conditions in the inner galaxy. The updated GCE spectral and morphological properties have an impact on its interpretations. In particular, a high-energy tail is found at higher significance than previously reported. This tail is very prominent in the northern hemisphere, and less so in the southern hemisphere. This strongly affects one prominent interpretation of the excess: known millisecond pulsars are incapable of producing this high-energy emission, even in the relatively softer southern hemisphere, and are therefore disfavored as the sole explanation of the GCE. The annihilation of dark matter particles of mass 40_{-7}^{+10} GeV (95% CL) to b quarks with a cross-section of $\sigma v = 1.4_{-0.3}^{+0.6} \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ provides a good fit to the excess especially in the relatively cleaner southern sky. Dark matter of the same mass range annihilating to b quarks or heavier dark matter particles annihilating to heavier Standard Model bosons can combine with a subdominant millisecond pulsars component to provide a good fit to the southern hemisphere emission as well, as can a broken power-law spectrum which would be related to recent cosmic-ray burst activity.

Collaboration name

Author: Dr CHOLIS, Ilias (Department of Physics, Oakland University)

Co-authors: Dr ZHONG, Yi-Ming (Kavli Institute for Cosmological Physics, University of Chicago); Dr MCDERMOTT, Samuel (Fermi National Accelerator Laboratory); Mr SURDUTOVICH, Joseph (Carleton College, Physics Department)

Presenter: Dr CHOLIS, Ilias (Department of Physics, Oakland University)

Session Classification: Dark Matter

Track Classification: Gamma Rays