

The General AntiParticle Spectrometer Search for Dark Matter using Cosmic-ray Antinuclei


**TeVPA
August 2022**

**Philip von Doetinchem
on behalf of the GAPS collaboration**

**philipvd@hawaii.edu
Department of Physics & Astronomy
University of Hawai'i at Mānoa
<http://www.phys.hawaii.edu/~philipvd>**




Unexplained features in cosmic rays




- unexplained feature in positrons:
 - astrophysical origin → pulsars
 - SNR acceleration
 - **dark matter annihilation**
- combined fit with antiproton and diffuse gamma-rays from the Galactic Center → 80GeV DM particle?
- understanding astrophysics background is a challenge better constraints on cosmic-ray propagation and astrophysical production are needed

antiproton




antiproton and gamma-ray results




Status Cosmic-ray Antinuclei Searches

- **Potential \bar{p} excess** in AMS-02 data above secondary background predictions at $R \sim 10$ GV was found in various studies → significance level unclear
- AMS-02 reported at conferences the observation of **antihelium candidates ($\sim 1/\text{year}$)** → interpretations are actively ongoing
- No explanation of antiproton nor antihelium **should overproduce antideuterons**
- **Possible physics models that explain antihelium candidates include:**
 - Secondary astrophysical background
 - Dark matter annihilation or decay
 - Nearby antistar: at distance of $\sim 1\text{pc}$
- **Search for antinuclei with independent technique is critical**
- Review based on 2nd Cosmic-ray Antideuteron Workshop “Cosmic-ray Antinuclei as Messengers of New Physics: Status and Outlook for the New Decade” [JCAP08(2020)035, arXiv:2002.04163]




The GAPS experiment



mass: ~2,500kg
power: 1.3kW

- The **General AntiParticle Spectrometer** is the first experiment dedicated and optimized for low-energy cosmic-ray antinuclei search
- Requirements: long flight time, large acceptance, large identification power
- **GAPS will deliver:**
 - a precision antiproton measurement in an unexplored energy range <0.25 GeV/n
 - antideuteron sensitivity 2 orders of magnitude below the current best limits, probing a variety of DM models across a wide mass range
 - leading sensitivity to low-energy cosmic antihelium nuclei
- **GAPS is under construction, preparing for first Antarctic Long Duration Balloon flight**

GAPS principle




- antiparticle slows down and stops in material
- large chance for creation of an excited exotic atom ($E_{\text{kin}} \sim E_I$)
- deexcitation:
 - fast ionization of bound electrons (Auger)
 - complete depletion of bound electrons
 - Hydrogen-like exotic atom (nucleus+antideuteron)
 - deexcites via characteristic X-ray transitions depending on antiparticle mass
- Nuclear annihilation with characteristic number of annihilation products

GAPS antideuteron model sensitivity

T. Aramaki et al., Astropart. Phys. 74, 6 (2016)


- **Low-energy antideuterons are essentially free of astrophysics background**
- GAPS is sensitive to a **wide range of dark matter models**, e.g.:
 - Generic 70GeV WIMP annihilation model that explains antiproton excess and γ -rays from Galactic center
 - Dark matter gravitino decay
 - Extra dimensions
 - Heavy DM models with Sommerfeld enhancement
 - Dark photons (inaccessible to other techniques)
- Selection of publications:
 - Braeuninger et al. Physics Letters B 678, 20–31 (2009)
 - Cui et al, JHEP 1011, 017 (2010)
 - Hryczuk et al., JCAP 1407, 031 (2014).
 - Korsmeier et al., Physical Review D 97, 103011 (2018)
 - Randall & Xu, JHEP (2020)



astrophysics background at $\sim 10^{-7}-10^{-8} (\text{s m}^2 \text{sr GeV/n})^{-1}$


Antiproton sensitivity

- Precision antiproton spectrum in unexplored low-energy range (<0.25 GeV/n): ~500 antiprotons for each long-duration balloon flight
- Validation of technique:
 - Reconstruction of annihilation signature
 - X-rays from exotic atom deexcitation
 - Test models for atmospheric effects→ Reduces the systematic uncertainties for antideuteron search
- Probe light dark matter models and primordial black hole evaporation



Antihelium-3 sensitivity

N. Saffold et al., Astropart. Phys. 130, 102580 (2021)




Finding low-energy antihelium would be truly revolutionary new physics

Time-of-Flight




- Tasks:
 - main trigger system, special antinuclei trigger achieves a manageable rate of ~ 500 Hz (down from 200 kHz individual TOF paddle rate)
 - velocity measurement
- Plastic scintillator (Eljen EJ-200: 160-180cm long, 0.6 cm thick) with SiPMs (Hamamatsu S13360-6050VE)
- fast sampling with DRS4 ASIC: <400 ps timing resolution achieved in test paddles (end-to-end time difference) and in GAPS functional prototype (GFP).






Tracker

- Tracker acts as target and tracking device
- GAPS can accommodate 1,440 4" Si(Li) detectors, 2.5mm thickness (1109 detectors calibrated for first flight)
- Operation at temperature of -35C to -45C, cooling system will use novel OHP approach
- Readout via custom ASIC: integrated low-noise preamplifier with large dynamic range: 10keV to 100MeV
- Publications:
 - Perez et al., NIM A 905, 12 (2018)
 - Kozai et al., NIM A 947, 162695 (2019)
 - Rogers et al., JINST 14, P10009 (2019)
 - Saffold et al., NIM A 997, 165015 (2021)
 - Xiao et al., in preparation (2022)




Oscillating heat pipe cooling system





- passive cooling approach developed at JAXA/ISAS:
- small capillary metal tubes filled with a phase-changing refrigeration liquid
- small vapor bubbles form in the fluid
→ expand in warm sections/contract in cool sections
- rapid expansion and contraction of these bubbles create thermo-contraction hydrodynamic waves that transport heat
- no active pump system is required
- First prototype was flown in 2012 and another prototype was flown from Ft. Sumner in 2019

Okazaki et al., J. Astr.. Instr. 3 (2014)
Fuke et al., J. Astron. Instrum. (2017)
Okazaki et al., Appl. Therm. Eng. (2018)

GAPS Functional Prototype (GFP)



~2.5m





- Prototype: 3 layers of Si(Li) tracker (36 modules): readout with flight ASIC, 2 layers of TOF above
- **Goals:** test and operate all components together, test readout chain, collect X-ray data, collect muon data

Integration of the flight instrument


- Si(Li) modules



- Gondola



- OHP



GAPS balloon payload (under integration)

Timeline

- Integration in fall 2022
- Ground testing in spring 2023
- Thermal vacuum test summer 2023
- **First flight in 2023/24 from McMurdo, Antarctica**




Image credit: NASA (cropped)

GAPS path forward




GAPS team - Oct 2019



- **GAPS will deliver:**
 - a precision antiproton measurement in an unexplored energy range <0.25 GeV/n
 - antideuteron sensitivity 2 orders of magnitude below the current best limits, probing a variety of DM models across a wide mass range
 - the only complementary probe of the AMS-02 antinuclei signal
- GAPS instrument integration is ongoing → **first flight in austral summer 2023**

GAPSimulator AR app



Get it on the PlayStore and App store


→ search for "GAPSimulator"

Developed by UH undergrads: Layne Fujioka, Ben Weiss, Zac Bailey

Event reconstruction


Primary track reconstruction

Y-Z Projection




Annihilation track reconstruction

Y-Z view




$\bar{p}+p$ annihilation at rest



- test of annihilation physics in Geant4 is ongoing
- use antiproton data for validation
- work with Geant4 developers


Trigger design




- main background: protons, alpha, carbon
- High-speed trigger and veto:
 - stopping events deposit more energy (lower beta)
 - annihilation events produce more TOF hits
 - paddle combinations can be used to constrain to zenith angle
- **smart combination reduces trigger rate to be below 500Hz**

Si(Li) detector development


- Lithium is applied to the front surface of B-doped p-type Si and diffused through short depth
- Li atoms donate electrons, resulting in an n-type Si lattice layer and leftover free positive Li ions
- under reverse bias, positive Li ions move away from the n-type region
 - compensate acceptor atoms in the p-type bulk
 - compensate impurities in the Si
- drifting procedure creates a thick compensated region (4.6days at 600V and 100C)
- ultrasonic machining on the n+(Li) contact → guard ring structure, reduces leakage current, much better energy resolution
- electrodes are thermal-evaporated ohmic/blocking contacts
- Passivation is applied




B-doped, p-type substrate wafers



Evaporate and diffuse Li for n+-layer




Form top-hat structure to control drift
Evaporate Ni + Au electrodes




Drift Li through wafer






Form guard ring + strips



Apply polyimide passivation




Tracker qualification



- Single detector test shows the required resolution of the detectors
- Detector module calibration facilities are set up at MIT and UHM
- All modules calibrated

Oscillating heat pipe cooling system



- Most of the Si(Li) detectors are cooled down to lower than -40C . Thermal design will be further optimized to satisfy all detector temperature requirements