



University
of Regina

Superallowed Fermi β Decay

The low-energy precision frontier of nuclear physics



Dr. Gwen Grinyer she/her

Department of Physics, University of Regina, Regina, SK S4S 0A2, Canada



Gwen.Grinyer@uregina.ca



@gwendoesscience

Hi! I'm Gwen!

- My academic journey:
 - B.Sc. McMaster University (2002)
 - M.Sc. University of Guelph (2004)
 - Ph.D. University of Guelph (2008)
 - PDF Michigan State (2008-2010)
 - Staff scientist CEA (2010-2017)
 - Professor U Regina (since 2017)
- My lived experience:
 - Experimental nuclear physicist
 - First generation academic
 - Mom of 3 kids (ages 7, 14, 16)
 - Woman and LGBT in physics
- Passionate about EDI in STEM




@gwendoesscience




Nuclear β decay

- A neutron turns into a proton (or vice versa)



neutron \rightarrow proton




proton \rightarrow neutron

- Momentum conservation & selection rules:

$$\vec{J}_P = \vec{J}_D + \vec{L} + \vec{S} \quad \pi_P = \pi_D (-1)^L$$

- Allowed decays ($L=0$)
- Forbidden decays ($L=1,2,3,\dots$)
- Fermi decays ($S=0$)
- Gamow-Teller decays ($S=1$)



Half-lives and *ft* values

B.Singh et al. Nucl. Data Sheets 84, 487 (1998)

Case	$J^\pi (P \rightarrow D)$	Classification	$T_{1/2}$	Fraction
$^{18}\text{N} \rightarrow ^{18}\text{C}$	$1^- \rightarrow 1^-$	Allowed (GT&F)	624 ms	
$^6\text{He} \rightarrow ^6\text{Li}$	$0^+ \rightarrow 1^+$	Allowed (GT only)	807 ms	64%
$^{10}\text{C} \rightarrow ^{10}\text{B}$	$0^+ \rightarrow 0^+$	Allowed (F only)	19 s	1%
$^{38}\text{Cl} \rightarrow ^{38}\text{Ar}$	$2^- \rightarrow 2^+$	1 st Forbidden	37 min	33%
$^{36}\text{Cl} \rightarrow ^{36}\text{Ar}$	$2^+ \rightarrow 0^+$	2 nd Forbidden	3×10^5 years	1%
$^{40}\text{K} \rightarrow ^{40}\text{Ca}$	$4^- \rightarrow 0^+$	3 rd Forbidden	1×10^9 years	0.1%
$^{50}\text{V} \rightarrow ^{50}\text{Cr}$	$6^+ \rightarrow 2^+$	4 th Forbidden	1×10^{17} years	0.1%

- The *ft* value is a convenient way to characterize nuclear β decay


$$ft = \frac{fT_{1/2}}{BR} = \frac{K}{g^2 |M_{fi}|^2}$$

Half-life

Strength

Nuclear Isotopic Spin (Isospin)

- Introduced by Heisenberg in 1932
 - Protons and neutrons – (iso)spin projections of the “nucleon”




$$t_z(p) = -\frac{1}{2}$$



$$t_z(n) = +\frac{1}{2}$$



- Total isospin (T) and isospin projection T_z of the nucleus



$$T_z = \frac{1}{2}(N - Z) \quad T = |T_z|, |T_z| + 1, \dots, \frac{N + Z}{2}$$

- Nuclear β decay is a neutron changing into a proton (or vice versa)
 - Fermi decay between “isobaric analogue states” is a ladder operator

$$|M_F|^2 = (T \mp T_z)(T \pm T_z + 1)$$

For $T = 1$ decays



$$|M_F|^2 = 2$$

Exact!
(to extent that isospin valid)

Conserved Vector Current Hypothesis (CVC)


- The ft values for superallowed Fermi decays...

$$ft = \frac{fT_{1/2}}{BR} = \frac{K}{g^2 |M_{fi}|^2}$$

strength? ← constants ✓
isospin ✓


- CVC hypothesis (based on analogy to electrodynamics)

- A universal coupling constant – the electric charge “e”



- The weak interaction is also thought to have a universal coupling constant!

$$G_V = 1.13621 \times 10^{-5} \text{ GeV}^{-2}$$




R.P.Feynman and M.Gell-Man PR 109, 193 (1958)

Conserved Vector Current Hypothesis (CVC)

- The ft values for superallowed Fermi decays... **should be constant!**


$$ft = \frac{fT_{1/2}}{BR} = \frac{K}{2G_V^2} = \text{constant } ?$$

- CVC hypothesis (based on analogy to electrodynamics)
 - A universal coupling constant – the electric charge “e”



- The weak interaction is also thought to have a universal coupling constant!

$$G_V = 1.13621 \times 10^{-5} \text{ GeV}^{-2}$$





R.P.Feynman and M.Gell-Man PR 109, 193 (1958)

Superallowed *ft* values

J.C.Hardy and I.S. Towner PRC 102, 045501 (2020)

- World survey of superallowed decays
 - > 220 independent measurements
- Superallowed *ft* values
 - Range from 3040 s to 3100 s (2%)
 - Higher-order effects (theory)




Superallowed *ft* values

J.C.Hardy and I.S. Towner PRC 102, 045501 (2020)

- World survey of superallowed decays
 - > 220 independent measurements
- Superallowed *ft* values
 - Range from 3040 s to 3100 s (2%)
 - Higher-order effects (theory)
- Isospin symmetry is not exact
 - Broken by charge dependent forces

$$|M_F|^2 = 2(1 - \delta_C)$$




Superallowed *ft* values

J.C.Hardy and I.S. Towner PRC 102, 045501 (2020)


- World survey of superallowed decays
 - > 220 independent measurements
- Superallowed *ft* values
 - Range from 3040 s to 3100 s (2%)
 - Higher-order effects (theory)
- Isospin symmetry is not exact
 - Broken by charge dependent forces
- Corrected Ft values
 - Constant at the level of 9×10^{-5}
 - Validation of the CVC hypothesis
 - Strong constraint on “new physics”

$$|M_F|^2 = 2(1 - \delta_C)$$



Cabibbo-Kobayashi-Maskawa (CKM) Matrix

- The CKM matrix plays a central role in the Standard Model
 - It describes *all* quark flavour changing interactions (including β decay)
 - Given that there are 3 quark generations, CKM is a 3x3 matrix



$$\begin{bmatrix} d' \\ s' \\ b' \end{bmatrix} = \begin{bmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{bmatrix} \begin{bmatrix} d \\ s \\ b \end{bmatrix}$$

- In the Standard Model the CKM matrix describes a *unitary* transformation

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$$

CKM Unitarity Test

J.C.Hardy and I.S. Towner PRC 102, 045501 (2020)


- The most precise test of CKM unitarity comes from the *top row*
 - V_{ud} is by far the largest and is obtained precisely from superallowed decays

$$|V_{ud}|^2 = \frac{2912.95 \pm 0.54}{\overline{Ft}}$$

↓
Constants

↗

Average Ft value from 15 superallowed Fermi transitions between ^{10}C and ^{74}Rb



$$\begin{aligned} |V_{ud}| &= 0.97373(31) \\ |V_{us}| &= 0.2243(5) \\ |V_{ub}| &= 0.00394(36) \end{aligned}$$

- Present status of the test of CKM unitarity:



$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9985(6)$$

2.5 σ deviation
from unity!!!

TRIUMF's ISAC Facility





- Canada's National Laboratory for Nuclear and Particle Physics
 - Isotope Separator and Accelerator (ISAC)



High-Precision Half-Life Measurements


- We use a gas proportional counter and a fast tape transport system
 - Implant radioactive isotopes from ISAC onto a tape (collection period)
 - Rapidly move the sample into a gas counter (beta particles ionize the gas)
 - Record the radioactive decay of the sample (exponential decay law)



Half-life of ^{10}C

M.R.Dunlop et al. PRL 116, 172501 (2016)

- Beam of radioactive ^{10}C
 - Intensity $\sim 10^5$ ions/s
- Data from 1 cycle (~ 8 mins)
 - Precision $\pm 0.07\%$
- Total of 550 cycles (4 days)
 - $T_{1/2} = 19.3009(17)$ s




$$A(t) = A_0 e^{-\lambda t} + B$$

Half-life of ^{10}C

M.R.Dunlop et al. PRL 116, 172501 (2016)



- Beam of radioactive ^{10}C
 - Intensity $\sim 10^5$ ions/s
- Data from 1 cycle (~ 8 mins)
 - Precision $\pm 0.07\%$
- Total of 550 cycles (4 days)
 - $T_{1/2} = 19.3009(17)$ s
- Systematic uncertainties
 - The most important part!
- Half-life of ^{10}C @ TRIUMF
 - Overall precision $\pm 0.009\%$
 - Most precise $T_{1/2}$ ever reported!



Half-life of ^{14}O

S.Sharma et al. E. Phys. J. A 58, 83 (2022)


- Beam of radioactive ^{14}O
 - Intensity $\sim 10^5$ ions/s
- Data from 1 cycle (~ 25 mins)
 - Precision $\pm 0.10\%$
- Total of 124 cycles (3 days)
 - $T_{1/2} = 70.6192(76)$ s



Half-life of ^{14}O

S.Sharma et al. E. Phys. J. A 58, 83 (2022)

- Beam of radioactive ^{14}O
 - Intensity $\sim 10^5$ ions/s
- Data from 1 cycle (~ 25 mins)
 - Precision $\pm 0.10\%$
- Total of 124 cycles (3 days)
 - $T_{1/2} = 70.6192(76)$ s
- Half-life of ^{14}O @ TRIUMF
 - Overall precision $\pm 0.010\%$
 - Comparable precision to ^{10}C !
- Article published in April!
 - Shivani Sharma, U of R
 - Now at Sunnybrook Hospital



Next generation: GRIFFIN



- New HPGe γ -ray spectrometer
 - 16 large volume “clover” detectors
 - Fully operational since 2015
- Experiment S1140: Half-life of ^{14}O
 - Statistical precision $\pm 0.03\%$
- Regina students lead the analysis!
 - Ugrad Dhruval Shah
 - M.Sc. Nastaran Saei
 - M.Sc. Jizhong Liu
 - Ph.D. Eric Gyabeng Fuakye
- Experiment S1848: BR of ^{34}Ar
 - New experiment at ISAC!
 - Scheduled in June/July 2023




My newest adventure!

LEGEND

- Large Enriched Germanium Experiment for Neutrinoless Double β decay
 - Deep underground ton-scale detector



- Physics program
 - Search for $0\nu\beta\beta$ decay in ^{76}Ge
 - Are neutrinos their own antiparticle?
 - Baryon asymmetry (matter/antimatter)
 - Lepton number violation (beyond SM)
- Legend 1000 baseline design
 - To probe $0\nu\beta\beta$ with 99.7% discovery CL
 - For a ^{76}Ge half-life $> 10^{28}$ years
 - Background : 1 count per FWHM ton year!
 - Considering SNOLAB as a possible site



LEGEND PCDR arXiv: 2107.11462 (2021)

LEGEND Canada 

Queen's, Regina, SFU, SNOLAB

Thank you so much!

- Superallowed Fermi β Decay
 - The low-energy precision frontier
 - Constrain the Standard Model
 - Demanding test of CKM unitarity
- Experiments at TRIUMF-ISAC
 - The best place for these studies
 - World-leading detectors/expertise
- Many other projects in my group!
 - Looking for students (Fall 2024)
- Please contact me if interested!!



Gwen.Grinyer@uregina.ca



@gwendoesscience

