



Contribution ID: 29

Type: **Talk**

## Effects of an induced three-body force in the incident channel of (d,p) reactions

*Friday 6 September 2019 11:20 (20 minutes)*

### Effects of an induced three-body force in the incident channel of (d,p) reactions

M. J. Dinmore, N. K. Timofeyuk, J. S. Al-Khalili, R. C. Johnson.  
University of Surrey, United Kingdom

A widely accepted practice for treating deuteron breakup in  $A(d,p)B$  reactions relies on solving a three-body  $A+n+p$  Schrodinger equation with pairwise  $A-n$ ,  $A-p$  and  $n-p$  interactions. However, it was shown in [1] that projection of the many-body  $A+2$  wave function into the three-body  $A+n+p$  channel results in a complicated three-body operator that cannot be reduced to a sum of pairwise potentials. It contains explicit contributions from terms that include interactions between the neutron and proton via excitation of the target  $A$ . Such terms are normally neglected. We estimate the first order contribution of these induced three-body terms and show that applying the adiabatic approximation to solving the  $A+n+p$  model results in a simple modification of the two-body nucleon optical potentials. We illustrate the role of these terms for the case of  $^{40}\text{Ca}(d,p)^{41}\text{Ca}$  transfer reactions at incident deuteron energies of 11.8, 20 and 56 MeV, using several parameterisations of nonlocal optical potentials.

[1] R.C. Johnson and N.K. Timofeyuk, Phys. Rev. C 89, 024605 (2014).

**Authors:** Mr DINMORE, Michael (University of Surrey); Dr TIMOFEYUK, Natalia (University of Surrey); Prof. AL-KHALILI, Jim (University of Surrey); Prof. JOHNSON, Ron (University of Surrey)

**Presenter:** Mr DINMORE, Michael (University of Surrey)

**Session Classification:** Parallel Session Friday: Nuclear Reactions

**Track Classification:** Nuclei