

Contribution ID: 90

Type: Talk

## An investigation for the appearance of long range nuclear potential on the ultra low energy nuclear synthesis

Friday 6 September 2019 12:35 (20 minutes)

On the three-body kinematics, the author proposed a general particle transfer (GPT) potential which appears, not only at the three-body break-up threshold, but also at the quasi two-body threshold. The new potential represents a Yukawa-type potential for shorter range, but a  $1/r^n$ -type potential for longer range. The longer range part of the GPT potential for n = 1 indicates an attractive Coulomb-like or a gravitational potential. While, n = 2 includes the Efimov-like potential. The GPT-potential implies the existence of Efimov-like states in the hadron systems. In order to confirm the GPT potential, we investigate the possibility of lanthanum (La)-nucleus creation via the *ultralowenergyreaction*  $Cs(2d,\gamma)La$  on the three-ion quasi-molecule CsD2 in the CsD2Pd12-cluster where D-Cs-D (d-Cs-d) three-body bound states and wave functions are calculated. The wavefunction overlap value (WFO:  $W_{n,m}$ ) between the La highest nuclear excited state with the quantum number n = 5, and the lowest CsD<sub>2</sub> quasi molecular modified states with the quantum number m = 6 is of critical importance for the existence of the electro-magnetic (EM) transition in the Cs(2d, $\gamma$ )La reaction. We found that the  $W_{5,6}$  value is very sensitive to the nuclear potential tail with a two- or three-body  $1/r^2$ -type long range hadron potential. It gives the ratio:  $W_{5,6}^L/W_{5,6}^S \approx 10^7$ , where  $W_{5,6}^L$  is given by the nuclear force with the long range GPT potential, and  $W^S_{5,6}$  is obtained by the usual short range nuclear potential. The ratio corresponds to the E2-transition ratio, where the E1-transition is forbidden in this reaction. This fact means that the quasi molecular state is stable only for the short range nuclear potential, but the nuclear synthesis occurs for the short range nuclear potential with the GPT long range potential. We can conclude that if the reaction occurs, then the GPT potential is confirmed.

Author: Prof. ORYU, Shinsho (Tokyo university of science)

Presenter: Prof. ORYU, Shinsho (Tokyo university of science)

Session Classification: Parallel Session Friday: Atoms and Molecules

Track Classification: Nuclei