



Contribution ID: 36

Type: Talk

Response functions and cross sections for inclusive neutrino scattering off ^2H , ^3H and ^3He

Thursday 5 September 2019 14:30 (20 minutes)

Neutrino scattering on nuclei has been investigated for several decades. It played an important role in establishing fundamentals of the theory of weak interactions and electroweak unification. Later it became a tool to study the much more subtle properties of neutrinos, such as masses and oscillations.

Most of the calculations for neutrino scattering on light nuclei were performed in coordinate space, see important Refs. (1-4). In a recent paper (5) we presented momentum-space based results for several (anti)neutrino induced reactions on light nuclei: $\nu(\bar{\nu}) + ^2\text{H} \rightarrow \nu(\bar{\nu}) + p + n$,

$$\bar{\nu} + ^2\text{H} \rightarrow e^+ + n + n,$$

$$\nu(\bar{\nu}) + ^3\text{He} \rightarrow \nu(\bar{\nu}) + p + ^2\text{H},$$

$$\nu(\bar{\nu}) + ^3\text{He} \rightarrow \nu(\bar{\nu}) + p + p + n,$$

$$\nu(\bar{\nu}) + ^3\text{H} \rightarrow \nu(\bar{\nu}) + p + ^2\text{H},$$

$$\nu(\bar{\nu}) + ^3\text{H} \rightarrow \nu(\bar{\nu}) + n + n + p,$$

$$\nu + ^3\text{He} \rightarrow e^+ + n + ^2\text{H},$$

$$\nu + ^3\text{He} \rightarrow e^+ + n + n + p,$$

$$\nu + ^3\text{H} \rightarrow e^+ + n + n + n. \text{ We restricted ourselves to neutrino energies up to } 300 \text{ MeV and pointed out problems arising from merging}$$

at many incoming neutrino energies. The essential building block for the total cross section at a given (anti)neutrino energy E is the threefold differential cross section $d^3\sigma/d\Omega' dE'$, which depends on E , on the (anti)neutrino scattering angle θ' and on its final energy E' .

Namely, the total cross section is most easily calculated as $\sigma_{tot}(E) = \int d\Omega' \int dE' \frac{d^3\sigma}{d\Omega' dE'} = 2\pi \int d\theta' \sin\theta' \int dE' \frac{d^3\sigma}{d\Omega' dE'}$. The cross section can be expressed in terms of the so-called response functions R_{jk} (5), which depend on two parameters only: on the internal energy of the nuclear system E_{CM} and on the magnitude of the three-momentum transfer Q : $d^3\sigma/d\Omega' dE' = \sum_{jk} v_{jk}(E, \theta', E') R_{jk}(E_{CM}, Q)$. Since the v_{jk} functions, stemming from the lepton matrix element,

are analytically known, it is thus much more efficient to calculate just the response functions on a sufficiently dense two dimensional grid in the (E_{CM}, Q) plane and use interpolations to obtain these values of the response functions which are needed to compute $\sigma_{tot}(E)$ at a required energy E . Note also that in the case of the neutral current driven reactions, exactly the same response functions can be used to obtain cross sections with neutrinos and antineutrinos. This approach has been already successfully tested for the (anti)neutrino reactions on the deuteron and one example of the response function for the $\nu(\bar{\nu}) + ^2\text{H} \rightarrow \nu(\bar{\nu}) + p + n$ reaction is shown in Fig. 1. The corresponding work for several reactions with the three-nucleon systems is in progress and will be reported at the conference. As in Ref. (5), also in the present calculations the single nucleon current from Ref. (6) and the AV18 two-nucleon potential (7) is employed.

!Fig.1 [The R_{00} response function (see Ref. (1) for the definition and more details) for the neutral-current driven process $\nu(\bar{\nu}) + ^2\text{H} \rightarrow \nu(\bar{\nu}) + p + n$]

References

- (1) S. Nakamura, T. Sato, V. Gudkov, and K. Kubodera, Phys. Rev. C **63**, 034617 (2001);
Erratum: [Phys. Rev. C **73**, 049904 (2006)].
- (2) S. Nakamura *et al.*, Nucl. Phys. A **707**, 561 (2002);
<http://www-nuclth.phys.sci.osaka-u.ac.jp/top/Netal/total/index.html>.
- (3) G. Shen, L. E. Marcucci, J. Carlson, S. Gandolfi, and R. Schiavilla, Phys. Rev. C **86**, 035503 (2012).
- (4) A. Baroni and R. Schiavilla, Phys. Rev. C **96**, 014002 (2017).
- (5) J. Golak *et al.*, Phys. Rev. C **98**, 015501 (2018).

(6) J. Golak *et al.*, Phys. Rev. C **90**, 024001 (2014).

(7) R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).

Author: GRASSI, Alessandro (for LENPIC Collaboration) (M. Smoluchowski Institute of Physics, Jagiellonian University, PL-30348 Cracow, Poland)

Presenter: GRASSI, Alessandro (for LENPIC Collaboration) (M. Smoluchowski Institute of Physics, Jagiellonian University, PL-30348 Cracow, Poland)

Session Classification: Parallel Session Thursday: Few-Nucleon Systems

Track Classification: Nuclei