

The $^{39,41,42}\text{Ar}$ nuclides as probes of neutron-induced reactions in a high-density plasma at the National Ignition Facility: a proposed experiment and calibration measurements

Inertial fusion laser-induced implosions at the National Ignition Facility (NIF) are a unique environment to reproduce astrophysical conditions in the laboratory. The laser energy is used to compress and heat a capsule filled with deuterium-tritium fuel to conditions (density, temperature, and pressure) comparable to or exceeding those in the center of stars. Recent experiments at NIF first passed the burning-plasma threshold [1,2], where self-heating exceeded the external heating applied to the fuel and produced record fusion yields of ≈ 1 MJ. Neutrons are produced in a volume with a radius of ~ 50 μm within ~ 100 ps, representing a uniquely high neutron density approaching 10^{22} cm^{-3} close to those of the astrophysical r process and fluxes of 10^{31} $\text{cm}^{-2}\text{s}^{-1}$. In a dedicated NIF high-power laser shot, we plan to investigate the following neutron-induced reactions on ^{40}Ar incorporated in the capsule gas; the chemical inertness of noble gas Ar allows for reliable collection of reaction products. The $^{40}\text{Ar}(n, 2n)^{39}\text{Ar}$ reaction is a direct monitor of the fast-neutron flux and the $^{40}\text{Ar}(n, \gamma)^{41}\text{Ar}$ and $^{40}\text{Ar}(2n, \gamma)^{42}\text{Ar}$ reactions are sensitive to energy downgraded neutrons. The latter reaction is a monitor of extreme neutron densities produced in the process and may provide an indication of the feasibility to study the important $^{58}\text{Fe}(2n, \gamma)^{60}\text{Fe}$ reaction [3] in the laboratory. The long-lived ^{39}Ar ($t_{1/2} = 268$ y) and ^{42}Ar (33 y) nuclides are detected via noble-gas accelerator mass spectrometry at Argonne National Laboratory. We report here on calibration measurements of the total yield of the $^{40}\text{Ar}(n, 2n)^{39}\text{Ar}$ reaction in a 14 MeV neutron activation, investigated for the first time. The neutron activation was performed with the DT neutron generator of Technical University Dresden located at Helmholtz-Zentrum Dresden-Rossendorf. Direct detection of the ^{42}Ar nuclide in a ^{40}Ar sample activated by the slow double-neutron capture reaction $^{40}\text{Ar}(n, \gamma)^{41}\text{Ar}(n, \gamma)^{42}\text{Ar}$ at the high flux reactor of Institut Laue-Langevin was successfully demonstrated for the first time. Preliminary results of these calibration experiments are presented.

Support from the Pazy Foundation (Israel) and USA-Israel Binational Science Foundation is gratefully acknowledged. This work was supported by the U.S. Department of Energy, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357. This research used resources of ANL's ATLAS facility, which is a DOE Office of Science User Facility.

[1] A. B. Zylstra et al., Nature 93, 542 (2022).

[2] A. L. Kritcher et al., Nature Phys. 18, 251 (2022).

[3] W. Wang et al., Astrophys. J. 889, 169 (2020)

Length of presentation requested

Oral presentation: 17 min + 3 min questions

Please select between one and three keywords related to your abstract

Nuclear physics - experimental

2nd keyword (optional)

Nucleosynthesis

3rd keyword (optional)

Instrumentation

Authors: CLARK, Adam; Dr ZYLSTRA, Alex (Lawrence Livermore National Laboratory); Dr VELSKO, Carol (Lawrence Livermore National Laboratory); Prof. PAUL, Michael (The Hebrew University of Jerusalem); Mr NEL-

SON, Austin (University of Notre Dame); DICKERSON, Clayton (Argonne National Lab); Mr HOFFMANN, Hans F. R. (Technical University Dresden); JAYATISSA, Heshani; Dr TOLTSUKHIN, Ivan (Argonne National Laboratory); MCLAIN, Jake; ZUBER, Kai (Technische Universitaet Dresden); REHM, Karl Ernst; Mrs CALLAHAN, Lauren (University of Notre Dame); Mrs PICHOTTA, Marie (Technical University Dresden); AVILA, Melina; Dr TESSLER, Moshe (Soreq Nuclear Research Center); Prof. COLLON, Philippe (University of Notre Dame); PARDO, Richard (Argonne National Laboratory); SCOTT, Robert (Argonne National Laboratory); SCHWENGNER, Ronald (Helmholtz-Zentrum Dresden-Rossendorf); Ms SAHOO, Rudra N. (The Hebrew University of Jerusalem); Mr BAI-LEY, Thomas (University of Notre Dame); Mr DOERING, Toralf (Helmholtz-Zentrum Dresden-Rossendorf); KOESTER, Ulli (Institut Laue-Langevin (FR)); Dr KASHIV, Yoav (University of Notre Dame); VONDRASEK, richard (Argonne National Laboratory)

Presenter: Prof. PAUL, Michael (The Hebrew University of Jerusalem)