



Contribution ID: 161

Type: Talk

Peiwen Wu (Korea Inst. for Advanced Study): Heavy Quark Flavored Scalar Dark Matter

Thursday 22 February 2018 11:37 (14 minutes)

The absence of confirmed signal in dark matter (DM) direct detection (DD) may suggest a weak coupling between DM and the first generation quarks. In this work we consider a real scalar dark matter S which has new Yukawa interactions with charm c and top quark t via a vector-like fermion mediator ψ . By setting the Higgs portal to be negligible, we focus on the new Yukawa interactions. Since there is no valence c, t quark in nucleons, DM-gluon scattering at loop level becomes important. We found that renormalization group equation (RGE) effects are crucial in calculating the DM-nucleon scattering rate at $\mu_{had} \sim 1$ GeV if one constructs the effective theory at $\mu_{EFT} \sim m_Z$. For the perturbative benchmark couplings we choose, combined results from relic abundance requirement $\Omega h^2 = 0.12$, direct/indirect detection constraints, 13 TeV LHC data have excluded a thermal relic DM with $m_S < m_t/2$ in this model. FCNC processes of top quark can be generated at both tree level $t \rightarrow \psi^{(*)} S \rightarrow c S S$ and loop level $t \rightarrow c + \gamma/g/Z$, of which the branching fractions are usually below 10^{-9} after passing the other constraints, which are still safe from the current top quark width measurements.

Presenter: WU, Peiwen (Korea Institute for Advanced Study (KIAS))

Session Classification: Session 7