



Contribution ID: 183

Type: Talk

## Paul Shapiro (U of Texas at Austin): Cosmology in a Universe with Complex Scalar-Field Dark Matter

Thursday 22 February 2018 18:14 (14 minutes)

We consider an alternative to WIMP cold dark matter (CDM), ultralight bosonic dark matter ( $m \geq 10^{-22}$  eV) described by a complex scalar field (SFDM) with global U(1) symmetry, for which the comoving particle number density is conserved after particle production during standard reheating. We allow for a repulsive self-interaction. For complex SFDM, structure formation is CDM-like on large scales but suppressed on small-scales by quantum effects, just as in the case of a real scalar field. However, in the  $\Lambda$ SFDM universe, the background evolution of the complex field differs from that of the real field; complex SFDM starts relativistic, evolving from stiff ( $w=1$ ) to radiationlike ( $w=1/3$ ), before becoming nonrelativistic at late times ( $w=0$ ). Thus, before the familiar radiation-dominated era, there is an earlier era of stiff-SFDM-domination, during which the expansion rate is higher than in  $\Lambda$ CDM. SFDM particle mass  $m$  and coupling strength  $\lambda$ , of a quartic self-interaction, are therefore constrained by cosmological observables, particularly  $N_{\text{eff}}$ , the effective number of neutrino species during BBN, and  $z_{\text{eq}}$ , the redshift of matter-radiation equality. Furthermore, since the stochastic gravitational-wave background (SGWB) from inflation is amplified during the stiff era, it can contribute a radiationlike component large enough to affect these observables by further boosting the expansion rate. Remarkably, this same amplification makes detection of the SGWB possible at high frequencies by current laser interferometer experiments, e.g., aLIGO/Virgo and LISA. For SFDM particle parameters that satisfy these cosmological constraints, the amplified SGWB is detectable by LIGO for a broad range of reheat temperatures  $T_{\text{reheat}}$ , for values of tensor-to-scalar ratio  $r$  currently allowed by CMB polarization measurements. The SGWB is maximally detectable if modes that reentered the horizon when reheating ended have frequencies today in the LIGO sensitive band. For  $r=0.01$ , if SFDM parameters are chosen which marginally satisfy the above constraints, the maximally detectable model for  $(\lambda/(mc^2)^2, m)=(10^{-18}\text{eV}^{-1}\text{cm}^3, 8 \times 10^{-20}\text{eV})$  corresponds to  $T_{\text{reheat}} \approx 10^4$  GeV, for which we predict an aLIGO O1 run detection with  $\text{SNR} \sim 10$ . Upper limits on the SGWB reported by aLIGO O1 already place a new kind of cosmological constraint on SFDM. A wider range of SFDM parameters and  $T_{\text{reheat}}$  should be accessible to aLIGO/Virgo O5. For  $r=0.01$  and  $\lambda/(mc^2)^2=10^{-18}\text{eV}^{-1}\text{cm}^3$ ,  $3\sigma$  detection is predicted for  $600 \leq T_{\text{reheat}}(\text{GeV}) \leq 10^7$  by O5.

**Presenter:** SHAPIRO, Paul (The University of Texas at Austin)

**Session Classification:** Session 10