

## Results from the KDK experiment

*Friday 31 March 2023 17:45 (15 minutes)*

Potassium-40 ( $^{40}\text{K}$ ) is a naturally-occurring radioactive isotope. It is a background in rare-event searches, plays a role in geochronology, and has a nuclear structure of interest to theorists. This radionuclide decays mainly by beta emission to calcium, and by electron-capture to an excited state of argon. The electron-capture decay of  $^{40}\text{K}$  directly to the ground state of argon has never been measured, and predicted intensities are highly variable (0–0.22%). This poorly understood intensity may impact the interpretation of the DAMA claim of dark matter discovery by constraining the signal modulation fraction [1]. The KDK (potassium decay) experiment has carried out the first measurement of this electron-capture branch, using a novel setup at Oak Ridge National Labs [2]. KDK deployed a very sensitive inner detector to trigger on the  $\gamma$ -radiation emitted by both forms of electron capture, surrounded by a very efficient veto to distinguish between the decays to ground state and those to the excited state. We present result of the experiment [3].

[1] Pradler et al, Physics Letters B 720 (2013) 399–404, <http://dx.doi.org/10.1016/j.physletb.2013.02.033>

[2] Stukel et al, Nuclear Inst. and Methods in Physics Research, A 1012 (2021) 165593, <https://doi.org/10.1016/j.nima.2021.165593>

[3] Stukel et al, <https://doi.org/10.48550/arXiv.2211.10319>

**Authors:** DI STEFANO, Philippe; STUKEL, Matthew Jake (Gran Sasso Science Institute); HARIASZ, Lilianna; KDK EXPERIMENT

**Presenter:** DI STEFANO, Philippe

**Session Classification:** SESSION 14: Direct detection: Technical Development-2 (CHAIR: Rafael Lang - Purdue University)