



Contribution ID: 53

Type: not specified

Quark-hadron pasta phases for two-phase approaches and the third family of compact stars

Friday 20 September 2019 09:40 (40 minutes)

The effect of pasta phases on the quark-hadron phase transition is investigated for a set of relativistic mean-field equations of state for both hadron and quark matter. The results of the full numerical solution with pasta phases are compared with those of an interpolating construction used in previous works, for which we demonstrate an adequate description of the numerical results. A one-to-one mapping of the free parameter of the construction to the physical surface tension of the quark-hadron interface is obtained for which a fit formula is given. For each pair of quark and hadron matter models the critical value of the surface tension is determined, above which the phase transition becomes close to the Maxwell construction. This result agrees well with earlier theoretical estimates. The study is extended to neutron star matter in beta equilibrium with electrons and muons and is applied to investigate the effect of pasta phases on the structure of hybrid compact stars and the robustness of a possible third family solution.

[1] K. Maslov et al., Phys. Rev. C 100, 025802 (2019)

Authors: BLASCHKE, David (University of Wroclaw); MASLOV, Konstantin (NRNU MEPhI); GRIGORIAN, Hovik; AYRIYAN, Alexander (JINR & AANL); TATSUMI, Toshitaka (Kyoto U.); MARUYAMA, Toshiki (Japan Atomic Energy Agency); YASUTAKE, Nobutoshi (Chiba Institute of Technology)

Presenter: BLASCHKE, David (University of Wroclaw)