

Contribution ID: 116

Type: Oral presentation

## Time Synchronization Schemes for the Future Hyper-Kamiokande Experiment

Tuesday 20 October 2020 08:20 (20 minutes)

On the strength of a double Nobel prize-winning experiment (Super)Kamiokande and an extremely successful long-baseline neutrino program, the third generation Water Cherenkov detector, Hyper-Kamiokande, is being developed by an international collaboration as a leading worldwide experiment based in Japan. Hyper-Kamiokande will be able to measure with the highest precision the leptonic CP violation, will have excellent capability to search for proton decay and ability to precisely test the three-flavor neutrino oscillation paradigm in conjunction with a strong astrophysical program.

The Hyper-Kamiokande detector will be hosted in the Tochibora mine, about 295 km away from the J-PARC proton accelerator research complex in Tokai, Japan and will be the largest underground water Cherenkov detector with a 68 m diameter and 72 m height cylindrical tank, approximately 8 times larger fiducial volume than Super-Kamiokande. It will be equipped with about 50,000 photo-sensors. Synchronization of the timing of each PMT signal is crucial for precise measurement of the timing of photon arrival. In Hyper-Kamiokande, timing resolution of the photo-sensor is expected to be sub-nanosecond and the jitter less than 100 ps. The CERN designed White Rabbit protocol and a full custom FPGA based solution are under evaluation to reach this objective. This contribution will show the architectural solutions envisaged for the experiment and will focus on the R&D program comparing the solutions under evaluation and showing the first measures conducted on the prototypes.

## Minioral

Yes

## **IEEE Member**

No

## Are you a student?

No

Author: RUSSO, Stefano (CNRS (new))

Presenter: RUSSO, Stefano (CNRS (new))

Session Classification: Oral presentations TRIG01

Track Classification: Data Acquisition System Architectures