

## Measuring hadronic corrections to the muon $g-2$ at BESIII

*Thursday 30 September 2021 11:15 (20 minutes)*

The recent Fermilab result has confirmed the long standing discrepancy between the direct measurement and the Standard Model (SM) prediction of the anomalous magnetic moment of the muon  $a_\mu=(g_\mu-2)/2$ . The possibility to improve the SM prediction using data-driven approaches has motivated the BESIII collaboration to embark on a dedicated experimental program. The high statistics data samples collected with the BESIII experiment in  $e^+e^-$  collisions in the tau-charm region are analyzed exploiting the initial state radiation technique in order to measure hadronic cross sections needed in the dispersive analysis of the hadronic vacuum polarization contribution to  $a_\mu$ . The same data enable investigations of two-photon collisions. These allow the determination of the momentum dependence of transition form factors of light mesons in the relevant kinematic region, which dominate the hadronic Light-by-Light contribution to  $a_\mu$ . The current status and ongoing investigations will be discussed.

### What is your topic?

Anomalous Magnetic Moment of the muon

**Authors:** LIU, Bei Jiang; DENIG, Achim (Univ. Mainz)

**Presenters:** DENIG, Achim (Univ. Mainz); DENIG, Achim (Universität Mainz)

**Session Classification:** Session 6:  $g-2$

**Track Classification:** Tau2021 Abstracts