

Precision measurement of the $B[Y(3S) \rightarrow \tau^+\tau^-] / B[Y(3S) \rightarrow \mu^+\mu^-]$ ratio at BABAR

Tuesday 28 September 2021 13:40 (20 minutes)

We report on a precision measurement of the ratio $R^{\Upsilon(3S)}_{\tau\mu} = B[\Upsilon(3S) \rightarrow \tau^+\tau^-] / B[\Upsilon(3S) \rightarrow \mu^+\mu^-]$ using data collected with the BaBar detector at the SLAC PEP-II e^+e^- collider. The measurement is based on a 28 fb^{-1} data sample collected at a center-of-mass energy of 10.355 GeV corresponding to a sample of 122 million $\Upsilon(3S)$ mesons. The ratio is measured to be $R^{\Upsilon(3S)}_{\tau\mu} = 0.966 \pm 0.008 \text{ (stat)} \pm 0.014 \text{ (syst)}$ and is in agreement with the Standard Model prediction of 0.9948 within 2 standard deviations. The uncertainty in $R^{\Upsilon(3S)}_{\tau\mu}$ is almost an order of magnitude smaller than the only previous measurement.

What is your topic?

Lepton universality and flavour violation

Authors: LUSIANI, Alberto (Scuola Normale Superiore and INFN, sezione di Pisa); SIBIDANOV, Alexei (University of Victoria)

Presenter: SIBIDANOV, Alexei (University of Victoria)

Session Classification: Session 2c: Test of fundamental symmetries with tau lepton

Track Classification: Tau2021 Abstracts