

# Top2015 - 8th International Workshop on Top Quark Physics

Sunday 13 September 2015 - Friday 18 September 2015

Continental Hotel, Ischia



## Book of Abstracts



# Contents

Welcome . . . . .	1
Theory keynote . . . . .	1
Status of the LHC machine . . . . .	1
Status and performances of physics objects at 13 TeV: CMS . . . . .	1
Status and performances of physics objects at 13 TeV: ATLAS . . . . .	1
Top quark physics in the forward region: LHCb measurement and prospects for Run II . . . . .	1
Standard Model results (excl. top) from ATLAS at 13 TeV . . . . .	1
Standard Model results (excl. top) from CMS at 13 TeV . . . . .	1
Searches for new physics at 13 TeV: ATLAS . . . . .	2
Searches for new physics at 13 TeV: CMS . . . . .	2
Top reconstruction and boosted top: theoretical overview . . . . .	2
Top reconstruction and boosted top: experimental overview . . . . .	2
Experimental results using boosted top and novel top reconstruction tools . . . . .	2
TBD . . . . .	2
Latest theory developments for top pair production, generators and showering . . . . .	2
Inclusive top pair production at 7, 8 and 13 TeV in CMS . . . . .	3
Inclusive top pair production at 7, 8 and 13 TeV in ATLAS . . . . .	3
Theory overview of recent progress on single-top production predictions and tools . . . . .	3
Single top production measurements at the LHC: t-channel . . . . .	3
Studies of single-top event properties . . . . .	3
Search for new physics with single top . . . . .	3
Spin measurements in top-quark events at the LHC . . . . .	4
Theory overview of top-quark properties . . . . .	4

Asymmetry measurements in top-quark events at the LHC . . . . .	4
Top production and properties (except mass) at the Tevatron . . . . .	4
ttbar+Boson (except Higgs) . . . . .	4
ttbar+jets/b-bbar . . . . .	4
Top-quark mass measurements at the LHC: standard methods . . . . .	4
Top-quark mass measurements at the LHC: alternative methods . . . . .	4
Top-quark mass measurements at the Tevatron (incl. Tevatron combination) . . . . .	5
Statistical and systematic treatment issues in top mass combinations . . . . .	5
Other combinations than top quark mass measurements at the LHC . . . . .	5
Combination of top quark mass measurements at the LHC . . . . .	5
Theory developments in differential top-quark distributions . . . . .	5
Electroweak corrections in top physics (1) . . . . .	5
Electroweak corrections in top physics (2) . . . . .	6
Theory overview for ttH and tH production . . . . .	6
Search for ttH and tH production with H->bb . . . . .	6
Search for ttH and tH production (other channels) . . . . .	6
Theory overview of BSM top and Higgs interactions . . . . .	6
Search for vector-like quark with top quarks . . . . .	6
Search for tHq production . . . . .	6
Search for new resonances with top quarks . . . . .	6
Searches for vector-like quarks with top quarks . . . . .	7
Searches for SUSY/dark matter with top quarks . . . . .	7
Searches for FCNC with top quarks . . . . .	7
Search for exotica with top quark . . . . .	7
Perspectives for top physics at high luminosity LHC . . . . .	7
Perspectives for top physics at new colliders . . . . .	7
Organization of TOP2016 . . . . .	7
Theory summary . . . . .	8
Experimental summary . . . . .	8
High Pt top and tail modeling (fixed order, resummation) . . . . .	8

PDF . . . . .	8
Generators for top physics and showering . . . . .	8
Young Scientist Forum . . . . .	8
Differential Cross-Section Measurements of Top Quark Pair Production at 8 TeV using the ATLAS Detector . . . . .	8
Measurement of the top quark pair production cross section at 13 TeV with the CMS detector . . . . .	9
Measurement of the inclusive $t\bar{t}$ production cross section in $p\bar{p}$ collisions at 1.96 TeV and determination of the top quark pole mass . . . . .	9
Measuring polarizations of bottom, charm, strange, up and down quarks in top decays . . . . .	9
Measurement of the t-channel single-top quark production cross section at 13 TeV with the CMS detector . . . . .	9
Searches for flavour-changing neutral current top quark decays $t$ to $Hq$ in $pp$ collisions with the ATLAS detector . . . . .	9
Differential cross section measurements (as a function of kinematics variables) . . . . .	9
Differential measurements, II . . . . .	10
Differential cross section measurements at the LHC (as a function of kinematics variables) . . . . .	10
Differential cross section measurements at the LHC (final state objects: gap fraction, vs $n$ jets,... including $t\bar{t}+HF$ ) . . . . .	10
Interpretation of top-quark mass measurements: a theory overview . . . . .	10
Implication of the top (and Higgs) mass to the vacuum stability . . . . .	10
Precise top mass determination using lepton distribution at LHC . . . . .	10
Top production and properties (except mass) at the Tevatron . . . . .	11
Spin measurements in top-quark events at the LHC . . . . .	11
Asymmetry measurements in top-quark events at the LHC . . . . .	11
Single top production measurements at the LHC: other channels . . . . .	11
Status of the LHC machine . . . . .	11
$t\gamma$ , $tH$ and $tZ$ Production Through FCNCs. . . . .	11
Towards NLO QCD radiative corrections for the associated production of heavy quarks and vector bosons . . . . .	12
Associated production of single top plus Higgs and $W$ at NLO-QCD accuracy . . . . .	12
Measurement of top quark polarization with the D0 detector . . . . .	13
Poster and YSF award winners announced . . . . .	13

Announcement of poster awards winners . . . . .	13
Measurement of the $t\bar{t}b\bar{b}$ cross section and the ratio $\sigma(t\bar{t}b\bar{b})/\sigma(t\bar{t}j\bar{j})$ in the lepton+jets final state at $\sqrt{s}=8$ TeV with the CMS experiment . . . . .	13
Measurement of associated top-quark-pair and b-jet production at CMS . . . . .	13
Inclusive and differential measurements of the $t\bar{t}$ charge asymmetry at 8 TeV with the CMS experiment . . . . .	14
Using the Standard Model Effective Field Theory to Search for BSM Physics in the Top Sector . . . . .	14
Measurement of the differential top quark pair production cross section in pp collisions at 8 TeV . . . . .	15
Measurement of the top-quark mass . . . . .	15
$t\bar{t}W$ and $t\bar{t}Z$ production cross sections in leptonic final states in 8 TeV pp collisions at ATLAS . . . . .	15
Search for single top-quark production via flavour-changing neutral currents at 8 TeV with the ATLAS detector . . . . .	15
Measurement of the charge asymmetry in top quark pair production in 8 TeV pp collision data collected by the ATLAS experiment . . . . .	16
Measurement of colour flow with the jet pull angle in $t\bar{t}b\bar{a}$ events using the ATLAS detector . . . . .	16
Measurement of $t\bar{t}b\bar{a}$ production with a veto on additional central jet activity in pp collisions at $\sqrt{s}=8$ TeV using the ATLAS detector . . . . .	17
Measurement of $t\bar{t}$ differential cross-section for highly boosted top quarks produced at ATLAS in proton proton collisions at $\sqrt{s} = 8$ TeV . . . . .	17
Top Tagging at ATLAS . . . . .	18
Studies of top quark pair modelling using ATLAS measurements . . . . .	18
First cross section measurements of $t\bar{t}$ pairs at $\sqrt{s} = 13$ TeV in the same-flavour dilepton events . . . . .	19
Unravelling the non-standard top and Higgs couplings in associated top-Higgs production at the High-luminosity LHC . . . . .	19
search for single top quark production in the $s$ -channel by the CMS at 8 TeV . . . . .	19
First measurement of the differential cross section for $t\bar{t}b\bar{a}$ production in the dilepton final state at a center-of-mass energy of 13 TeV . . . . .	20
Constraining QCD multijet background in single top t-channel production at 13 TeV . . . . .	20

**Session 1: Welcome and keynote / 1**

**Welcome**

**Corresponding Authors:** fabrizio.margaroli@cern.ch, luca.lista@cern.ch

**Session 1: Welcome and keynote / 2**

**Theory keynote**

**Corresponding Author:** mpeskin@slac.stanford.edu

3

**Status of the LHC machine**

**Corresponding Author:** mike.lamont@cern.ch

**Session 2': Status of LHC, physics objects and performances at 13 TeV / 4**

**Status and performances of physics objects at 13 TeV: CMS**

**Session 2': Status of LHC, physics objects and performances at 13 TeV / 5**

**Status and performances of physics objects at 13 TeV: ATLAS**

**Corresponding Author:** imma.riu@cern.ch

6

**Top quark physics in the forward region: LHCb measurement and prospects for Run II**

**Session 3: First physics results from run II at 13 TeV (non-top phys. relevant for top) / 7**

**Standard Model results (excl. top) from ATLAS at 13 TeV**

**Corresponding Author:** paul.james.laycock@cern.ch

**Session 3: First physics results from run II at 13 TeV (non-top phys. relevant for top) / 8**

## **Standard Model results (excl. top) from CMS at 13 TeV**

**Corresponding Author:** [jesus.manuel.vizan.garcia@cern.ch](mailto:jesus.manuel.vizan.garcia@cern.ch)

**Session 3: First physics results from run II at 13 TeV (non-top phys. relevant for top) / 9**

## **Searches for new physics at 13 TeV: ATLAS**

**Corresponding Author:** [jlove@anl.gov](mailto:jlove@anl.gov)

**Session 3: First physics results from run II at 13 TeV (non-top phys. relevant for top) / 10**

## **Searches for new physics at 13 TeV: CMS**

**Corresponding Author:** [jandrea@cern.ch](mailto:jandrea@cern.ch)

**Session 4: Top reconstruction tools, incl. boosted top / 11**

## **Top reconstruction and boosted top: theoretical overview**

**Corresponding Author:** [michael.spannowsky@cern.ch](mailto:michael.spannowsky@cern.ch)

**Session 4: Top reconstruction tools, incl. boosted top / 12**

## **Top reconstruction and boosted top: experimental overview**

**Corresponding Author:** [louise.skinnari@cern.ch](mailto:louise.skinnari@cern.ch)

13

## **Experimental results using boosted top and novel top reconstruction tools**

14

**TBD**



**Session 5: Top-pair cross section / 15**

## **Latest theory developments for top pair production, generators and showering**

**Corresponding Author:** emanuele.re@lapth.cnrs.fr

**Session 5: Top-pair cross section / 16**

## **Inclusive top pair production at 7, 8 and 13 TeV in CMS**

**Corresponding Author:** shannonrebecca.walchcrucy@ugent.be

**Session 5: Top-pair cross section / 17**

## **Inclusive top pair production at 7, 8 and 13 TeV in ATLAS**

**Corresponding Author:** julian.glatzer@cern.ch

**Session 7: Single top / 18**

## **Theory overview of recent progress on single-top production predictions and tools**

**Corresponding Author:** andrew.papanastasiou@desy.de

**Session 7: Single top / 19**

## **Single top production measurements at the LHC: t-channel**

**Corresponding Author:** matthias.komm@cern.ch

20

## **Studies of single-top event properties**

21

## **Search for new physics with single top**

22

## **Spin measurements in top-quark events at the LHC**

Session 6: Top-quark properties (incl. single top) / 23

### **Theory overview of top-quark properties**

Corresponding Author: markus.schulze@cern.ch

24

## **Asymmetry measurements in top-quark events at the LHC**

25

## **Top production and properties (except mass) at the Tevatron**

Session 6': Top-quark properties (incl. single top) / 26

### **$t\bar{t}$ +Boson (except Higgs)**

Corresponding Author: sjolin@physto.se

27

### **$t\bar{t}$ +jets/ $b$ - $b\bar{b}$**

Session 9: Top mass / 28

## **Top-quark mass measurements at the LHC: standard methods**

Corresponding Author: andrea.castro@cern.ch

Session 9: Top mass / 29

## **Top-quark mass measurements at the LHC: alternative methods**

**Corresponding Author:** marcel.vos@cern.ch

**Session 9: Top mass / 30**

## **Top-quark mass measurements at the Tevatron (incl. Tevatron combination)**

**Session 10: Top mass and combinations / 31**

## **Statistical and systematic treatment issues in top mass combinations**

**Corresponding Author:** andreas.alexander.maier@cern.ch

**Session 10: Top mass and combinations / 32**

## **Other combinations than top quark mass measurements at the LHC**

**Corresponding Author:** mara.senghi@ciemat.es

33

## **Combination of top quark mass measurements at the LHC**

**Session 14: Differential cross sections mini-workshop / 34**

## **Theory developments in differential top-quark distributions**

**Corresponding Author:** dheymes@physik.rwth-aachen.de

**Session 14: Differential cross sections mini-workshop / 35**

## **Electroweak corrections in top physics (1)**

**Corresponding Author:** peter.uwer@physik.hu-berlin.de

**Session 14: Differential cross sections mini-workshop / 36**

## **Electroweak corrections in top physics (2)**

**Corresponding Author:** [davide.pagani@cern.ch](mailto:davide.pagani@cern.ch)

**Session 11: Top and Higgs in the SM / 37**

## **Theory overview for ttH and tH production**

**Corresponding Author:** [marco.zaro@lpthe.jussieu.fr](mailto:marco.zaro@lpthe.jussieu.fr)

**Session 11: Top and Higgs in the SM / 38**

## **Search for ttH and tH production with H->bb**

**Corresponding Author:** [darren.michael.puigh@cern.ch](mailto:darren.michael.puigh@cern.ch)

**Session 11: Top and Higgs in the SM / 39**

## **Search for ttH and tH production (other channels)**

**Corresponding Author:** [mcfayden@cern.ch](mailto:mcfayden@cern.ch)

**Session 12: Top and Higgs boson beyond SM / 40**

## **Theory overview of BSM top and Higgs interactions**

**Corresponding Author:** [aldo.deandrea@cern.ch](mailto:aldo.deandrea@cern.ch)

41

## **Search for vector-like quark with top quarks**

42

## **Search for tHq production**

**Session 12: Top and Higgs boson beyond SM / 43**

## **Search for new resonances with top quarks**

**Corresponding Author:** min.suk.kim@cern.ch

**Session 13: Top and Exotica/SuSy / 44**

## **Searches for vector-like quarks with top quarks**

**Corresponding Author:** jeff.tseng@physics.ox.ac.uk

**Session 13: Top and Exotica/SuSy / 45**

## **Searches for SUSY/dark matter with top quarks**

**Corresponding Author:** sophio.patarai@cern.ch

**Session 13: Top and Exotica/SuSy / 46**

## **Searches for FCNC with top quarks**

**Corresponding Author:** kirill.skovpen@cern.ch

47

## **Search for exotica with top quark**

**Session 15: Perspectives / 48**

## **Perspectives for top physics at high luminosity LHC**

**Session 15: Perspectives / 49**

## **Perspectives for top physics at new colliders**

**Corresponding Author:** michelangelo.mangano@cern.ch

**Session 16: Organization of TOP2016 and summary / 50**

## **Organization of TOP2016**

**Session 16: Organization of TOP2016 and summary / 51**

## **Theory summary**

**Corresponding Author:** paolo.nason@cern.ch

**Session 16: Organization of TOP2016 and summary / 52**

## **Experimental summary**

**Corresponding Author:** martijn.mulders@cern.ch

**Session 14: Differential cross sections mini-workshop / 53**

## **High Pt top and tail modeling (fixed order, resummation)**

**Session 14: Differential cross sections mini-workshop / 54**

## **PDF**

**Corresponding Author:** thorne@hep.ucl.ac.uk

55

## **Generators for top physics and showering**

56

## **Young Scientist Forum**

**Session 8: Young Scientists Forum / 57**

## **Differential Cross-Section Measurements of Top Quark Pair Production at 8 TeV using the ATLAS Detector**

Corresponding Author: benjamin.tannenwald@cern.ch

Session 8: Young Scientists Forum / 58

## **Measurement of the top quark pair production cross section at 13 TeV with the CMS detector**

Corresponding Author: till.michael.arndt@cern.ch

Session 8: Young Scientists Forum / 59

## **Measurement of the inclusive $t\bar{t}$ production cross section in $p\bar{p}$ collisions at 1.96 TeV and determination of the top quark pole mass**

Corresponding Author: jiri.franc@fjfi.cvut.cz

Session 8: Young Scientists Forum / 60

## **Measuring polarizations of bottom, charm, strange, up and down quarks in top decays**

Corresponding Author: yevgeny.kats@gmail.com

Session 8: Young Scientists Forum / 61

## **Measurement of the t-channel single-top quark production cross section at 13 TeV with the CMS detector**

Corresponding Author: nils.faltermann@cern.ch

Session 8: Young Scientists Forum / 62

## **Searches for flavour-changing neutral current top quark decays $t$ to $Hq$ in $pp$ collisions with the ATLAS detector**

Corresponding Author: shota.tsiskaridze@cern.ch

63

**Differential cross section measurements (as a function of kinematics variables)**

65

**Differential measurements, II**

Session 5: Top-pair cross section / 66

**Differential cross section measurements at the LHC (as a function of kinematics variables)**

Corresponding Author: otto.heinz.hindrichs@cern.ch

Session 5: Top-pair cross section / 67

**Differential cross section measurements at the LHC (final state objects: gap fraction, vs njets,... including tt+HF)**

Corresponding Author: matthias.danninger@cern.ch

Session 10: Top mass and combinations / 68

**Interpretation of top-quark mass measurements: a theory overview**

Corresponding Author: gennaro.corcella@lnf.infn.it

Session 12: Top and Higgs boson beyond SM / 69

**Implication of the top (and Higgs) mass to the vacuum stability**

Session 8: Young Scientists Forum / 70

**Precise top mass determination using lepton distribution at LHC**



**Corresponding Author:** skawabata@seoultech.ac.kr

**Session 6: Top-quark properties (incl. single top) / 71**

## **Top production and properties (except mass) at the Tevatron**

**Corresponding Author:** giorgio.chiarelli@pi.infn.it

**Session 6': Top-quark properties (incl. single top) / 72**

## **Spin measurements in top-quark events at the LHC**

**Corresponding Author:** linacre@fnal.gov

**Session 6': Top-quark properties (incl. single top) / 73**

## **Asymmetry measurements in top-quark events at the LHC**

**Corresponding Author:** schwarz@umich.edu

**Session 7: Single top / 74**

## **Single top production measurements at the LHC: other channels**

**Corresponding Author:** carolina.gabaldon.ruiz@cern.ch

**Session 2: Status of LHC, physics objects and performances at 13 TeV / 75**

## **Status of the LHC machine**

**Corresponding Author:** mike.lamont@cern.ch

76

## **$t\gamma$ , $tH$ and $tZ$ Production Through FCNCs.**

**Author:** Artur Amorim<sup>1</sup>

**Co-authors:** Jose Santiago<sup>2</sup>; Juan Pedro Araque Espinosa<sup>3</sup>; Nuno Filipe Castro<sup>4</sup>; Rui Alberto Serra Ribeiro Dos Santos<sup>5</sup>

<sup>1</sup> *University of Porto*

<sup>2</sup> *Granada University*

<sup>3</sup> *LIP Laboratorio de Instrumentacao e Fisica Experimental de Part*

<sup>4</sup> *LIP and University of Porto (PT)*

<sup>5</sup> *ISEL Instituto Superior de Engenharia (PT)*

We study the  $t\gamma$ ,  $tH$  and  $tZ$  production through Flavour Changing Neutral Currents (FCNCs) anomalous couplings, in particular how the distributions of physical observables depend on the anomalous couplings. We find that the angular distributions of the decay products depend on the chirality of the anomalous coupling and that the kinematic distributions of the top quark, photon, Higgs boson and Z boson change according to the nature of the anomalous coupling. We also study the expected sensitivity of the LHC experiments to the  $tZ$  production via FCNC at the run-2.

80

## Towards NLO QCD radiative corrections for the associated production of heavy quarks and vector bosons

**Author:** Felix Anger<sup>1</sup>

**Co-author:** Harald Ita<sup>2</sup>

<sup>1</sup> *Albert-Ludwigs-Universität Freiburg*

<sup>2</sup> *Universität Freiburg*

**Corresponding Author:** felix.anger@physik.uni-freiburg.de

The production of vector bosons in association with heavy quarks constitutes an irreducible background to many processes relevant for the ongoing physics program at the LHC, for both SM measurements and BSM searches. In order to provide NLO QCD radiative corrections to these processes, we employ the framework of generalized unitarity, where loop amplitudes are constructed by combining tree amplitudes. The latter are generated by the use of a Berends-Giele off-shell recursion. With this set of methods, we aim to extend existing predictions for the associated production of heavy quarks and vector bosons to higher multiplicities. Here we report recent progress with this process class.

83

## Associated production of single top plus Higgs and W at NLO-QCD accuracy

**Authors:** Benedikt Maier<sup>1</sup>; Federico Demartin<sup>2</sup>

**Co-authors:** Fabio Maltoni<sup>3</sup>; Marco Zaro<sup>4</sup>

<sup>1</sup> *KIT - Karlsruhe Institute of Technology (DE)*

<sup>2</sup> *Universite Catholique de Louvain*

<sup>3</sup> *Universite Catholique de Louvain (UCL) (BE)*

<sup>4</sup> *LPTHE Jussieu, Paris*

**Corresponding Author:** federico.demartin@uclouvain.be

Like single top plus W production, associated production of a top quark together with a Higgs and a W boson ( $tHW$ ) at the LHC is well defined only at LO in QCD and in the 5-flavour scheme. NLO-QCD corrections to the  $tHW$  process feature an overlap with the associated production of a top quark pair and a Higgs boson ( $t\bar{t}H$ ).

I present results from various techniques that can be employed to separate the ttH contribution from the other NLO corrections to tHW (diagram removal, diagram subtraction) and discuss under which conditions this separation can provide sensible predictions, both for the total cross section and for differential distributions.

**Session 8: Young Scientists Forum / 84**

## Measurement of top quark polarization with the D0 detector

**Corresponding Author:** kamil.augsten@fjfi.cvut.cz

85

## Poster and YSF award winners announced

**Session 8: Young Scientists Forum / 86**

## Announcement of poster awards winners

90

## Measurement of the ttbb cross section and the ratio $\sigma(\text{ttbb})/\sigma(\text{ttjj})$ in the lepton+jets final state at $\sqrt{s}=8$ TeV with the CMS experiment

**Author:** Tobias Anton Verlage<sup>1</sup>

<sup>1</sup> Rheinisch-Westfaelische Tech. Hoch. (DE)

**Corresponding Author:** tobias.verlage@cern.ch

The associated production of a top-quark pair with at least two additional jets is studied in the single lepton+jets final state for an integrated luminosity of 19.6/fb at 8 TeV. The jets from the top-quark decay are identified using a constrained kinematic fit and multivariate classifiers in different categories split by the jet multiplicity. The contributions of ttjj and ttbb with at least two additional jets of any flavor or b jets to the inclusive top-quark pair production are extracted with a simultaneous template fit of b-tag discriminants. The measured cross sections  $\sigma(\text{ttbb}) = 271.0 \pm 103.0(\text{stat}) \pm 32.2(\text{syst}) \pm 7.0(\text{lumi})$  fb,  $\sigma(\text{ttjj}) = 23.1 \pm 2.3(\text{stat}) \pm 2.9(\text{syst}) \pm 0.6(\text{lumi})$  pb and the cross section ratio  $\sigma(\text{ttbb})/\sigma(\text{ttjj}) = 0.0117 \pm 0.0040(\text{stat}) \pm 0.0003(\text{syst})$  correspond to the phase space of the additional jets to the top-quark pair defined at generator-level as:  $p_t > 40$  GeV/c,  $|\eta| < 2.5$ ,  $\Delta R > 0.5$ . The flavor of the generated jets is defined by the flavor of the leading quark in the jet. The obtained results are in good agreement with NLO calculations and the CMS measurement in the dilepton channel when an appropriate jet definition is used.

91

## Measurement of associated top-quark-pair and b-jet production at CMS

**Author:** Tae Jeong Kim<sup>1</sup>

**Co-author:** Nazar Bartosik<sup>2</sup>

<sup>1</sup> *Chonbuk National University (KR)*

<sup>2</sup> *Deutsches Elektronen-Synchrotron Hamburg and Zeuthen (DE)*

**Corresponding Author:** nazar.bartosik@cern.ch

Presented is the measurement of  $\sigma(\text{t}\bar{\text{t}}\text{b}\bar{\text{b}})$  and  $\sigma(\text{t}\bar{\text{t}}\text{jj})$ , as well as their ratio, using pp collisions at centre-of-mass energy of 8 TeV recorded with the CMS detector at the LHC. The data used in the measurement corresponds to the integrated luminosity of 19.6 inverse femtobarn. The measurement is performed in the dileptonic final state of the  $\text{t}\bar{\text{t}}$  system for two thresholds of additional jet transverse momentum: 20 GeV and 40 GeV. The measured ratio for  $p_T > 40$  GeV is higher but compatible with a theoretical NLO prediction within 1.6 standard deviations.

92

## Inclusive and differential measurements of the $t\bar{t}$ charge asymmetry at 8 TeV with the CMS experiment

**Author:** Thorsten Chwalek<sup>1</sup>

<sup>1</sup> *KIT - Karlsruhe Institute of Technology (DE)*

**Corresponding Author:** thorsten.chwalek@cern.ch

The  $\text{t}\bar{\text{t}}$  charge asymmetry is measured in proton-proton collisions at a centre-of-mass energy of 8 TeV. The data, collected with the CMS experiment at the LHC, correspond to an integrated luminosity of 19.7/fb. Selected events contain an electron or a muon and four or more jets, where at least one jet is identified as originating from b-quark hadronization. The  $\text{t}\bar{\text{t}}$  charge asymmetry is measured inclusively and differentially as a function of rapidity, transverse momentum, and invariant mass of the  $\text{t}\bar{\text{t}}$  system. For the first time at the LHC, the measurements are also performed in a reduced fiducial phase space of top quark pair production.

95

## Using the Standard Model Effective Field Theory to Search for BSM Physics in the Top Sector

**Author:** Liam Moore<sup>1</sup>

<sup>1</sup> *University of Glasgow*

**Corresponding Author:** l.moore.1@research.gla.ac.uk

This poster will summarize some results from the Standard Model Effective Field Theory (EFT), in particular, how dimension-six operators can modify top quark observables and affect the fundamental parameters of the SM. Results will be shown from a recent global fit of the Wilson Coefficients affecting top-pair and single-top production to current collider data using novel fast-fitting methods. Finally, there will be some information on progress towards a more detailed analysis, including efforts to incorporate NLO QCD corrections to the higher dimensional operators.

96

## Measurement of the differential top quark pair production cross section in pp collisions at 8 TeV

**Author:** Ievgen Korol<sup>1</sup>

<sup>1</sup> *Deutsches Elektronen-Synchrotron Hamburg and Zeuthen (DE)*

**Corresponding Author:** ievgen.korol@cern.ch

Normalized differential top quark pair production cross sections are measured in  $p\bar{p}$  collisions at a centre-of-mass energy of 8 TeV at the LHC using the CMS detector. The dataset used for these measurements corresponds to an integrated luminosity of  $19.7 \text{ fb}^{-1}$ . The measurements are performed in the lepton+jets (e+jets and  $\mu$ +jets) and in the dilepton (ee,  $\mu\mu$ , and  $e\mu$ ) decay channels. The  $t\bar{t}$  production cross section is measured as a function of kinematic properties of the charged leptons, the jets associated to  $b$  quarks, the top quarks, and the  $t\bar{t}$  system. The data are compared with several predictions from perturbative QCD calculations up to approximate next-to-next-to-leading-order precision. No significant deviations are observed relative to the standard model predictions.

97

## Measurement of the top-quark mass

**Author:** Markus Seidel<sup>1</sup>

<sup>1</sup> *CERN*

**Corresponding Author:** markus.seidel@cern.ch

The mass of the top quark is measured using a sample of  $t\bar{t}$  candidate events collected by CMS in pp collisions at  $\sqrt{s}=8$  TeV at the LHC, corresponding to an integrated luminosity of up to  $19.7/\text{fb}$ . This poster presents the results in the lepton+jets, all-jets, and dilepton channels, and their combination.

98

## $t\bar{t}W$ and $t\bar{t}Z$ production cross sections in leptonic final states in 8 TeV pp collisions at ATLAS

**Author:** Carl Joseph Edmund Suster<sup>1</sup>

<sup>1</sup> *University of Sydney (AU)*

**Corresponding Author:** carl.suster@cern.ch

A measurement of the production cross section of a pair of top quarks in association with a  $W$  or  $Z$  boson is presented, using  $20.3 \text{ fb}^{-1}$  of  $pp$  collision data collected by the ATLAS detector at  $\sqrt{s} = 8$  TeV. The measurement combines four separate analyses which consider final states with two opposite sign, two same sign, three and four leptons. A simultaneous fit to data of the  $t\bar{t}W$  and  $t\bar{t}Z$  signals measures their cross sections to be  $\sigma_{t\bar{t}Z} = 176^{+58}_{-52} \text{ fb}$  and  $\sigma_{t\bar{t}W} = 369^{+100}_{-91} \text{ fb}$ , consistent with next-to-leading order theoretical calculations. Tested against the background-only hypothesis, these correspond to an observed (expected) significance of  $4.2\sigma$  ( $4.5\sigma$ ) for  $t\bar{t}Z$  and  $5.0\sigma$  ( $3.2\sigma$ ) for  $t\bar{t}W$ .

99

## Search for single top-quark production via flavour-changing neutral currents at 8 TeV with the ATLAS detector

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Dominic Hirschebuehl<sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *Bergische Universitaet Wuppertal (DE)*

**Corresponding Author:** lroos@lpnhe.in2p3.fr

This poster presents a search for single top-quark production via flavour-changing neutral current processes from gluon plus up- or charm-quark initial states in proton-proton collisions at the LHC. Data collected with the ATLAS detector at a centre-of-mass energy of 8 TeV and corresponding to an integrated luminosity of 20.3 fb<sup>-1</sup> are used. Candidate events for a top quark decaying into a lepton, a neutrino and a jet are selected and classified into signal- and background-like candidates using a neural network. No signal is observed and an upper limit on the production cross-section multiplied by the  $t \rightarrow Wb$  branching fraction is set. The observed 95% CL limit is  $\sigma_{qg \rightarrow t} \times BR(t \rightarrow Wb) < 3.4$  pb. The observed limit can be interpreted as upper limits on the coupling constants of the flavour-changing neutral current interactions divided by the scale of new physics  $\kappa_{ugt}/\Lambda < 5.8 \cdot 10^{-3} \text{TeV}^{-1}$  and  $\kappa_{cgt}/\Lambda < 13 \cdot 10^{-3} \text{TeV}^{-1}$  and on the branching fractions  $BR(t \rightarrow ug) < 4.0 \cdot 10^{-5}$  and  $BR(t \rightarrow cg) < 17 \cdot 10^{-5}$ .

100

## Measurement of the charge asymmetry in top quark pair production in 8 TeV pp collision data collected by the ATLAS experiment

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Daniel Marley<sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *University of Michigan (US)*

**Corresponding Author:** lroos@lpnhe.in2p3.fr

The charge asymmetry in top quark pair production,  $AC$ , measured using 20.3 fb<sup>-1</sup> of data recorded with the ATLAS detector at a center-of-mass energy  $\sqrt{s} = 8$  TeV is presented. Events where one top quark decays semi-leptonically ( $e$ +jets or  $\mu$ +jets) are considered. The  $t\bar{t}$  system is fully reconstructed using a likelihood fit. A Bayesian unfolding procedure is performed to determine  $AC$  at the parton level. An inclusive result is presented along with differential measurements with respect to the velocity of the  $t\bar{t}$  system,  $\beta_{z,t\bar{t}}$ , the transverse momentum of the  $t\bar{t}$  system,  $p_{T,t\bar{t}}$ , and the invariant mass of the  $t\bar{t}$  system,  $m_{t\bar{t}}$ . All measurements are consistent with Standard Model predictions.

101

## Measurement of colour flow with the jet pull angle in $t\bar{t}$ events using the ATLAS detector

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Tom Neep<sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *University of Manchester (GB)*

**Corresponding Author:** lroos@lfnhe.in2p3.fr

The distribution and orientation of energy inside jets is predicted to be an experimental handle on colour connections between the hard-scatter quarks and gluons initiating the jets. This poster presents a measurement of the distribution of one such variable, the jet pull angle. The pull angle is measured for jets produced in  $t\bar{t}$  events with one W boson decaying leptonically and the other decaying to jets using 20.3 inverse fb of data recorded with the ATLAS detector at a centre-of-mass energy of 8 TeV at the LHC. The jet pull angle distribution is corrected for detector resolution and acceptance effects and is compared to various models.

102

## Measurement of $t\bar{t}$ production with a veto on additional central jet activity in pp collisions at $\sqrt{s}=8$ TeV using the ATLAS detector

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Callie Bertsche<sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *University of Oklahoma (US)*

**Corresponding Author:** lroos@lfnhe.in2p3.fr

This poster presents a measurement of the amount of QCD radiation in top quark pair-production ( $t\bar{t}$ ) events, using dilepton  $t\bar{t}$  events that have an opposite-sign  $e\mu$  pair and two  $b$ -tagged jets in the final state. The measurement includes the complete 2012 ATLAS data sample of  $20.3 \text{ fb}^{-1}$  integrated luminosity of  $pp$  collision data at  $\sqrt{s} = 8 \text{ TeV}$ . The fraction of signal events that does not contain additional jet activity in a central rapidity region is measured as a function of (a) the minimum transverse momentum of any additional jet in the event, and (b) the minimum scalar transverse momentum sum of all additional jets in the event, and the results shown for four central rapidity regions and four regions of the invariant mass of the  $e\mu bb$  system. Compensation for detector effects is applied to the measurement and the results compared at the particle level to prediction from simulation by several next-to-leading order (NLO) and multi-leg leading order (LO) Monte Carlo generators.

103

## Measurement of $t\bar{t}$ differential cross-section for highly boosted top quarks produced at ATLAS in proton proton collisions at $\sqrt{s} = 8 \text{ TeV}$

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Federica Fabbri <sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *Universita e INFN, Bologna (IT)*

**Corresponding Author:** lroos@lpnhe.in2p3.fr

The measurement of the differential cross-section for top quark pair production as function of top transverse momentum will be presented.

The used dataset has been collected in 2012 in proton-proton collisions at a center of mass energy of 8 TeV.

The measurement is performed for  $t\bar{t}$  events in the semileptonic channel decay where the hadronically decaying top quark has a transverse momentum above 300 GeV.

The hadronic top quark decay is reconstructed as a single large radius jet and identified using the jet substructure properties.

This technique allows to increase the detection efficiency extending the cross-section measurement to high  $p_T$  region never reached before.

The main background sources are evaluated both with data driven methods (for  $W$ +jets and fake leptons contributions) and using Monte Carlo simulations ( $Z$ +jets, diboson,  $t\bar{t}$  dilepton channel, single top).

The observed yield, after the background subtraction, is corrected for efficiency and resolution effects to obtain the result both in a fiducial region close to the event selection and also extrapolated to the full partonic phase-space.

The final differential cross section has been compared with several theoretical predictions that generally slightly exceed the measured distribution.

104

## Top Tagging at ATLAS

**Author:** Julien Caudron<sup>1</sup>

<sup>1</sup> *Johannes-Gutenberg-Universitaet Mainz (DE)*

**Corresponding Author:** julien.caudron@cern.ch

Studies of the boosted sector in top-quark physics have known a fast-growing development with the arrival of high-energy data at LHC. Different techniques to identify high- $p_T$  top quarks based on substructure analyses of large radius jets have been developed for Run-1 and Run-2 data. New results are presented on the optimization and performance comparisons, for the different techniques (HEPTopTagger, Shower Deconstruction and substructure variables cut-based taggers), using pp collision data and MC simulations at  $\sqrt{s}=8$  TeV. In addition, prospects for Run-2 analyses, using MC simulations at  $\sqrt{s}=13$  TeV with 2015 data-taking conditions, are also presented.

105

## Studies of top quark pair modelling using ATLAS measurements

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** Maria Moreno Llacer <sup>2</sup>



<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *Georg-August-Universitaet Goettingen (DE)*

**Corresponding Author:** lroos@lpnhe.in2p3.fr

The data collected during the Run 1 at the LHC have given a large boost to measurements involving top quarks and triggered tremendous activities on the experimental and theoretical side.

The precision of most of these measurements are limited by the systematic uncertainties related to the modelling of the  $t\bar{t}$  signal process, to the event modelling and the description of the hard scattering environment.

Choices to be made in the signal simulation are the proton distribution functions, the Monte Carlo generator and the hadronisation model. On the event modelling side, important ingredients are related to the description of the underlying event, via Monte Carlo tunes, and the settings adopted for the modelling of colour reconnection, extra QCD radiation and the description of additional interactions accompanying the hard scatter (pile-up).

A huge experimental effort is underway with the aim of reducing the signal modelling uncertainties in close collaboration with the theory community. Recently new Monte Carlo generators based on NLO matrix elements for the hard scattering process matched with parton showers have become available. In addition, a new frontier has been reached with the advent of Monte Carlo generators based on merged NLO matrix elements with up to 4 partons hard scattering process and matched to parton showers.

This kind of studies not only for  $t\bar{t}$  process but also the production of top quark pairs in association with bosons, photons, Z and Higgs, which become now accessible at the LHC, are also of extremely interest. The predictions of various Monte Carlo generators are compared to data from Run 1 and presented in this work.

106

## First cross section measurements of $t\bar{t}$ pairs at $\sqrt{s} = 13$ TeV in the same—flavour dilepton events

**Author:** Lydia Roos<sup>1</sup>

**Co-author:** James William Howarth<sup>2</sup>

<sup>1</sup> *Centre National de la Recherche Scientifique (FR)*

<sup>2</sup> *Deutsches Elektronen-Synchrotron Hamburg and Zeuthen (DE)*

**Corresponding Author:** lroos@lpnhe.in2p3.fr

First results of the inclusive cross-section measurement of  $t\bar{t}$  events in the same-flavour dilepton decay mode are presented using data collected by the ATLAS experiment in Summer 2015 at a centre of mass energy of 13 TeV.

Session 8: Young Scientists Forum / 107

## Unravelling the non-standard top and Higgs couplings in associated top-Higgs production at the High-luminosity LHC

**Corresponding Author:** pankaj.sharma@adelaide.edu.au

108

## search for single top quark production in the $s$ -channel by the CMS at 8 TeV

**Author:** Ferdos Rezaei Hosseinabadi<sup>1</sup>

<sup>1</sup> *Institute for Research in Fundamental Sciences(IPM)*

Results of the search for single top-quark production in the  $s$ -channel in p-p collisions at a center-of-mass energy of 8TeV by the CMS experiment at the LHC will be presented. Leptonic decays mode of top quarks with an electron or muon in the final state are considered. In order to separate the expected signal from background processes, a multivariate discriminant is defined using Boosted Decision Trees. This analysis leads to an upper limit on the cross section of 11.5 Pb, corresponding to 2.1 times the standard model cross section, at 95% Confidence Level (TOP-13-009)

109

## First measurement of the differential cross section for $t\bar{t}$ production in the dilepton final state at a center-of-mass energy of 13 TeV

**Author:** Youn Jung Roh<sup>1</sup>

<sup>1</sup> *Korea University (KR)*

**Corresponding Author:** youn.jung.roh@cern.ch

Differential cross sections of top-quark pair production are measured in dilepton decay channel with proton-proton collisions at a center-of-mass energy of 13 TeV. The measurement is performed with a RunII data using CMS detector at the Large Hadron Collider. In this analysis, we measure the differential cross sections with respect to kinematic variables of leptons,  $b$ jets, and top-quarks.

110

## Constraining QCD multijet background in single top $t$ -channel production at 13 TeV

**Author:** Georgios Krintiras<sup>1</sup>

<sup>1</sup> *Universite Catholique de Louvain (UCL) (BE)*

Precision measurement of the cross section for single top production in  $t$  channel is an important test of the Standard Model (SM). The purity of the collected data in single top events is limited by the understanding of the shape and yield of background contributions. Besides electroweak and  $t\bar{t}$  processes, QCD multijet events are a non-negligible background for this measurement, with large intrinsic modeling uncertainties. We report on how we control this contamination in the context of the measurement of  $t$ -channel single top cross section at 13 TeV.