

# Dynamical Relaxation of Axion Misalignment in the Early Universe

Eric Gonzalez

University of Michigan

Phenomenology Symposium 2019

# Background

# Why QCD Axion Dark Matter?

- Three questions for any dark matter (DM) candidate:
  - **Does it solve other problems in physics?**
  - **Can it be produced in the early universe?**
  - **Is it detectable?**

# Why QCD Axion Dark Matter?

- Three questions for any dark matter (DM) candidate:
  - **Does it solve other problems in physics?**
  - **QCD axion solves Strong CP problem**
  - **Can it be produced in the early universe?**
  
- **Is it detectable?**


# Why QCD Axion Dark Matter?

- Three questions for any dark matter (DM) candidate:
  - **Does it solve other problems in physics?**
  - QCD axion solves Strong CP problem
  - **Can it be produced in the early universe?**
  - QCD Axion DM can be produced non-thermally
    - Misalignment, Topological Defects
  - **Is it detectable?**

# Why QCD Axion Dark Matter?

- Three questions for any dark matter (DM) candidate:
  - **Does it solve other problems in physics?**
  - QCD axion solves Strong CP problem
  - **Can it be produced in the early universe?**
  - QCD Axion DM can be produced non-thermally
    - Misalignment, Topological Defects
  - **Is it detectable?**
  - Many experiments, current and planned.


# QCD Axion Experiments



Arvanitaki et. Al  
1411.2263

Raffelt  
0611350

# Misalignment Mechanism (Pre-Inflationary PQ Breaking)




Arvanitaki et. Al  
1411.2263

Raffelt  
0611350

# Standard Misalignment Mechanism

$$\ddot{\theta}_a + 3H \dot{\theta} = -m_a^2 \sin\theta_a$$


$H > m_a$  axion is frozen



# Standard Misalignment Mechanism

$$\ddot{\theta}_a + 3H \dot{\theta} = -m_a^2 \sin\theta_a$$

$H < m_a$  axion oscillates




$$\Omega_a h^2 = 0.12 \theta_{mis}^2 \mathcal{F}(\theta_{mis}) \left( \frac{f_a}{5 \times 10^{11} \text{GeV}} \right)^{\frac{7}{6}}$$

# Standard Misalignment Mechanism

$$\ddot{\theta}_a + 3H \dot{\theta} = -m_a^2 \sin\theta_a$$

$H < m_a$  axion oscillates



$$\Omega_a h^2 = 0.12 \theta_{mis}^2 \mathcal{F}(\theta_{mis}) \left( \frac{f_a}{5 \times 10^{11} \text{GeV}} \right)^{\frac{7}{6}}$$

# Case 1


## Large Decay Constant

# Standard Misalignment Mechanism

— Early Universe  $V_a$   
— Low Temp.  $V_a$

Assumed same potential  
in early universe:

No Early Universe  
Dynamics  
in Standard picture



$H > m_a$  axion is frozen

# New Early Universe Axion Dynamics


— Early Universe  $V_a$   
— Low Temp.  $V_a$

Lift Assumption:

Axion can relax in Early Universe

Lifted assumption provides

scenario where  $\theta_{mis} \ll 1$



$H > m_a$  axion is frozen


$H < m_a$  axion oscillates

# Axion Mass and $\Lambda_{\text{QCD}}$

$$m_a \propto \frac{\Lambda_{\text{QCD}}^2}{f_a}$$

$$\mu_r \frac{d}{d\mu_r} \frac{8\pi^2}{g^2} = 3N - F$$


$$\mathcal{L} \supset \lambda_u u^c Q \phi_u + \lambda_d d^c Q \phi_d$$



# $\Lambda_{\text{QCD}}$ and the Higgs VEV

$$m_a \propto \frac{\Lambda_{\text{QCD}}^2}{f_a}$$

$$\Lambda_{\text{QCD}}^3 \propto \langle \phi \rangle^2$$




Dvali 9505253  
Banks and Dine 9608197

# Hubble Induced Mass

$$\Delta K \sim \frac{|X|^2}{M^2} |\phi|^2 - \frac{|X|^2}{M^4} |\phi|^4$$

$$V_\phi \sim (m_\phi^2 - cH_I^2) |\phi|^2 + \frac{cH_I^2}{M^2} |\phi|^4 + \dots$$




# Phase of the Axion Potential

— Early Universe  $V_a$   
— Low Temp.  $V_a$

$$\bar{\theta} = \theta_{\text{QCD}} + \arg(\det[m_u m_d]) + \dots$$

$$\bar{\theta}_{\text{MSSM}} = \theta_{\text{QCD}} + \arg(\det[\lambda_u \lambda_d]) + 3\arg(B\mu) + \dots$$



# Dynamical Axion Misalignment Production


Two criteria for DAMP:

1) Axion must relax in the early universe

2) CP structure relates Early/Late axion potential minima

— Early Universe  $V_a$

— Low Temp.  $V_a$




# Case 2

## Small Decay Constant

# Phase of the Axion Potential

$$\bar{\theta}_{\text{MSSM}} = \theta_{\text{QCD}} + \arg(\det[\lambda_u \lambda_d]) + 3\arg(B\mu)$$

— Early Universe  $V_a$   
— Low Temp.  $V_a$




# Mass of the Higgs Phase

$$\bar{\theta}_{\text{MSSM}} = \theta_{\text{QCD}} + \arg(\det[\lambda_u \lambda_d]) + 3\arg(B\mu)$$

$$V_\phi \sim (B\mu - cH_I^2)\phi_u\phi_d$$

— Early Universe  $V_a$   
— Low Temp.  $V_a$




# Shifting the Potential


$$\bar{\theta}_{\text{MSSM}} = \theta_{\text{QCD}} + \arg(\det[\lambda_u \lambda_d]) + 3\arg(B\mu)$$

$$V_\phi \sim (B\mu - cH_I^2)\phi_u\phi_d$$

$\pi$  shift if  $B\mu < cH_I^2$




# Anharmonic Effect



CPV from Yukawa Couplings

Electron EDM in TeV scale SUSY


# New Motivated Regions




Arvanitaki et. Al  
1411.2263

Raffelt  
0611350

# Summary Table





DAMP <sub>0</sub>	DAMP <sub>π</sub>
$\theta_i \rightarrow 0$	$\theta_i \rightarrow \pi$
Decay constant predicted by CP structure	
Isocurvature perturbations are suppressed.	
Jeong and Takahashi 1308.8131	
$f_a \gtrsim 10^{16} \text{ GeV}$ typical string axion scale. <small>Svrcek and Witten 0605206</small>	Anharmonic effect could enhance structure formation. <small>Greene, Kofman, and Starobinsky 9808477</small>



Thank you!

# Cosmology Bounds



# Early Universe Axion Dynamics

— Early Universe  $V_a$   
— Low Temp.  $V_a$

$$\bar{\theta} = \theta_{\text{QCD}} + \arg(\det[m_u m_d]) + \dots$$

$$\mathcal{L} \subset \lambda_u u^c Q \phi_u + \lambda_d d^c Q \phi_d$$

$$\bar{\theta}_{\text{MSSM}} = \theta_{\text{QCD}} + \arg(\det[\lambda_u \lambda_d]) + 3\arg(B\mu)$$

