

Double Field Theory at $SL(2)$ angles


Adolfo Guarino

Université Libre de Bruxelles

The String Theory Universe 2017
February 21st, Milano

Based on [arXiv:1604.08602](https://arxiv.org/abs/1604.08602) & [arXiv:1612.05230](https://arxiv.org/abs/1612.05230)

w/ F. Ciceri & G. Inverso (and G. Dibitetto & J. Melgarejo)



Duality covariant approaches to strings

- Different strings related by dualities: IIA/IIB T-duality, IIB S-duality, ...

- *String dualities* realised as global symmetries in SUGRA

lower-dimensional phenomenon

embedding tensor, non-geometry...

vs

higher-dimensional phenomenon

β -supergravity, γ -supergravity...

Today's talk : **Extended Field Theories** [extend internal coords to transform under duality]

- Double Field Theory (DFT) \Rightarrow Orthogonal groups $O(d,d)$ [half-max SUGRA (T-duality)]
- Exceptional Field Theory (EFT) \Rightarrow Exceptional groups $E_{d+1(d+1)}$ [max SUGRA (U-duality)]

[Siegel '93] [Hull & Zwiebach (Hohm) '09 '10] [Hohm & Samtleben '13]

Dualities in SUGRA and Extended Field Theory


D	Maximal sugra / EFT	Half-maximal sugra	DFT
9	$\mathbb{R}^+ \times \text{SL}(2)$	$\mathbb{R}^+ \times \text{O}(1, 1 + n)$	$\mathbb{R}^+ \times \text{O}(1, 1 + n)$
8	$\text{SL}(2) \times \text{SL}(3)$	$\mathbb{R}^+ \times \text{O}(2, 2 + n)$	$\mathbb{R}^+ \times \text{O}(2, 2 + n)$
7	$\text{SL}(5)$	$\mathbb{R}^+ \times \text{O}(3, 3 + n)$	$\mathbb{R}^+ \times \text{O}(3, 3 + n)$
6	$\text{SO}(5, 5)$	$\mathbb{R}^+ \times \text{O}(4, 4 + n)$	$\mathbb{R}^+ \times \text{O}(4, 4 + n)$
5	$\text{E}_{6(6)}$	$\mathbb{R}^+ \times \text{O}(5, 5 + n)$	$\mathbb{R}^+ \times \text{O}(5, 5 + n)$
4	$\text{E}_{7(7)}$	$\text{SL}(2) \times \text{O}(6, 6 + n)$	$\mathbb{R}^+ \times \text{O}(6, 6 + n)$
3	$\text{E}_{8(8)}$	$\text{O}(8, 8 + n)$	$\mathbb{R}^+ \times \text{O}(7, 7 + n)$

Duality groups of half-maximal SUGRA and DFT differ for $\mathbf{D} \leq 4$

... in this talk we will look at $D=4$:

EFT with $E_{7(7)}$ duality group

[Hohm & Samtleben '13]



SL(2)-DFT with $SL(2) \times O(6,6+n)$ duality group

[arXiv:1612.05230]



DFT with $R^+ \times O(6,6+n)$ duality group

[Siegel '93]

[Hull & Zwiebach '09]

[Hohm, Hull & Zwiebach '10]

[Hohm & Kwak '11]

$E_{7(7)}$ -EFT

[momentum, winding, ...]

- Space-time : external ($D=4$) + **generalised internal** ($y^{\mathcal{M}}$ coordinates in **56** of $E_{7(7)}$)

Generalised diffs = ordinary internal diffs + internal gauge transfos

- Generalised Lie derivative built from an $E_{7(7)}$ -invariant **structure Y-tensor**

$$\mathbb{L}_\Lambda U^{\mathcal{M}} = \Lambda^{\mathcal{N}} \partial_{\mathcal{N}} U^{\mathcal{M}} - U^{\mathcal{N}} \partial_{\mathcal{N}} \Lambda^{\mathcal{M}} + Y^{\mathcal{M}\mathcal{N}}{}_{\mathcal{P}\mathcal{Q}} \partial_{\mathcal{N}} \Lambda^{\mathcal{P}} U^{\mathcal{Q}}$$

Closure requires a **section constraint** : $Y^{\mathcal{P}\mathcal{Q}}{}_{\mathcal{M}\mathcal{N}} \partial_{\mathcal{P}} \otimes \partial_{\mathcal{Q}} = 0$

Two maximal solutions : M-theory (**7** dimensional) & Type IIB (**6** dimensional)

[massless theories]

Massive IIA arises as a *deformation of EFT*

[Ciceri, A.G. & Inverso '16]

[Hohm & Kwak '11 (sec const violated)]

[Romans '86]

[Cassani, de Felice, Petrini, Strickland-Constable & Waldram '16 (generalised geometry)]

E₇₍₇₎-EFT

- E₇₍₇₎-EFT action [$\mathcal{D}_\mu = \partial_\mu - \mathbb{L}_{A_\mu}$]

$$S_{\text{EFT}} = \int d^4x d^{56}y e \left[\hat{R} + \frac{1}{48} g^{\mu\nu} \mathcal{D}_\mu \mathcal{M}^{\mathcal{MN}} \mathcal{D}_\nu \mathcal{M}_{\mathcal{MN}} - \frac{1}{8} \mathcal{M}_{\mathcal{MN}} \mathcal{F}^{\mu\nu\mathcal{M}} \mathcal{F}_{\mu\nu}{}^{\mathcal{N}} + e^{-1} \mathcal{L}_{\text{top}} - V_{\text{EFT}}(\mathcal{M}, g) \right]$$

with *field strengths* & *potential term* given by

$$\mathcal{F}_{\mu\nu}{}^{\mathcal{M}} = 2 \partial_{[\mu} A_{\nu]}{}^{\mathcal{M}} - [A_\mu, A_\nu]_{\text{E}}{}^{\mathcal{M}} + \text{two-form terms} \quad (\text{tensor hierarchy})$$

$$V_{\text{EFT}}(\mathcal{M}, g) = -\frac{1}{48} \mathcal{M}^{\mathcal{MN}} \partial_{\mathcal{M}} \mathcal{M}^{\mathcal{KL}} \partial_{\mathcal{N}} \mathcal{M}_{\mathcal{KL}} + \frac{1}{2} \mathcal{M}^{\mathcal{MN}} \partial_{\mathcal{M}} \mathcal{M}^{\mathcal{KL}} \partial_{\mathcal{L}} \mathcal{M}_{\mathcal{NK}} - \frac{1}{2} g^{-1} \partial_{\mathcal{M}} g \partial_{\mathcal{N}} \mathcal{M}^{\mathcal{MN}} - \frac{1}{4} \mathcal{M}^{\mathcal{MN}} g^{-1} \partial_{\mathcal{M}} g g^{-1} \partial_{\mathcal{N}} g - \frac{1}{4} \mathcal{M}^{\mathcal{MN}} \partial_{\mathcal{M}} g^{\mu\nu} \partial_{\mathcal{N}} g_{\mu\nu}$$

- **Two-derivative** potential : **ungauged** N=8 D=4 SUGRA when $\Phi(x, y) = \Phi(x)$

From $E_{7(7)}$ -EFT to $SL(2)$ -DFT

- Halving EFT with $E_{7(7)}$ symmetry to obtain $SL(2)$ -DFT with $SL(2) \times O(6,6)$ symmetry

$E_{7(7)}$	\rightarrow	$SL(2) \times SO(6,6)$	$\alpha = (+, -)$ vector index of $SL(2)$
56	\rightarrow	(2, 12) + (1, 32)	M vector index of $SO(6,6)$
$y^{\mathcal{M}}$	\rightarrow	$y^{\alpha M} + \cancel{y^A}$	A M-W spinor index of $SO(6,6)$
<hr style="width: 100%; border: 0.5px solid blue;"/>		<hr style="width: 100%; border: 0.5px solid red;"/>	
EFT		SL(2)-DFT	[see Dibitetto, A.G. & Roest '11 for SUGRA]

via a **Z₂ truncation** (*vector* = +1 , *spinor* = -1) on coordinates, fields, etc.

- $SL(2)$ -DFT generalised Lie derivative [DFT corresponds to an $\alpha = +$ orientation]

$$\mathbb{L}_\Lambda U^{\alpha M} = \Lambda^{\beta N} \partial_{\beta N} U^{\alpha M} - U^{\beta N} \partial_{\beta N} \Lambda^{\alpha M} + \eta^{MN} \eta_{PQ} \partial_{\beta N} \Lambda^{\beta P} U^{\alpha Q} + 2 \epsilon^{\alpha\beta} \epsilon_{\gamma\delta} \partial_{\beta N} \Lambda^{\gamma[M} U^{\delta|N]}$$

- $SL(2)$ -DFT section constraints :

$$\eta^{MN} \partial_{\alpha M} \otimes \partial_{\beta N} = 0 \quad , \quad \epsilon^{\alpha\beta} \partial_{\alpha[M} \otimes \partial_{\beta|N]} = 0$$

N=1 SUGRA in D=10 as in DFT

SL(2)-DFT with SL(2) x O(6,6) symmetry

- SL(2)-DFT action [$\mathcal{D}_\mu = \partial_\mu - \mathbb{L}_{A_\mu}$]

$$S_{\text{SL}(2)\text{-DFT}} = \int d^4x d^{24}y e \left[\hat{R} + \frac{1}{4} g^{\mu\nu} \mathcal{D}_\mu M^{\alpha\beta} \mathcal{D}_\nu M_{\alpha\beta} + \frac{1}{8} g^{\mu\nu} \mathcal{D}_\mu M^{MN} \mathcal{D}_\nu M_{MN} \right. \\ \left. - \frac{1}{8} M_{\alpha\beta} M_{MN} \mathcal{F}^{\mu\nu\alpha M} \mathcal{F}_{\mu\nu}{}^{\beta N} + e^{-1} \mathcal{L}_{\text{top}} - V_{\text{SL}(2)\text{-DFT}}(M, g) \right]$$

with field strengths & potential term given by

$$\mathcal{F}_{\mu\nu}{}^{\alpha M} = 2 \partial_{[\mu} A_{\nu]}{}^{\alpha M} - [A_\mu, A_\nu]_S{}^{\alpha M} + \text{two-form terms} \quad (\text{tensor hierarchy})$$

$$V_{\text{SL}(2)\text{-DFT}}(M, g) = M^{\alpha\beta} M^{MN} \left[-\frac{1}{4} (\partial_{\alpha M} M^{\gamma\delta})(\partial_{\beta N} M_{\gamma\delta}) - \frac{1}{8} (\partial_{\alpha M} M^{PQ})(\partial_{\beta N} M_{PQ}) \right. \\ \left. + \frac{1}{2} (\partial_{\alpha M} M^{\gamma\delta})(\partial_{\delta N} M_{\beta\gamma}) + \frac{1}{2} (\partial_{\alpha M} M^{PQ})(\partial_{\beta Q} M_{NP}) \right] \\ + \frac{1}{2} M^{MN} M^{PQ} (\partial_{\alpha M} M^{\alpha\delta})(\partial_{\delta Q} M_{NP}) + \frac{1}{2} M^{\alpha\beta} M^{\gamma\delta} (\partial_{\alpha M} M^{MQ})(\partial_{\delta Q} M_{\beta\gamma}) \\ - \frac{1}{4} M^{\alpha\beta} M^{MN} \left[g^{-1}(\partial_{\alpha M} g) g^{-1}(\partial_{\beta N} g) + (\partial_{\alpha M} g^{\mu\nu})(\partial_{\beta N} g_{\mu\nu}) \right] \\ - \frac{1}{2} g^{-1}(\partial_{\alpha M} g) \partial_{\beta N}(M^{\alpha\beta} M^{MN})$$

- Two-derivative potential : **ungauged** N=4 D=4 SUGRA when $\Phi(x, y) = \Phi(x)$

SL(2) angles & moduli stabilisation


- Scherk-Schwarz (SS) reductions with $SL(2) \times O(6,6)$ twist matrices $U_{\alpha M}{}^{\beta N} = e^\lambda e_\alpha{}^\beta U_M{}^N$ yield **N=4** , **D=4** gaugings [Schön & Weidner '06]

$$f_{\alpha MNP} = -3 e^{-\lambda} e_\alpha{}^\beta \eta_{Q[M} U_N{}^R U_P]{}^S \partial_{\beta R} U_S{}^Q$$

$$\xi_{\alpha M} = 2 U_M{}^N \partial_{\beta N} (e^{-\lambda} e_\alpha{}^\beta)$$

[de Roo & Wagemans '85]

- Moduli stabilisation **requires** gaugings $G = G_1 \times G_2$ at **relative** SL(2) angles



(sec. constraint **violated**)

$$\epsilon^{\alpha\beta} \partial_{\alpha[M} \otimes \partial_{\beta|N]} \neq 0$$

- Dependence on **both** type + & type - coordinates [**not possible** in DFT]

Example : SO(4) x SO(4) gaugings and non-geometry

- SS with $U(y^{\alpha M}) \in O(6,6)$: Half of the coords of **type +** & half of **type -**
- SL(2)-superposition of *two chains* of **non-geometric** fluxes $(H, \omega, Q, R)_{\pm}$

$$f_+ \quad f_{+mnp} = H^+_{mnp} \quad , \quad f_{+mn\bar{p}} = \omega^+_{mn}{}^p \quad , \quad f_{+\bar{m}\bar{n}p} = Q^{+mn}{}_p \quad , \quad f_{+\bar{m}\bar{n}\bar{p}} = R^{+mnp}$$

$$f_- \quad f_{-mnp} = H^-_{mnp} \quad , \quad f_{-mn\bar{p}} = \omega^-_{mn}{}^p \quad , \quad f_{-\bar{m}\bar{n}p} = Q^{-mn}{}_p \quad , \quad f_{-\bar{m}\bar{n}\bar{p}} = R^{-mnp}$$

Most general family (8 params) of SO(4) x SO(4) gaugings of N=4 SUGRA

- SO(4) x SO(4) SUGRA : **AdS₄** & **dS₄** vacua (sphere/hyperboloid reductions)

[de Roo, Westra, Panda & Trigiante '03] [Dibitetto, A.G. & Roest '12]

- "Hybrid \pm " sources to cancel flux-induced tadpoles : SL(2)-dual NS-NS branes

Summary & Future directions


- **SL(2)-DFT** captures the duality group of N=4 SUGRA in D=4
 - **SL(2)-DFT** sec. constraints : N=1 SUGRA in D=10 & N=(2,0) SUGRA in D=6
 - **SL(2)-DFT** action extendable to $SL(2) \times SO(6,6+n)$ and *deformable as EFT*
[Ciceri, A.G. & Inverso '16]
 - Non-geometric gaugings at non-trivial SL(2) angles : *full moduli AdS₄ stabilisation*
[**not** possible in DFT]
-
- **Flux formulation** of SL(2)-DFT : sec. cons violating terms & SL(2)-NS-NS branes
[Aldazabal, Graña, Marqués & Rosabal '13]
 - **Cosmological applications** of SL(2)-DFT (stable dS₄ , inflation, ...)
[Hassler, Lüst & Massai '14]

Grazie mille!!

Thanks a lot!!

Extra material

A family of exotic branes



[Sakatani '15]

Dualities in SUGRA and Extended Field Theory

D	Maximal sugra / EFT	Half-maximal sugra	DFT
9	$\mathbb{R}^+ \times \text{SL}(2)$	$\mathbb{R}^+ \times \text{O}(1, 1 + n)$	$\mathbb{R}^+ \times \text{O}(1, 1 + n)$
8	$\text{SL}(2) \times \text{SL}(3)$	$\mathbb{R}^+ \times \text{O}(2, 2 + n)$	$\mathbb{R}^+ \times \text{O}(2, 2 + n)$
7	$\text{SL}(5)$	$\mathbb{R}^+ \times \text{O}(3, 3 + n)$	$\mathbb{R}^+ \times \text{O}(3, 3 + n)$
6	$\text{SO}(5, 5)$	$\mathbb{R}^+ \times \text{O}(4, 4 + n)^*$	$\mathbb{R}^+ \times \text{O}(4, 4 + n)$
5	$\text{E}_{6(6)}$	$\mathbb{R}^+ \times \text{O}(5, 5 + n)$	$\mathbb{R}^+ \times \text{O}(5, 5 + n)$
4	$\text{E}_{7(7)}$	$\text{SL}(2) \times \text{O}(6, 6 + n)$	$\mathbb{R}^+ \times \text{O}(6, 6 + n)$
3	$\text{E}_{8(8)}$	$\text{O}(8, 8 + n)$	$\mathbb{R}^+ \times \text{O}(7, 7 + n)$

Duality groups of half-maximal SUGRA and DFT differ for $\mathbf{D} \leq 4$

* There is also the chiral N=(2,0) SUGRA in D=6 with $\mathbb{R}^+ \times \text{O}(5, n)$ duality group

SO(4) x SO(4) twist matrices

- O(6,6) twist :
$$U_M^N(y^{\alpha M}) = \begin{pmatrix} \mathbb{I}_6 & 0_6 \\ \beta & \mathbb{I}_6 \end{pmatrix} \begin{pmatrix} \mathbb{I}_6 & b \\ 0_6 & \mathbb{I}_6 \end{pmatrix} \begin{pmatrix} u & 0_6 \\ 0_6 & u^{-t} \end{pmatrix} = \begin{pmatrix} u_m^{\underline{n}} & b_{mp} (u^{-t})^p_{\underline{n}} \\ \beta^{mp} u_p^{\underline{n}} & (u^{-t})^m_{\underline{n}} + \beta^{mp} b_{pq} (u^{-t})^q_{\underline{n}} \end{pmatrix}$$

where
$$\beta^{mn} = \begin{pmatrix} (\beta_{(1)})^{ab} & 0_3 \\ 0_3 & (\beta_{(2)})^{ij} \end{pmatrix}, \quad b_{mn} = \begin{pmatrix} (b_{(1)})_{ab} & 0_3 \\ 0_3 & (b_{(2)})_{ij} \end{pmatrix}, \quad u_m^{\underline{n}} = \begin{pmatrix} (u_{(1)})_a^b & 0_3 \\ 0_3 & (u_{(2)})_i^j \end{pmatrix}$$

$$u_{(1),(2)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} (\cos Y_{(1),(2)} + \cos \tilde{Y}_{(1),(2)}) & -\frac{1}{2} (\sin Y_{(1),(2)} + \sin \tilde{Y}_{(1),(2)}) \\ 0 & \frac{1}{2} (\sin Y_{(1),(2)} + \sin \tilde{Y}_{(1),(2)}) & \frac{1}{2} (\cos Y_{(1),(2)} + \cos \tilde{Y}_{(1),(2)}) \end{pmatrix},$$

$$b_{(1),(2)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} \sin(Y_{(1),(2)} - \tilde{Y}_{(1),(2)}) \\ 0 & -\frac{1}{2} \sin(Y_{(1),(2)} - \tilde{Y}_{(1),(2)}) & 0 \end{pmatrix},$$

$$\beta_{(1),(2)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \tan\left(\frac{1}{2}(Y_{(1),(2)} - \tilde{Y}_{(1),(2)})\right) \\ 0 & -\tan\left(\frac{1}{2}(Y_{(1),(2)} - \tilde{Y}_{(1),(2)})\right) & 0 \end{pmatrix},$$

$$Y_{(1)} = (\tilde{c}'_1 - a'_0) (y^{+1} - y^{+\bar{1}}) + (\tilde{d}'_1 - b'_0) (y^{-1} - y^{-\bar{1}})$$

$$\tilde{Y}_{(1)} = (\tilde{c}'_1 + a'_0) (y^{+1} + y^{+\bar{1}}) + (\tilde{d}'_1 + b'_0) (y^{-1} + y^{-\bar{1}})$$

$$Y_{(2)} = (\tilde{c}'_2 - a'_3) (y^{+4} - y^{+\bar{4}}) + (\tilde{d}'_2 - b'_3) (y^{-4} - y^{-\bar{4}})$$

$$\tilde{Y}_{(2)} = (\tilde{c}'_2 + a'_3) (y^{+4} + y^{+\bar{4}}) + (\tilde{d}'_2 + b'_3) (y^{-4} + y^{-\bar{4}})$$

Deformed EFT (XFT)

- Generalised Lie derivative

[no density term]

$$\mathbb{L}_\Lambda U^\mathcal{M} = \Lambda^\mathcal{N} \partial_\mathcal{N} U^\mathcal{M} - U^\mathcal{N} \partial_\mathcal{N} \Lambda^\mathcal{M} + Y^{\mathcal{MN}}_{\mathcal{PQ}} \partial_\mathcal{N} \Lambda^\mathcal{P} U^\mathcal{Q}$$

in terms of an $E_{n(n)}$ -invariant **structure Y-tensor**. Closure requires **sec. constraint**

- **Deformed** generalised Lie derivative

$$\tilde{\mathbb{L}}_\Lambda U^\mathcal{M} = \Lambda^\mathcal{N} \partial_\mathcal{N} U^\mathcal{M} - U^\mathcal{N} \partial_\mathcal{N} \Lambda^\mathcal{M} + Y^{\mathcal{MN}}_{\mathcal{PQ}} \partial_\mathcal{N} \Lambda^\mathcal{P} U^\mathcal{Q} - \underbrace{X_{\mathcal{NP}}{}^\mathcal{M}}_{\text{non-derivative}} \Lambda^\mathcal{N} U^\mathcal{P}$$

in terms of an **X deformation** which is $E_{n(n)}$ -algebra valued

non-derivative

- Closure & triviality of the Jacobiator require (together with **sec. constraint**)

$$X_{\mathcal{MN}}{}^\mathcal{P} \partial_\mathcal{P} = 0$$

X constraint

$$X_{\mathcal{MP}}{}^\mathcal{Q} X_{\mathcal{NQ}}{}^\mathcal{R} - X_{\mathcal{NP}}{}^\mathcal{Q} X_{\mathcal{MQ}}{}^\mathcal{R} + X_{\mathcal{MN}}{}^\mathcal{Q} X_{\mathcal{QP}}{}^\mathcal{R} = 0$$

Quadratic constraint (gauged max. supergravity)

X deformation : background fluxes & Romans mass


$$Y^{\mathcal{P}\mathcal{Q}}{}_{MN} \partial_{\mathcal{P}} \otimes \partial_{\mathcal{Q}} = 0$$

section constraint

$$X_{MN}{}^{\mathcal{P}} \partial_{\mathcal{P}} = 0$$

X constraint

[algebraic system]



M-theory (n coords)

- SL(n) orbit
- Freund-Rubin param. (n = 4 and n = 7)
- *massless* IIA (subcase)

Type IIB (n-1 coords)

- SL(n-1) orbit
- p-form fluxes compatible with SL(n-1)
- **SL(2)-triplet of 1-form flux** (includes compact SO(2))

+

New massive Type IIA (n-1 coords)

- SL(n-1) orbit
- p-form fluxes compatible with SL(n-1)
- dilaton flux
- **Romans mass parameter** (kills the M-theory coord)

Massive Type IIA described in a purely geometric manner !!

[QC = flux-induced tadpoles]

E₇₍₇₎-XFT action

- E₇₍₇₎-XFT action [$\mathcal{D}_\mu = \partial_\mu - \tilde{\mathbb{L}}_{A_\mu}$] [$y^{\mathcal{M}}$ coords in the **56** of E₇₍₇₎]

$$S_{\text{XFT}} = \int d^4x d^{56}y e \left[\hat{R} + \frac{1}{48} g^{\mu\nu} \mathcal{D}_\mu \mathcal{M}^{\mathcal{MN}} \mathcal{D}_\nu \mathcal{M}_{\mathcal{MN}} - \frac{1}{8} \mathcal{M}_{\mathcal{MN}} \mathcal{F}^{\mu\nu\mathcal{M}} \mathcal{F}_{\mu\nu}{}^{\mathcal{N}} + e^{-1} \mathcal{L}_{\text{top}} - V_{\text{XFT}}(\mathcal{M}, g) \right]$$

with field strengths & potential given by

(deformed tensor hierarchy)

$$\mathcal{F}_{\mu\nu}{}^{\mathcal{M}} = 2 \partial_{[\mu} A_{\nu]}{}^{\mathcal{M}} + X_{[\mathcal{P}\mathcal{Q}]}{}^{\mathcal{M}} A_\mu{}^{\mathcal{P}} A_\nu{}^{\mathcal{Q}} - [A_\mu, A_\nu]_{\mathbb{E}}{}^{\mathcal{M}} + \text{two-form terms}$$

$$V_{\text{XFT}}(\mathcal{M}, g, X) = V_{\text{EFT}}(\mathcal{M}, g) + \underbrace{\frac{1}{12} \mathcal{M}^{\mathcal{MN}} \mathcal{M}^{\mathcal{KL}} X_{\mathcal{MK}}{}^{\mathcal{P}} \partial_{\mathcal{N}} \mathcal{M}_{\mathcal{PL}}}_{\text{cross term}} + \underbrace{V_{\text{SUGRA}}(\mathcal{M}, X)}_{\text{gauged max. sugra}}$$

- Two-One-**Zero**-derivative potential : **gauged** 4D max. sugra when $\Phi(x, y) = \Phi(x)$

Extended (super) Poincaré superalgebra

- Central charges (internal symmetries) $Z_{IJ} = (a_{IJ}^a) T^a$

- The algebra :

$$[P_\mu, P_\nu] = 0 \quad [M_{\mu\nu}, M_{\rho\sigma}] = i(\eta_{\nu\rho} M_{\mu\sigma} - \eta_{\nu\sigma} M_{\mu\rho} - \eta_{\mu\rho} M_{\nu\sigma} + \eta_{\mu\sigma} M_{\nu\rho})$$

$$[P_\mu, M_{\rho\sigma}] = i(\eta_{\mu\rho} P_\sigma - \eta_{\mu\sigma} P_\rho)$$

$$[T^a, T^b] = if_{ab}^c T^c \quad [T^a, P_\mu] = [T^a, M_{\mu\nu}] = 0$$

$$[Q'_\alpha, P_\mu] = [\bar{Q}'_{\dot{\alpha}}, P_\mu] = 0 \quad [Q'_\alpha, T^a] = (b_a)'_J Q'^J_\alpha \quad [\bar{Q}'_{\dot{\alpha}}, T^a] = -\bar{Q}'_{\dot{\alpha}} (b_a)_{J'}^J$$

$$[Q'_\alpha, M_{\mu\nu}] = \frac{1}{2}(\sigma_{\mu\nu})_\alpha^\beta Q'_\beta \quad [\bar{Q}'_{\dot{\alpha}}, M_{\mu\nu}] = -\frac{1}{2}\bar{Q}'_{\dot{\beta}} (\bar{\sigma}_{\mu\nu})^{\dot{\beta}}_{\dot{\alpha}}$$

$$\{\bar{Q}'_{\dot{\alpha}}, \bar{Q}'_{\dot{\beta}}\} = -2\epsilon_{\dot{\alpha}\dot{\beta}} Z^{IJ\dagger} \quad \{Q'_\alpha, Q'_\beta\} = 2\epsilon_{\alpha\beta} Z^{IJ} \quad \{Q'_\alpha, \bar{Q}'_{\dot{\beta}}\} = 2\delta^{IJ} (\sigma^\mu)_{\alpha\dot{\beta}} P_\mu$$