

Phenomenology 2021 Symposium



Contribution ID: 1289

Type: **Higgs**

A suppressed Higgs coupling in a classically conformal extension of the standard model

Wednesday 26 May 2021 16:30 (15 minutes)

We consider a classically conformal $U(1)$ extension of the Standard Model (SM). The $U(1)$ symmetry is radiatively broken by the Coleman-Weinberg mechanism, after which the $U(1)$ Higgs field ϕ drives electroweak symmetry breaking through a mixed quartic coupling with the SM Higgs doublet with coupling constant λ_{mix} . We calculate the Higgs triple couplings in this system and find a suppression of the coupling $g_{h\phi\phi}$ when compared to the naively expected value $g_{h\phi\phi} \sim \lambda_{mix} v_h$ ($v_h = 246$ GeV), likely due to the unique nature of the classically conformal potential. The suppression opens up parameter space for the mixing angle θ between SM Higgs and $U(1)$ Higgs eigenstates. We consider experimental signals for such conformal structure via the anomalous Higgs decay $h \rightarrow \phi\phi$ and anomalous SM Higgs couplings. The conformal structure would allow for a sizeable anomalous SM Higgs coupling alongside a heavily suppressed $h \rightarrow \phi\phi$ decay mode.

Summary

Authors: BAULES, Victor (University of Alabama, The); OKADA, Nobuchika (University of Alabama)

Presenter: BAULES, Victor (University of Alabama, The)

Session Classification: Higgs IV