


Tuning Pythia for Forward Physics




Felix Kling -
SLAC




Holger Schulz -
Durham University




Max Fieg -
UC Irvine



[mfieg at uci dot edu](mailto:mfieg@uci.edu)







LS2 Report: FASER is born

FASER, the Forward Search Experiment, has been installed in the LHC tunnel during Long Shutdown 2. It is currently being tested and will start taking data next year

24 MARCH, 2021 | By Anaïs Schaeffer




The final elements of FASER were put into place this month. (Image: CERN)



LS2 Report: FASER is born

FASER, the Forward Search Experiment, has been installed in the LHC tunnel during Long Shutdown 2. It is currently being tested and will start taking data next year

24 MARCH, 2021 | By Anaïs Schaefer




The final elements of FASER were put into place this month. (Image: CERN)

“If you are the 100th person to look under a rock, you are unlikely to find anything...

...but if you are using a new tool, or find a rock that's been left unturned, you don't have to be smart to find something”




Dr. Steven
Chu




LS2 Report: FASER is born

FASER, the Forward Search Experiment, has been installed in the LHC tunnel during Long Shutdown 2. It is currently being tested and will start taking data next year

24 MARCH, 2021 | By Anaïs Schaefer



The final elements of FASER were put into place this month. (Image: CERN)



“If you are the 100th person to look under a rock, you are unlikely to find anything...”



...but if you are using a new tool, or find a rock that's been left unturned, you don't have to be smart to find something”




Dr. Steven
Chu

Main Questions

1. Can we tune Pythia for forward experiments?
2. Which experiments can we hope to tune to?
3. How can we estimate the uncertainties in our tune?




High Energy Physics - Experiment
 [Submitted on 13 May 2021]
First neutrino interaction candidates at the LHC




Main Problem

- Usual Pythia tunes don't describe LHCf data
 - * other generators aren't that great either ...




default tunes

LHCf pion spectrum




LHCf neutron spectrum



plots made with Rivet

Importance for Forward Neutrino Fluxes



Most neutrinos come from pion / kaon decays inside the LHC's beam pipe.
→ Neutrino spectrum sensitive to forward pion / kaon production.

Main Questions

1. Can we tune Pythia for forward experiments?
2. Which experiments can we hope to tune to?
3. How can we estimate the uncertainties in our tune?

Tuning Pipeline

PYTHIA



Generate Events
with sets of
tuning
parameters

Tuning Pipeline

PYTHIA



Generate Events
with sets of
tuning
parameters



Rivet — the particle-physics MC analysis toolkit

Combine experimental analyses with Pythia simulations



YODA — Yet more Objects for Data Analysis
small, mean and full of Jedi magic

Tuning Pipeline

PYTHIA



Generate Events
with sets of
tuning
parameters



YODA — Yet more Objects for Data Analysis
small, mean and full of Jedi magic

Rivet — the particle-physics MC analysis toolkit

Combine experimental analyses with Pythia simulations

pyapprentice 1.0.6


Predict Pythia response
to tuning parameters
and fit to experimental
data

Main Questions


1. Can we tune Pythia for forward experiments?
2. Which experiments can we hope to tune to?
3. How can we estimate the uncertainties in our tune?

Before Tuning

Poor predictions in the forward region.





After Tuning



Poor predictions in the forward region.

An improvement, but some analyses are problematic

How much forward physics data can we fit at once?



Poor predictions in the forward region.

An improvement, but some analyses are problematic

Problematic analyses *can* be generated by Pythia

Main Questions

1. Can we tune Pythia for forward experiments?
2. Which experiments can we hope to tune to?
3. How can we estimate the uncertainties in our tune?


Forward Experiments

LHCf ($\eta > 8.81$)				
Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
forward π^0 or γ	7	✓	[1]	LHCF_2012_I1115479
	2.76, 7	✓	[2]	LHCF_2016_I1385877
	13	✓	[3]	LHCF_2018_I1518782
forward γ (diffractive)	13	—	[4]	—
forward neutrons	7	✓	[5]	LHCF_2015_I1351909
	13	✓	[6]	LHCF_2018_I1692008
	13	—	[7]	

CASTOR ($5.2 < \eta < 6.6$)				
Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
forward E	13	✓	[14]	CMS_2017_I1511284
forward E vs central N_{ch}	0.9, 2.76, 7	✓	[15]	CMS_2013_I1218372
	13	—	[16]	CMS_2019_I1747892 ⁽¹⁾
$dE/d\eta$	13	—	[17]	CMS_2018_I1708620

TOTEM (L2) ($5.3 < \eta < 6.5$)				
Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
$dN_{ch}/d\eta$	7	✓	[10]	TOTEM_2012_I1115294
	8	✓	[11]	TOTEM_2014_I1328627
	8	✓	[12]	CMSTOTEM_2014_I1294140
σ_{DD}	7	✓	[13]	

Analyses Targeting Diffractive Processes



ATLAS ($|\eta| < 5$)

Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
MB: dN_{ch} , η and pT	0.9, 2.36, 7	✓	[22]	ATLAS_2010_S8918562
	8	✓	[23]	ATLAS_2016_11426695
	13	✓	[24]	ATLAS_2016_11419652
MB: $\sum E_T$	7	✓	[25]	ATLAS_2012_11183818
	σ_{inel}	7	✓	[26]
13		✓	[27]	ATLAS_2016_11468167
η gap	7	✓	[28]	ATLAS_2012_11084540
	7	✓	[29]	
ALFA: tagged p SD (unpublished)	8	✓	[30]	ATLAS_2019_11762584 ⁽¹⁾
	13		[31]	

ALICE ($|\eta| < 5$)

Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
$\sigma_{SD}, \sigma_{DD}, \sigma_{inel}$	7	✓	[18]	ALICE_2012_11181770
incl. photons	0.9, 2.76, 7	✓	[19]	—
N_{ch}	0.9, 7, 8 TeV	✓	[20]	—
ϕ	2.76 TeV	✓	[21]	—


CMS ($|\eta| < 5$)

Analysis	\sqrt{s} [TeV]	HD	Refs.	RIVET
η gap	7	—	[32]	CMS_2015_11356998
TOTEM SD σ_{2j} w. tagged p	8	✓	[33]	
Strange Production	0.9, 7	✓	[34]	CMS_2011_S8978280
	13	✓	[35]	CMS_2017_11608166 ⁽¹⁾
σ_{inel} (incl. SD enhanced) diffractive (unpublished)	13	✓	[36]	CMS_2018_11653948
	7	—	[37]	
	8	—	[38]	
	13	—	[39]	

Main Questions

1. Can we tune Pythia for forward experiments?
2. Which experiments can we hope to tune to?
3. How can we estimate the uncertainties in our tune?

Estimating Uncertainty




- Naively, one could take the error band defined by multiple generators' predictions


This is problematic:
uncertainty strongly depends
on the weakest generator

Want something more robust


Estimating Uncertainty - Replica Tunes



1. Create replica datasets
2. Create replica *tunes* from these
3. Use these replica tunes to explore parameter space effectively







Estimating Uncertainty - Replica Tunes



- Preliminary error bars

Main Questions

1. Can we tune Pythia for forward experiments? 
2. Which experiments can (or should) we tune to?
Are there other experiments we should focus on?
Are there other experiments we are ignoring?
3. How can we estimate the uncertainties in our tune?



Forward Physics



Dr. Steven
Chu

Thank You!

References

1. Faser_nu Technical Proposal: <https://arxiv.org/abs/1812.09139>
2. Faser Physics Reach for LLP's: <https://arxiv.org/abs/1811.12522>
3. Rivet - <https://rivet.hepforge.org/>
4. Apprentice - <https://iamholger.gitbook.io/apprentice/installation>
5. Pythia - <http://home.thep.lu.se/Pythia/>

Backup

First Results


- we tested 76 Pythia8 parameters, plotted key distribution, and identified 9 relevant parameters
- We tuned them to the LHCf pion and neutron analyses
- First results look promising

Parameters:

```
SigmaDiffractive:mMin  
SigmaDiffractive:lowMEnhance  
SigmaDiffractive:maxAX  
SigmaDiffractive:maxXX  
SigmaDiffractive:mResMax  
SigmaDiffractive:maxXB  
SigmaDiffractive:maxAXB  
SigmaDiffractive:SaSepsilon  
StringPT:sigma
```

Analyses:


```
LHCF_2016_I1385877  
LHCF_2015_I1351909
```




Tuning Parameters



Parameter	Def.
SigmaDiffractive:maxXB	65
SigmaDiffractive:maxAX	65
SigmaDiffractive:maxXX	65
SigmaDiffractive:maxAXB	3.0
SigmaDiffractive:mMin	0.28
SigmaDiffractive:lowMEEnhance	2.0
SigmaDiffractive:mResMax	1.062
SigmaDiffractive:SaSepsilon	0.0
StringPT:sigma	0.335
Diffraction:mMinPert	10.
Diffraction:mWidthPert	10.
Diffraction:probMaxPert	1.0
Diffraction:pickQuarkNorm	5.0
Diffraction:pickQuarkPower	1.0
Diffraction:primKTwidth	0.5
Diffraction:largeMassSuppress	4.0
Diffraction:sigmaRefPomP	10.
Diffraction:mRefPomP	100
Diffraction:mPowPomP	0.0
SigmaDiffractive:PomFlux	1.0
SigmaDiffractive:PomFluxEpsilon	0.085
SigmaDiffractive:PomFluxAlphaPrime	0.25

TOTEM



LHCf Neutrons





LHCb

