

Twin Higgs Portal Dark Matter

Based on work with David Curtin, SG

2101.11019 [hep-ph]

— ONE SLIDE SUMMARY —

We extend the Twin Higgs framework to include a **singlet scalar particle** that acts like dark matter.

We treat the scalar as a **thermal relic** and compute its expected direct detection cross section

The pNGB nature of the (SM-like) Higgs **suppresses the direct detection cross section**, relative to benchmark Higgs portal WIMP models

— ONE SLIDE SUMMARY —

We extend the Twin Higgs framework to include a **singlet scalar particle** that acts like dark matter.

We treat the scalar as a **thermal relic** and compute its expected direct detection cross section

The pNGB nature of the (SM-like) Higgs **suppresses the direct detection cross section**, relative to benchmark Higgs portal WIMP models

+ extra suppression from asymmetric reheating!

TWIN HIGGS MODELS

Motivation:

Solve the (little) hierarchy problem via
color-neutral top partners

Lagrangian:

$$\mathcal{L} = \mathcal{L}_{\text{SM}_A} + \mathcal{L}_{\text{SM}_B}$$

- ▶ Complete or partial SM copy
- ▶ **singlet** under SM charges

$$\mathbb{Z}_2 : A \leftrightarrow B$$

$$H = (H_A, H_B)^T$$

$$V = \lambda \left(H^\dagger H - \frac{f_0^2}{2} \right)^2 + \text{breaking terms}$$

SU(4) symmetric

explicit or dynamical, required
for Higgs pheno

Twin Higgs Mechanism

1) $\langle H \rangle = \frac{1}{\sqrt{2}} f$  $SU(4) \rightarrow SU(3)$


125 GeV Higgs is pNGB of
broken $SU(4)$

Twin Higgs Mechanism

1) $\langle H \rangle = \frac{1}{\sqrt{2}} f \quad \longrightarrow \quad SU(4) \rightarrow SU(3)$

125 GeV Higgs is pNGB of broken $SU(4)$

2)



$$\Delta V = \frac{3}{8\pi^2} \Lambda^2 \left(\lambda_A^2 H_A^\dagger H_A + \lambda_B^2 H_B^\dagger H_B \right) = \frac{3\lambda^2}{8\pi^2} \Lambda^2 H^\dagger H$$

accidental $SU(4)$ symmetry!

125 GeV Higgs protected at 1 loop

Introducing:

TWIN HIGGS PORTAL DARK MATTER

Start with Twin Higgs potential:

$$V = \lambda \left(H_A^\dagger H_A + H_B^\dagger H_B - \frac{f_0^2}{2} \right)^2 + \kappa \left(\left(H_A^\dagger H_A \right)^2 + \left(H_B^\dagger H_B \right)^2 \right) + \sigma f_0^2 H_A^\dagger H_A$$

SU(4) symmetric

explicit breaking terms

$$V = \lambda \left(H_A^\dagger H_A + H_B^\dagger H_B - \frac{f_0^2}{2} \right)^2 + \kappa \left(\left(H_A^\dagger H_A \right)^2 + \left(H_B^\dagger H_B \right)^2 \right) + \sigma f_0^2 H_A^\dagger H_A$$

$$+ \frac{1}{2} \mu_S^2 S^2 + \frac{1}{2} \lambda_{HS} S^2 \left(H_A^\dagger H_A + H_B^\dagger H_B \right)$$


Add a singlet scalar



$$V = \lambda \left(H_A^\dagger H_A + H_B^\dagger H_B - \frac{f_0^2}{2} \right)^2 + \kappa \left(\left(H_A^\dagger H_A \right)^2 + \left(H_B^\dagger H_B \right)^2 \right) + \sigma f_0^2 H_A^\dagger H_A$$

$$+ \frac{1}{2} \mu_S^2 S^2 + \frac{1}{2} \lambda_{HS} S^2 \left(H_A^\dagger H_A + H_B^\dagger H_B \right)$$

Get *suppressed coupling* to pNGB Higgs




↓

$$= i\lambda_{HS} \left(\frac{\kappa}{\lambda} \right) v \left(1 - 2v^2/f^2 \right) \left(1 - v^2/f^2 \right) + \mathcal{O} \left(\frac{\kappa^2}{\lambda^2} \right)$$

$$V = \lambda \left(H_A^\dagger H_A + H_B^\dagger H_B - \frac{f_0^2}{2} \right)^2 + \kappa \left(\left(H_A^\dagger H_A \right)^2 + \left(H_B^\dagger H_B \right)^2 \right) + \sigma f_0^2 H_A^\dagger H_A$$

$$+ \frac{1}{2} \mu_S^2 S^2 + \frac{1}{2} \lambda_{HS} S^2 \left(H_A^\dagger H_A + H_B^\dagger H_B \right)$$

Get *suppressed coupling* to pNGB Higgs



$$= i \lambda_{HS} \left(\frac{\kappa}{\lambda} \right) v \left(1 - 2v^2 / f^2 \right) \left(1 - v^2 / f^2 \right) + \mathcal{O} \left(\frac{\kappa^2}{\lambda^2} \right)$$


explicit $SU(4)$
breaking
explicit \mathbb{Z}_2 breaking
($v = f/\sqrt{2}$ in \mathbb{Z}_2 limit)

... proportional to explicit breaking parameters.

$$V = \lambda \left(H_A^\dagger H_A + H_B^\dagger H_B - \frac{f_0^2}{2} \right)^2 + \kappa \left(\left(H_A^\dagger H_A \right)^2 + \left(H_B^\dagger H_B \right)^2 \right) + \sigma f_0^2 H_A^\dagger H_A$$

$$+ \frac{1}{2} \mu_S^2 S^2 + \frac{1}{2} \lambda_{HS} S^2 \left(H_A^\dagger H_A + H_B^\dagger H_B \right)$$

Get *suppressed coupling* to pNGB Higgs



... proportional to explicit breaking parameters.

Why does this matter?



Why does this matter?

- We treat dark matter as a thermal relic
- pNGB Higgs mediates *both freeze out and direct detection*

Why does this matter?

- We treat dark matter as a thermal relic
- pNGB Higgs mediates *both freeze out and direct detection*



Low energy ($m_S \ll m_h$):



Why does this matter?

- We treat dark matter as a thermal relic
- pNGB Higgs mediates *both freeze out and direct detection*

Low energy ($m_S \ll m_h$):





Same small coupling mediates **both**

Why does this matter?

- We treat dark matter as a thermal relic
- pNGB Higgs mediates *both freeze out and direct detection*



High energy ($m_S \gtrsim m_h$):



Why does this matter?

- We treat dark matter as a thermal relic
- pNGB Higgs mediates *both freeze out and direct detection*


High energy ($m_S \gtrsim m_h$):



Reduced direct detection

$$\Omega_{\text{DM}} h^2 = 0.120 \pm .001$$


Planck 2018, astro-ph/1807.06209



Reduced direct detection

$$\Omega_{\text{DM}} h^2 = 0.120 \pm .001$$

Planck 2018, astro-ph/1807.06209




- Freezes out via λ_{SSh}
- DD proceeds via λ_{SSh}

Maps on to regular Higgs portal WIMP scenario

Reduced direct detection

$$\Omega_{\text{DM}} h^2 = 0.120 \pm .001$$

Planck 2018, astro-ph/1807.06209




- Freezes out via λ_{SSH}
- DD proceeds via λ_{SSH}

Maps on to regular Higgs portal WIMP scenario

- Freezes out through unsuppressed coupling λ_{HS}
- DD proceeds via λ_{SSH}

Direct detection suppressed — possibilities for **next gen** experiments!


Is there reason to assume we live here?



- Freezes out through unsuppressed coupling λ_{HS}
- DD proceeds via λ_{SSH}

Direct detection possible in
next generation experiments!

Is there reason to assume we live here?




Yes! Natural mass scale for S is $\mathcal{O}(f)$:

$$V_S = \frac{1}{2}\mu_S^2 S^2 + \frac{1}{2}\lambda_{HS} S^2 (H_A^\dagger H_A + H_B^\dagger H_B)$$

- want **natural** tree level mass: $m_S^2 = \mu^2 + \lambda_{HS} f^2$
- unsuppressed radial mode coupling generates $\mathcal{O}(f)$ mass corrections at loop level

Is there reason to assume we live here?



Vast majority of unexplored parameter space corresponds to most natural DM mass scale!

Last point:

Last point:

Some Twin Higgs models (e.g. MTH) have spectra in tension with ΔN_{eff} constraints

This can be solved by **diluting** the twin radiation through late-time decays to mostly SM final states, known as **asymmetric reheating**

This dilutes the DM, further reducing the expected coupling:

Last point:

Some Twin Higgs models (e.g. MTH) have spectra in tension with ΔN_{eff} constraints

This can be solved by **diluting** the twin radiation through late-time decays to mostly SM final states, known as **asymmetric reheating**

This dilutes the DM, further reducing the expected coupling:


$$\Omega \rightarrow \Omega/D \quad \longrightarrow \quad \lambda_{HS} \rightarrow \lambda_{HS}/\sqrt{D}$$

Last point:

Some Twin Higgs models (e.g. MTH) have spectra in tension with ΔN_{eff} constraints

This can be solved by **diluting** the twin radiation through late-time decays to mostly SM final states, known as **asymmetric reheating**

This dilutes the DM, further reducing the expected coupling:



THPDM + AR:
cosmologically valid
for all twin spectra!

CONCLUSION

We extended the Twin Higgs framework to include a **singlet scalar DM candidate**

CONCLUSION

We extended the Twin Higgs framework to include a **singlet scalar DM candidate**

The **pNGB nature** of the Higgs leads to **suppressed direct detection** signals for the most natural DM mass scale

CONCLUSION

We extended the Twin Higgs framework to include a **singlet scalar DM candidate**

The **pNGB nature** of the Higgs leads to **suppressed direct detection** signals for the most natural DM mass scale

CONCLUSION

The model is valid for **any Twin Higgs spectrum** — incorporating asymmetric reheating further suppresses direct detection

We extended the Twin Higgs framework to include a **singlet scalar DM candidate**

The **pNGB nature** of the Higgs leads to **suppressed direct detection** signals for the most natural DM mass scale

CONCLUSION

The model is valid for **any Twin Higgs spectrum** — incorporating asymmetric reheating further suppresses direct detection

Thanks!
hep-ph/2101.11019

EXTRAS

ASYMMETRIC REHEATING

Cosmology issues in Twin Higgs

- DD suppression valid for **all** Twin Higgs models...
- ... but some models (e.g. MTH) have spectra in tension with ΔN_{eff}
- Solve with *asymmetric reheating*:

nuMTH: Z. Chacko, N. Craig, P. J. Fox, R. Harnik [hep-ph/1611.07975](#)

XMTH: N. Craig, S. Koren, T. Trott [hep-ph/1611.07977](#)



Cosmology issues in Twin Higgs

- DD suppression valid for **all** Twin Higgs models...
- ... but some models (e.g. MTH) have spectra in tension with ΔN_{eff}
- Solve with *asymmetric reheating*:

nuMTH: Z. Chacko, N. Craig, P. J. Fox, R. Harnik [hep-ph/1611.07975](#)

XMTH: N. Craig, S. Koren, T. Trott [hep-ph/1611.07977](#)

- Introduce **weakly interacting**, massive, **long-lived** particle with **preferential decays** to SM
- **Freezes out relativistically**, dominates cosmology at late times
- **Decays** at late times
- **Dilutes twin sector** radiation contribution to ΔN_{eff}

Cosmology issues in Twin Higgs

- DD suppression valid for **all** Twin Higgs models...
- ... but some models (e.g. MTH) have spectra in tension with ΔN_{eff}
- Solve with *asymmetric reheating*:

nuMTH: Z. Chacko, N. Craig, P. J. Fox, R. Harnik [hep-ph/1611.07975](#)


XMTH: N. Craig, S. Koren, T. Trott [hep-ph/1611.07977](#)


Diluting twin radiation **dilutes DM**:

$$\Omega \rightarrow \Omega/D$$

Further reduces DD predictions:

$$\lambda_{HS} \rightarrow \lambda_{HS}/\sqrt{D}$$





Twin Higgs Portal DM
+
asymmetric reheating



Model of DM that is cosmologically valid for **any** twin spectrum