Phenomenology 2021 Symposium



Contribution ID: 1267 Type: Cosmology

The Hubble Constant in the Axi-Higgs Universe

Monday 24 May 2021 18:30 (15 minutes)

The Λ CDM model provides an excellent fit to the CMB data. However, a statistically significant tension emerges when its determination of the Hubble constant H_0 is compared to the local distance-redshift measurements. The axi-Higgs model, which couples ultralight axions to the Higgs field, offers a specific variation of the Λ CDM model. It relaxes the H_0 tension as well as explains the 7 Li puzzle in Big-Bang nucleosynthesis, the S_8 tension with the weak-lensing data, and the observed isotropic cosmic birefringence in CMB. In this letter, we demonstrate how the H_0 and S_8 tensions can be resolved simultaneously, by correlating the axion impacts on the early and late universe. In a benchmark scenario selected for experimental tests soon, the analysis combining the CMB+BAO+WL+SN data yields $H_0=71.1\pm1.1$ km/s/Mpc and $S_8=0.766\pm0.011$. Combining this (excluding the SN(supernovae) part) with the local distance-redshift measurements yields $H_0=72.3\pm0.7$ km/s/Mpc, while S_8 is unchanged.

Summary

Authors: LUU, Hoang Nhan (The Hong Kong University of Science and Technology); Dr LI, Lingfeng (The Hong Kong University of Science and Technology); Prof. TYE, S.-H. Henry (The Hong Kong University of Science and Technology); Prof. LIU, Tao (The Hong Kong University of Science and Technology); FUNG, WH Leo (The Hong Kong University of Science and Technology); QIU, Yu-Cheng (The Hong Kong University of Science and Technology)

Presenter: LUU, Hoang Nhan (The Hong Kong University of Science and Technology)

Session Classification: Cosmology II