



PHENO 2021

The 2021 Phenomenology Symposium

24th – 26th May 2021

University of Pittsburgh

ATLAS Measurements of CP-Violation and Rare Decays Processes with Beauty Mesons



Pavel Řezníček for the ATLAS Collaboration

24th May 2021




B-Physics at ATLAS

- ATLAS Run 2: 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ collected in 2015-2018
- Producing 2.5M $b\bar{b}$ pairs/second, B_s , B_c , Λ_b , etc. available
- Program focused mostly on muonic final states, fully reconstructable
- Typical trigger: low- p_T di-muons at low invariant mass, using information from tracker and muon detectors
- B-physics trigger rate up to $\sim 200 \text{ Hz}$



- ATLAS Run 2: 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ collected in 2015-2018
- Producing 2.5M $b\bar{b}$ pairs/second, B_s , B_c , Λ_b , etc. available
- Program focused mostly on muonic final states, fully reconstructable
- Typical trigger: low- p_T di-muons at low invariant mass, using information from tracker and muon detectors
- B-physics trigger rate up to $\sim 200 \text{ Hz}$




**Measurement of the CP -violating phase ϕ_s in $B_s^0 \rightarrow J/\psi\phi$ decays
in ATLAS at 13 TeV**

Eur. Phys. J. C 81 (2021) 342

CPV in $B_s^0 \rightarrow J/\psi\phi$ and the measurement


- Interference of direct decay and decay with mixing into the same final state of $B_s^0 \rightarrow J/\psi\phi$ gives rise to time-dependent CP violation



- In the Standard Model (SM) the ϕ_s is small: $\phi_s \simeq -2\beta_s = -0.03696^{+0.00072}_{-0.00082}$ rad
- New Physics (NP) could contribute to the mixing box diagrams, potentially enlarging ϕ_s
- Whole system described by:
 - weak phase ϕ_s and direct-CPV parameter λ
 - CP-state amplitudes (and their phases)
 - the mixing parameters Δm_s , $\Delta\Gamma_s$, Γ_s

Measurement

- Final state: admixture of CP-odd ($L = 1$) and CP-even ($L = 0, 2$) states
- Distinguishable through time-dependent angular analysis: $\frac{d^4\Gamma}{dt d\Omega} = \sum_{k=1}^{10} \mathcal{O}^{(k)}(t) g^{(k)}(\theta_T, \psi_T, \phi_T)$
- Analyzing signal final state $B_s^0 \rightarrow J/\psi(\mu^+\mu^-)\phi(K^+K^-)$
- Non-resonant S-wave decay $B_s^0 \rightarrow J/\psi K^+K^-$ contribution included in the differential decay rate




Opposite-side flavour tagging

- Use $b - \bar{b}$ correlation \Rightarrow initial B_s^0 flavour
 - $b(\bar{b}) \rightarrow l^{-(+)} \text{ transition}$
 - diluted by oscillations and $b \rightarrow c \rightarrow l$
- Key variables: charge of p_T -weighted tracks in cone $\Delta R(\phi, \eta)$ around the opposite side lepton

$$Q_x = \frac{\sum_i^N \text{tracks} q_i \cdot (p_{Ti})^\kappa}{\sum_i^N (p_{Ti})^\kappa}$$


- Building per-candidate tag probability $P(B|Q)$



Four taggers



- **Muon:** tight-ID or low- p_T , $\kappa = 1.1$, $\Delta R = 0.5$
- **Electron:** $p_T(e) > 0.5 \text{ GeV}$, $\kappa = 1.0$, $\Delta R = 0.5$
- **Jet:** b -tagged jets, $\kappa = 1.1$, $\Delta R = 0.5$
- Search order based on best purity

- Calibrated on self-tagged $B^\pm \rightarrow J/\psi K^\pm$ data



Tagging performance

Tag method	$\epsilon_x [\%]$	$D_x [\%]$	$T_x [\%]$
Tight muon	4.50 ± 0.01	43.8 ± 0.2	0.862 ± 0.009
Electron	1.57 ± 0.01	41.8 ± 0.2	0.274 ± 0.004
Low- p_T muon	3.12 ± 0.01	29.9 ± 0.2	0.278 ± 0.006
Jet	12.04 ± 0.02	16.6 ± 0.1	0.334 ± 0.006
Total	21.23 ± 0.03	28.7 ± 0.1	1.75 ± 0.01



Unbinned maximum likelihood fit


- An unbinned maximum likelihood (UML) fit performed in 10 D space

$$\ln \mathcal{L} = \sum_{i=1}^N \{ w_i \cdot \ln(f_s \mathcal{F}_s + f_s f_{B^0} \mathcal{F}_{B^0} + f_s f_{\Lambda_b} \mathcal{F}_{\Lambda_b} + (1 - f_s(1 + f_{B^0} + f_{\Lambda_b})) \mathcal{F}_{\text{bkg}}) \}$$

Observables

$\mathcal{F}_x(m_i, t_i, \sigma_{m_i}, \sigma_{t_i}(p_{T_i}), \theta_T, \psi_T, \phi_T, P(B|Q_i))$

- Base B_s^0 decay observables: mass, time, angles
 - Conditional observables: per-candidate tagging Q_x and mass/time resolutions ($p_T(B)$ dependent)
- Full time-angular PDF including S-wave
- Fixed parameters: $\Delta m_s = \text{PDG}$, direct CP $\lambda = 1$
- Trigger causing decay time inefficiency, modeled in MC




Results


Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.081	0.041	0.022
$\Delta\Gamma_s$ [ps $^{-1}$]	0.0607	0.0047	0.0043
Γ_s [ps $^{-1}$]	0.6687	0.0015	0.0022
$ A_s(0) ^2$	0.2213	0.0019	0.0023
$ A_0(0) ^2$	0.5131	0.0013	0.0038
$ A_S(0) ^2$	0.0321	0.0033	0.0046
$\delta_\perp - \delta_S$ [rad]	-0.25	0.05	0.04
Solution (a)			
δ_\perp [rad]	3.12	0.11	0.06
δ_\parallel [rad]	3.35	0.05	0.09
Solution (b)			
δ_\perp [rad]	2.91	0.11	0.06
δ_\parallel [rad]	2.94	0.05	0.09

- Almost 500k signal candidates
- Weak phase ϕ_s as well as decay width difference $\Delta\Gamma_s$ compatible with Standard Model
- Dominant systematics on ϕ_s measurement from tagging
 - Accounting for pile-up dependence, calibration curves model and MC precision, "Punzi" PDFs variations, difference between B^\pm and B_s^0 kinematics

- Two solutions in $\delta_\parallel - \delta_\perp$ plane, with negligible impact on other parameters





- Statistical (BLUE) combination with Run 1 result



Results

Parameter	Value	Statistical uncertainty	Systematic uncertainty
		0.001	0.004

Comparison with other experiments



- Dominant systematics on ϕ_s measurement from tagging


- Accounting for pile-up dependence, calibration curves model and MC precision, "Punzi" PDFs variations, difference between B^\pm and B_s^0 kinematics

**Study of the rare decays of B_s^0 and B^0 mesons into muon pairs
using data collected during 2015 and 2016 with the ATLAS detector**

JHEP 04 (2019) 098

Analysis of rare $B_{(s)}^0 \rightarrow \mu\mu$ decays


- FCNC in the SM proceeding via loop and box diagrams, and helicity suppressed $\implies \mathcal{B} \sim 10^{-9}$
- BSM can significantly contribute, modifying the branching ratio



Measurement

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = N_{d(s)} \cdot \frac{\mathcal{B}(B^\pm \rightarrow J/\psi K^\pm) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)}{N_{J/\psi K^\pm} \cdot \frac{\epsilon_{\mu^+ \mu^-}}{\epsilon_{J/\psi K^\pm}}} \cdot \frac{f_u}{f_{d(s)}}$$

- $\mathcal{B}(B_{(s)}^0 \rightarrow \mu\mu)$ measurement relative to $\mathcal{B}(B^\pm \rightarrow J/\psi K^\pm)$, $B_s^0 \rightarrow J/\psi \phi$ as control channel
- Blinded signal di-muon invariant mass region [5166, 5526] MeV
- BDT based background suppression, trained on sidebands data
- Yields $N_{d(s)}$ and $N_{J/\psi K^\pm}$ obtained from UML fits to the mass spectra
- Relative reconstruction efficiencies estimated from MC (corrected for data-MC differences): $\epsilon_{\mu^+ \mu^-} / \epsilon_{J/\psi K^\pm} = 0.1176 \pm 0.0009_{\text{stat.}} \pm 0.0047_{\text{syst.}}$
- Known branching ratios from PDG, $f_u/f_{d(s)}$ from HFLAV




- $B^\pm \rightarrow J/\psi K^\pm$ yield:
 $33435 \pm 0.3\%_{\text{stat.}} \pm 4.8\%_{\text{syst.}}$

Backgrounds


Partially reconstructed b -hadron decays

- Mostly in the low di-muon mass region
- Shape free in the mass fit




Peaking backgrounds

- Hadronic B_s^0 decays where hadrons are misidentified as muons
- Simulated and fixed in the mass fit





Continuum background

- Combinatorics of μ and uncorrelated hadron decays
- Reduced by BDT
- Linear shape constrained in the mass fit across BDT bins
- Systematics due to $B_c^\pm \rightarrow J/\psi \mu \nu$ and $B_{(s)}^0/\Lambda_b^0 \rightarrow h \mu \nu$ decays




BDT and signal yield extraction

- BDT formed from 15 variables
 - kinematics, isolation, B -vertex separation from PV
- BDT output validated on reference $B^\pm \rightarrow J/\psi K^\pm$ and control $B_s^0 \rightarrow J/\psi \phi$ channels, observed difference applied as a correction to signal channel
- Signal region divided into four BDT bins with constant signal efficiency
- Simultaneous extraction of $B_s^0 \rightarrow \mu\mu$ and $B^0 \rightarrow \mu\mu$ yields from unbinned maximum likelihood fit to di-muon mass distributions in the four BDT bins



Results

- Contours obtained using Neyman construction



- Combined measurement compatible with SM at 2.4σ
- Statistic uncertainties dominate
- Largest systematics contribution from di-muon mass fit procedure

Standard Model

$$\mathcal{B}(B_s^0 \rightarrow \mu\mu) = (3.66 \pm 0.14) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu\mu) = (1.03 \pm 0.05) \times 10^{-10}$$

ATLAS 2015 + 2016 data


$$\mathcal{B}(B_s^0 \rightarrow \mu\mu) = (3.2^{+1.1}_{-1.0}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu\mu) < 4.3 \times 10^{-10} \text{ at 95% CL}$$


ATLAS Run 1 + 2015 + 2016 data

$$\mathcal{B}(B_s^0 \rightarrow \mu\mu) = (2.8^{+0.8}_{-0.7}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu\mu) < 2.1 \times 10^{-10} \text{ at 95% CL}$$



- Combining binned 2D profile likelihoods,
 f_s/f_d the only source of correlation between experiments



Summary

- Latest ATLAS measurements of CP -violation in $B_s^0 \rightarrow J/\psi\phi$ decay and branching ratio measurement of rare $B_{(s)}^0 \rightarrow \mu\mu$ decays compatible with Standard Model predictions
- Full Run 2 data analyses in progress
 - CPV measurement releasing Δm_s and direct- CP λ , improvements in tagging and fit model
 - Rare decays including $B_s^0 \rightarrow \mu\mu$ lifetime analysis
- Program continuation in Run 3 and HL-LHC
 - HL-LHC projections [CERN Yellow Report Monograph 7 \(2019\) pp. 1–1418](#))

