

# Gamma-Ray Bursts at high and very high energies

*Lara Nava*


*Marie Skłodowska-Curie Fellow*

*INAF*

*Osservatorio Astronomico di Brera - Osservatorio Astronomico di Trieste*



# GRBs: powerful explosions at cosmological distances



## PROMPT emission

internal shocks or reconnection



Particle acceleration: non-thermal  
Radiative process: synchrotron??

Duration: seconds

Energy range: 10 keV - 10 MeV

## AFTERGLOW emission

external shocks




Particle acceleration: non-thermal  
Radiative process: synchrotron

Duration: weeks-months

Energy range: radio to soft X-rays

# GRBs: powerful explosions at cosmological distances



## PROMPT emission

internal shocks or reconnection



Particle acceleration: non-thermal  
Radiative process: synchrotron??

Duration: seconds

Energy range: 10 keV - 10 MeV

## AFTERGLOW emission

external shocks





Particle acceleration: non-thermal  
Radiative process: synchrotron

Duration: weeks-months

Energy range: radio to soft X-rays

# GRBs: powerful explosions at cosmological distances



## PROMPT emission

internal shocks or reconnection



Particle acceleration: non-thermal  
Radiative process: synchrotron??

Duration: seconds

Energy range: 10 keV - 10 MeV

## AFTERGLOW emission

external shocks




Particle acceleration: non-thermal  
Radiative process: synchrotron

Duration: weeks-months

Energy range: radio to soft X-rays

# GRBs: powerful explosions at cosmological distances



## PROMPT emission

internal shocks or reconnection



Particle acceleration: non-thermal  
Radiative process: synchrotron??

Duration: seconds

Energy range: 10 keV - 10 MeV

## AFTERGLOW emission

external shocks




Particle acceleration: non-thermal  
Radiative process: synchrotron

Duration: weeks-months

Energy range: radio to soft X-rays

# GRBs: powerful explosions at cosmological distances



## PROMPT emission

internal shocks or reconnection



Particle acceleration: non-thermal  
Radiative process: synchrotron??

Duration: seconds

Energy range: 10 keV - 10 MeV

## AFTERGLOW emission


external shocks




Particle acceleration: non-thermal  
Radiative process: synchrotron


Duration: weeks-months

Energy range: radio to soft X-rays




# The GRB electromagnetic spectrum







# The GRB electromagnetic spectrum








# The GRB electromagnetic spectrum




**EGRET, AGILE, Fermi-LAT**

**20 MeV - 300 GeV**

**130 GRBs detected in 9yr**




# The GRB electromagnetic spectrum




**EGRET, AGILE, Fermi-LAT**  
20 MeV - 300 GeV  
130 GRBs detected in 9yr


**MAGIC, HESS, VERITAS**  
50-100 GeV — 100 TeV  
no detections!



# High-energy emission: Fermi-LAT observations


High-energy LAT lightcurves  
for 10 GRBs, 0.1-10 GeV






# High-energy emission: Fermi-LAT observations

## High-energy LAT lightcurves for 10 GRBs, 0.1-10 GeV



## Optical lightcurves



# High-energy emission: modelling and interpretation


Issue with an external shock- synchrotron interpretation  
of the GeV emission

Electrons are  
accelerated up to  
some maximum  
energy  $\gamma_{\max}$



Synchrotron photons  
are emitted up to a  
maximum energy  
 $E_{\text{syn,max}}$

$$E_{\text{syn,max}} \propto \gamma_{\max}^2 B \Gamma$$
$$\approx 50 \text{ MeV} \times \Gamma$$



# High-energy emission: modelling and interpretation


Issue with a external shock- synchrotron interpretation  
of the GeV emission

Electrons are  
accelerated up to  
some maximum  
energy  $\gamma_{\max}$




Synchrotron photons  
are emitted up to a  
maximum energy  
 $E_{\text{syn,max}}$

$$E_{\text{syn,max}} \propto \gamma_{\max}^2 B \Gamma$$
$$\approx 50 \text{ MeV} \times \Gamma$$




SSC ???  
(synchrotron  
self-Compton)

# Inverse Compton component?





# Inverse Compton component?





# Inverse Compton component?



Tam et al. 2013



# Why should we care about what happens at high energies?

- ◆ nature of radiative processes
- ◆ energetics and efficiencies
- ◆ magnetic fields
- ◆ particle acceleration
- ◆ EBL (extragal. background light)
- ◆ quantum gravity

GRB physics


more general  
relevance



# Why should we care about what happens at high energies?


## CTA

Cherenkov Telescope Array



Large-Sized Telescopes (LST)  
30 - 200 GeV


Simulation GRB spectrum




huge number of events!

...but small detection rate:  
0.1 - 1 per year


# Inverse Compton component?



# Inverse Compton component?




# Inverse Compton component?



Prospects for GRB detection with the CTA  
...work in progress...



# What are we doing in Trieste



# Ongoing activity and future prospects

- ◆ Development of a code for modeling afterglow emission describing synchrotron and SSC emission, Klein-Nishina, pair production. Solution of two coupled equations for evolution of particle and photon spectra


Two examples of syn+SSC spectra and lightcurves from the code



- ◆ Application to Fermi-LAT GRBs: can high-energy photons be explained by SSC radiation?
- ◆ CTA GRB detection rate



# The collaboration




Thank you for the attention!



# High-energy emission: modelling and interpretation

Consequences on the inferred prompt mechanism efficiency!!




inferred from X-rays:  
internal shocks ruled out!

jet must be  
magnetically  
dominated

inferred from GeV:  
internal shocks possible!


internal shocks  
(and matter  
dominated jets)  
are still viable

# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$

# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$

# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$





inconsistency  
between X-ray  
and GeV  
observations!?!?!?

# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$





inconsistency  
between X-ray  
and GeV  
observations!?!?!?

# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$





inconsistency  
between X-ray  
and GeV  
observations!?!?!?




# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$






inconsistency  
between X-ray  
and GeV  
observations!?!?!?



# High-energy emission: modelling and interpretation




afterglow flux measured at a frequency  
above  $\nu_c$  is a proxy for  $E_{\text{blastwave}}$



inconsistency  
between X-ray  
and GeV  
observations!?!?!?

# High-energy emission: modelling and interpretation

upper limits to  $\epsilon_B$



Density

traditionally assumed values  $> 10^{-2}$

SSC ???  
(synchrotron  
self-Compton)

similar results found by:  
Kumar & Barniol Duran 2009  
Lemoine 2013a/b  
Santana et al., 2014

Possible explanation for  
such small values of the  
magnetic field

Lemoine 2013, Lemoine+, 2013

