

# Indirect Detection of Secluded Supersymmetric Dark Matter

Patrick Barnes

<https://arxiv.org/abs/2003.13744>

<https://arxiv.org/abs/2106.09740>

Patrick Barnes, Zachary Johnson, Aaron Pierce, Bibhushan Shakya

December 4, 2021



# Brief Outline

Overview and Motivation

Photon Spectra from Dark Matter Annihilations

Analysis and Indirect Detection Bounds

## Secluded WIMPS and Indirect Detection

The WIMP paradigm remains a popular model of dark matter.

Traditional WIMP candidates, such as MSSM neutralinos, are increasingly bounded by direct detection experiments.

WIMP dark matter within a secluded sector with small portal couplings to the Standard Model can evade direct detection and collider bounds.


Indirect detection signals, however, will not be suppressed.

# Supersymmetric Secluded Sectors and Portals

Supersymmetry can explain why the secluded particles are at the weak scale.


A SUSY kinetic mixing provides a gauge, gaugino, and Higgs portal,

$$\frac{\epsilon}{2} \int d^2\theta W_Y W' + h.c. = \epsilon D_Y D' - \frac{\epsilon}{2} F_Y^{\mu\nu} F'_{\mu\nu} + i\epsilon \tilde{B} \sigma^\mu \partial_\mu \tilde{B}'^\dagger + i\epsilon \tilde{B}' \sigma^\mu \partial_\mu \tilde{B}^\dagger.$$



## R-Parity Violation

R-Parity is sometimes postulated in the MSSM to stabilize the LSP.



We can add R-Parity violating couplings and investigate the results of different ones on our annihilation spectra.

# Photon Spectra from Annihilation

For R-Parity even final states, we have Dirac DM  $\psi$ , a dark photon  $Z'$ , and dark Higgs  $H'$ . We do not assume supersymmetry.

$$\psi\bar{\psi} \rightarrow Z'H' \text{ (Higgs Mechanism)}$$


$$\psi\bar{\psi} \rightarrow Z'Z' \text{ (Stueckelberg)}$$

$$SS^* \rightarrow H'H' \text{ (Scalar DM)}$$

Branching ratios set by

$$\mathcal{L} = \xi |H'|^2 |H|^2 - \frac{\epsilon}{2} F_Y^{\mu\nu} F'_{\mu\nu}. \quad (1)$$

# R-Parity Even Final States



**Figure 1: Left:** Spectra for  $\psi\bar{\psi}$  annihilation to either  $Z'Z'$ , or  $Z'H'$  in the degenerate case  $m_{Z'} = m_{H'}$ . **Right:** We now allow  $m_{H'}' \neq m_{Z'}$ .

$$H' \rightarrow b\bar{b} \text{ or } W^+W^-$$

$$Z' \rightarrow u\bar{u} \text{ and } \ell^+\ell^-$$

## R-Parity Odd Final States



If the secluded sector is supersymmetric, annihilation to neutralinos,  $\psi\bar{\psi} \rightarrow \chi'_1\chi'_1$ , is possible.

We assume  $H'$  is charged under  $U(1)'$ , so the Higgsino and gaugino mix to form Majorana mass eigenstates  $\chi'_1$  and  $\chi'_2$ .

These neutralinos will decay to SM states through the gaugino portal.




# LSP in the Visible Sector



**Figure 2: Left:** Effective DM annihilation through a neutralino cascade. “RPV” indicates the three fermion final state from RPV  $\chi_1$  decay, which differs based on the dominant RPV coupling. **Right:** The resulting spectra for specific examples of non-zero RPV couplings.

$$W_{RPV} = \frac{1}{2} \lambda_{ijk} L_i L_j E_k + \lambda'_{ijk} L_i Q_j D_k^c + \frac{1}{2} \lambda''_{ijk} U_i^c D_j^c D_k^c. \quad (2)$$

# LSP in the Secluded Sector



**Figure 3:** The photon spectra for direct  $\psi$  annihilation to  $\chi'_1$ , shown for multiple potential RPV mediated  $\chi'_1$  decays.

If the  $\chi'_1$  is lighter than its MSSM counterparts, it may decay directly to the SM via RPV couplings.

# Analysis



Fermi-LAT

6 years of data

15 dSph galaxies

<https://arxiv.org/abs/1503.02641>





CTA

Projected 525 hours


Milky Way galactic center

<https://arxiv.org/abs/2007.16129>



# R-Parity Even Final States



# R-Parity Even Final States



# R-Parity Odd Final States



## Takeaways

Indirect detection can provide a robust probe of DM models where small couplings will suppress direct and collider signals.

A well motivated example is a supersymmetric secluded sector.

For large areas of parameter space, CTA will probe the thermal relic cross section for such a model.