



Contribution ID: 183

Type: not specified

Inevitable Large non-Gaussianity from Curvaton Models

Tuesday 9 May 2023 15:00 (15 minutes)

Curvatons are light (compared to Hubble during inflation) spectator fields that potentially contribute adiabatic curvature perturbations post-inflation. They can thereby alter CMB observables such as the spectral index n_s , the tensor-to-scalar ratio r , and the local non-Gaussianity $f_{\text{NL}}^{(\text{loc})}$. We systematically explore the observable space of a curvaton with a quadratic potential. We find that when the underlying inflation model does not satisfy the n_s and r observational constraint but can be made viable with a significant curvaton contribution, a large $f_{\text{NL}}^{(\text{loc})}$ is inevitable without fine-tuning, therefore a lack of observation of said $f_{\text{NL}}^{(\text{loc})}$ in next generation experiments will rule out these models. On the other hand, when the underlying inflation model already satisfies the n_s and r observational constraint, $f_{\text{NL}}^{(\text{loc})}$ cannot be distinguished from the single-field inflation prediction in general.

Authors: LODMAN, Jacqueline (Harvard University); LU, Qianshu; RANDALL, lisa (harvard)

Presenter: LODMAN, Jacqueline (Harvard University)

Session Classification: BSM XIV