



Contribution ID: 116

Type: not specified

MAGPI: Measurement of Axion Gradients with Photon Interferometry

Tuesday 9 May 2023 18:15 (15 minutes)

I will present a novel search technique for axions with a CP-violating monopole coupling \tilde{g}_Q to bulk SM charges $Q \in \{B, L, B-L\}$. Gradients in the static axion field configurations sourced by matter induce achromatic circular photon birefringence via the axion-photon coupling $g_{\phi\gamma}$. Circularly polarized light fed into an optical or (open) radio-frequency (RF) Fabry-Perot (FP) cavity develops a phase shift that accumulates up to the cavity finesse: the fixed axion spatial gradient prevents a cancellation known to occur for an axion dark-matter search. The relative phase shift between two FP cavities fed with opposite circular polarizations can be detected interferometrically. This time-independent signal can be modulated up to non-zero frequency by altering the cavity orientations with respect to the field gradient. Multi-wavelength co-metrology techniques can be used to address chromatic measurement systematics and noise sources. I will discuss how, with Earth as the axion source, we project reach beyond current constraints on the product of couplings $\tilde{g}_Q g_{\phi\gamma}$ for axion masses $m_\phi < 10^{-5}$ eV. If shot-noise-limited sensitivity can be achieved, an experiment using high-finesse RF FP cavities could reach a factor of $\sim 10^6$ into new parameter space for $\tilde{g}_Q g_{\phi\gamma}$ for masses $m_\phi < 10^{-10}$ eV.

Authors: THOMPSON, Jedidiah O. (Stanford University); FEDDERKE, Michael A. (Johns Hopkins University)

Co-authors: GIACCONE, Bianca (Fermilab); KAPLAN, David E. (Johns Hopkins University); CERVANTES, Raphael (Fermilab); HARNIK, Roni (Fermilab); POSEN, Sam (Fermilab); RAJENDRAN, Surjeet (Johns Hopkins University)

Presenter: FEDDERKE, Michael A. (Johns Hopkins University)

Session Classification: Axion III

Track Classification: Axion and ALP