Contribution ID: 56

Type: Parallel Talk (Theory)

Torsional regularization and finite bare charge

Monday 26 November 2018 16:45 (15 minutes)

We show that in the presence of the torsion tensor S_{ij}^{k} , whose existence is required by the consistency of the conservation law for the total angular momentum of a Dirac particle in curved spacetime with relativistic quantum mechanics, the quantum commutation relation for the four-momentum is given by $[p_i, p_j] = 2i\hbar S_{ij}^{k}p_k$. We propose that this relation replaces the integration in the momentum space in Feynman diagrams with the summation over the discrete momentum eigenvalues.

We derive a prescription for this summation that agrees with convergent integrals:

$$\int \frac{d^4 p}{(p^2 + \Delta)^s} \to 4\pi U^{s-2} \sum_{l=1}^{\infty} \int_0^{\pi/2} d\phi \frac{\sin^4 \phi \, n^{s-3}}{[\sin \phi + U\Delta n]^s}$$

where $n = \sqrt{l(l+1)}$ and $1/\sqrt{U}$ is a constant on the order of the Planck mass, determined by the Einstein-Cartan theory of gravity.

We show that this prescription regularizes ultraviolet-divergent integrals in loop diagrams.

We extend this prescription to tensor integrals and apply it to vacuum polarization.

We derive a finite, gauge-invariant vacuum polarization tensor and a finite running coupling that agrees with the low-energy limit of the standard quantum electrodynamics.

Including loops from all charged fermions, we find a finite value for the bare electric charge of an electron: $\approx -1.22 e$.

Torsional regularization, originating from the noncommutativity of the momentum and spin-torsion coupling, therefore provides a realistic, physical mechanism for eliminating infinities in quantum field theory: quantum electrodynamics with torsion is ultraviolet complete.

arXiv

1712.09997

 Author:
 Dr POPLAWSKI, Nikodem (University of New Haven)

 Presenter:
 Dr POPLAWSKI, Nikodem (University of New Haven)

Session Classification: Parallel Talks B

Track Classification: QFT, Strings, AdS/CFT