Contribution ID: 66

## Visible neutrino decay and matter effects at future long-base line experiments

Thursday 29 November 2018 17:00 (15 minutes)

The consequences of introducing matter effects into the neutrino visible decay scheme are studied. To this end, we select two

baselines for which matter effects have to be considered:1300 km (DUNE) and 7650 km (considering an hypothetical

beam aimed towards ANDES). The matter effects are almost unnoticeable for the visible decay contribution DUNE, being sizable at ANDES. We also carry out a realistic analysis taking DUNE as a context, considering  $\nu_{\mu}$  disappearance and  $\nu_{e}$  appearance channels, for both FHC and RHC modes. The sensitivity to the decay constant  $\alpha_{3}$  can be as low as  $2 \times 10^{-6}$  eV<sup>2</sup> at 90% C.L., depending on the neutrino masses and type of coupling (scalar, pseudo-scalar or both). Lastly, we asses the impact of neutrino decay in the determination of  $\theta_{23}$  and  $\delta_{CP}$ , and find that the best-fit value of  $\theta_{23}$  can move from a true value at the lower octant towards the higher octant.

## arXiv

arXiv:1805.03279 [hep-ph]

**Authors:** ASCENCIO, Marvin (Pontificia Universidad Catolica del Perú); CALATAYUD CADENILLAS, Anthony Mard (Pontificia Universidad Católica del Perú); Dr GAGO MEDINA, Alberto Martín (Pontificia Universidad Católica del Perú); JONES-PEREZ, Joel (PUCP)

Presenter: CALATAYUD CADENILLAS, Anthony Mard (Pontificia Universidad Católica del Perú)

Session Classification: Parallel Talks A

Track Classification: Neutrinos