### PHOENIX-2023



Contribution ID: 44

Type: Talk

# Renormalization-group improved Higgs to two gluons decay rate

We investigate the renormalization group scale and scheme dependence of the  $H \rightarrow gg$  decay rate at the order N<sup>4</sup>LO in renormalization-group summed perturbative theory, which employs the summation of all renormalization-group accessible logarithms including the leading and subsequent four sub-leading logarithmic contributions to the full perturbative series expansion. The main advantage of this approach is the closed-form analytic expressions, which represent the summation of all RG-accessible logarithms in the perturbative series that is known to a given order. The new renormalization-group summed expansion for the  $H \rightarrow gg$  decay rate shows an improved behaviour by exhibiting a reduced sensitivity to the renormalization-group scale. Moreover, we study the higher-order behaviour of the  $H \rightarrow gg$  decay width using the asymptotic *Padé* approximant method in four different renormalization schemes. Furthermore, the higher-order behaviour is independently investigated in the framework of the asymptotic *Padé*-Borel approximant method where generalized Borel-transform is used as an analytic continuation of the original perturbative expansion. The predictions of the asymptotic *Padé*-Borel approximant method.

## Designation

Student

## **Reference** publication/preprint

arXiv.2205.06061v2 [hep-ph]

#### Institution

Indian Institute of Technology (BHU), Varanasi, 221005

Author: RES. SCHOLAR, PHYSICS, IIT(BHU), VARTIKA SINGH (Indian Institute of Technology, BHU)

**Co-authors:** Ms JAIN, Astha (Indian Institute of Technology (BHU)); Dr ABBAS, Gauhar (Indian Institute of Technology (BHU)); Ms SINGH, Neelam (Indian Institute of Technology (BHU))

Presenter: RES. SCHOLAR, PHYSICS, IIT(BHU), VARTIKA SINGH (Indian Institute of Technology, BHU)

Session Classification: Parallel: BSM + Cosmology