



Centre for
High Energy Physics

PPC 2024



Search for dark matter decay and annihilation using γ ray observation by Tibet AS $_{\gamma}$ and LHAASO


Abhishek Dubey

Centre for High Energy Physics
Indian Institute of Science, Bengaluru



Based on arXiv:2105.05680 (PRD Letter) & Dubey et al (In Prep.)

In collaboration with
Tarak Nath Maity, Akash Kumar Saha and Ranjan Laha

Dark Matter Indirect detection



Dark Matter Indirect detection



Flux of gamma rays from DM decay/annihilation

DM decay

$$\frac{d\Phi^G}{dE_\gamma} = \frac{1}{4\pi m_\chi \tau_\chi} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

m_χ = DM mass, τ_χ = DM lifetime,
 E_γ, E_e = energy of the prompt photons and prompt electrons/positron
 ρ = DM density profile, which we have taken as NFW profile
 s = line-of-sight distance taken for our galaxy, b, l are Galactic latitude and longitude
 $\tau_{\gamma\gamma}$ = optical depth of photons due to CMB, SL+IR and EBL

 HDMspectra

Flux of gamma rays from DM decay/annihilation

DM decay

$$\frac{d\Phi^G}{dE_\gamma} = \frac{1}{4\pi m_\chi \tau_\chi} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

 HDM Spectra

m_χ = DM mass, τ_χ = DM lifetime,
 E_γ, E_e = energy of the prompt photons and prompt electrons/positron
 ρ = DM density profile, which we have taken as NFW profile
 s = line-of-sight distance taken for our galaxy, b, l are Galactic latitude and longitude
 $\tau_{\gamma\gamma}$ = optical depth of photons due to CMB, SL+IR and EBL

DM annihilation

$$\frac{d\Phi^G}{dE_\gamma} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho^2(s, b, l) B_{sh}(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

Since the annihilation rate depends on the dark matter density squared (and $\langle\varrho^2\rangle \geq \langle\varrho\rangle^2$), the presence of the subhalos will boost the gamma-ray signatures from dark matter annihilation. It is given by B_{sh} (Boost factor).

Flux of gamma rays from DM decay/annihilation

DM decay

$$\frac{d\Phi^G}{dE_\gamma} = \frac{1}{4\pi m_\chi \tau_\chi} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

 HDM Spectra

m_χ = DM mass, τ_χ = DM lifetime,
 E_γ, E_e = energy of the prompt photons and prompt electrons/positron
 ρ = DM density profile, which we have taken as NFW profile
 s = line-of-sight distance taken for our galaxy, b, l are Galactic latitude and longitude
 $\tau_{\gamma\gamma}$ = optical depth of photons due to CMB, SL+IR and EBL

DM annihilation

$$\frac{d\Phi^G}{dE_\gamma} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho^2(s, b, l) B_{sh}(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

Since the annihilation rate depends on the dark matter density squared (and $\langle Q^2 \rangle \geq \langle Q \rangle^2$), the presence of the subhalos will boost the gamma-ray signatures from dark matter annihilation. It is given by B_{sh} (Boost factor).

In our analysis, we have taken both primary and inverse Compton scattering gamma ray flux from Galactic and Extragalactic domain into consideration.

Boost factor

Total Luminosity from DM annihilation

$$\leftarrow L(M) = [1 + B_{\text{sh}}(M)] L_{\text{host}}(M) \leftarrow \text{Luminosity from DM annihilation if there is no substructure.}$$

$$B_{\text{sh}}(M) = \frac{1}{L_{\text{host}}(M)} \int dm \frac{dN}{dm} L_{\text{sh}}(m) [1 + B_{\text{ssh}}(m)]$$

\rightarrow Subhalo mass function

Boost factor

Total Luminosity from DM annihilation


$$L(M) = [1 + B_{\text{sh}}(M)] L_{\text{host}}(M)$$

Luminosity from DM annihilation if there is no substructure.

$$B_{\text{sh}}(M) = \frac{1}{L_{\text{host}}(M)} \int dm \frac{dN}{dm} L_{\text{sh}}(m) [1 + B_{\text{ssh}}(m)]$$

Subhalo mass function

Shin'ichiro Ando et al. 2019



High Energy γ ray detectors


Tibet AS $_{\gamma}$



Present Performance


❑ # of detectors	0.5 m ² x 597
❑ Effective area	~65,700 m ²
❑ Angular resolution	~0.5° @10TeV ~0.2° @100TeV
❑ Energy resolution	~40%@10TeV γ ~20%@100TeV γ

LHAASO




Pointing accuracy ~0.1°
Angular resolution ~0.3°
Energy resolution <20%@6TeV

Photon vs Cosmic ray shower



Sub-PeV diffuse Gamma rays from the Galactic disk



First detection of sub PeV diffuse γ rays by Tibet AS $_{\gamma}$






Fig. γ rays observed by Tibet AS $_{\gamma}$ in Galactic plane in the regions of $|b| < 5^\circ, 25^\circ < l < 100^\circ$

Sub-PeV diffuse Gamma rays from the Galactic disk



Diffuse γ rays observed by LHAASO






Fig. γ rays observed by LHAASO in inner Galaxy plane region of $|b| < 5^\circ, 15^\circ < l < 125^\circ$ and Tibet AS $_\gamma$ in the regions of $|b| < 5^\circ, 25^\circ < l < 100^\circ$

Limit on high Galactic latitude PeV γ -ray flux from Tibet AS $_{\gamma}$


Due to the better sensitivity of Tibet-AS $_{\gamma}$ and higher energy reach compared to MILAGRO, HAWC, and ARGO-YBJ and also more efficient suppression of background EAS produced by protons and atomic nuclei, Tibet-AS $_{\gamma}$ observations can be used to constrain the γ ray flux from the sky outside the Galactic plane ($|b| > 20$ deg.).

Limit on high Galactic latitude PeV γ -ray flux from Tibet AS $_{\gamma}$


Due to the better sensitivity of Tibet-AS $_{\gamma}$ and higher energy reach compared to MILAGRO, HAWC, and ARGO-YBJ and also more efficient suppression of background EAS produced by protons and atomic nuclei, Tibet-AS $_{\gamma}$ observations can be used to constrain the γ ray flux from the sky outside the Galactic plane ($|b| > 20$ deg.).





Results



arXiv:2105.05680 (PRD Letter) + Dubey et al. (In Prep.)



Results



Conclusions


- We have obtained constraints on Dark Matter lifetime and annihilation cross section for different final states using Tibet AS _{γ} and LHAASO observation.
- We have studied the effect of inverse Compton scattering and dark matter substructure which helps put better constrain dark matter parameters.
- We get the most stringent constraints in large region of parameter space for both dark matter decay and annihilation.

Thank You

Email: abhishekd1@iisc.ac.in


Backup

Dark Matter (DM)





- No electric charge
- No or very little baryonic interactions
- Long-lived or stable

Mass scale of dark matter (not to scale)



Results

arXiv:2105.05680 (PRD Letter)



Cosmic Ray measurement by Tibet AS γ







Fig: Amenomori et al., EPJ Web of Conferences 208, 03001 (2019)

P_{IC} and Energy Loss for ICS and Synchrotron



P_{IC} and Energy Loss for ICS and Synchrotron



Esmaili & Serpico 2015

Does cosmic ray measurement by Tibet AS γ tell us anything about the diffuse gamma ray at this sky region ($|b| > 20$ deg.) ?



YES

!

It can give an upper limit on the diffuse gamma rays

Implication of Muon Cut for Tibet ASy





Fig : Gaisser et al., 1991

TABLE II. Implications for γ -ray detection of our calculation of the fluctuations in the number of muons N_μ for cosmic-ray showers with $N_e > 10^5$; see Fig. 3(b).

	$N_\mu < 75$	$N_\mu < 100$	$N_\mu < 200$	$N_\mu < 300$
Percentage of γ -ray signals retained	10%	20%	60%	83%
Level of cosmic-ray background				
Solid line fit	10^{-5}	1.5×10^{-5}	4×10^{-5}	10^{-4}
Dashed line fit	$< 10^{-7}$	10^{-7}	6.6×10^{-7}	4×10^{-6}



Implication of Muon Cut for Tibet ASy



Amenomori et al., PRL,
2019

Implication of Muon Cut for Tibet ASy

Muon
Cut



Amenomori et
al., PRL, 2019


Choose a Muon Cut

Take into account majority of photon induced events

Discard most of the background (CR induced) events

Our detector is not perfect ! Even after the tight muon cut some CR induced shower will get in

Fig:
2107.06541



Upper limits on diffuse gamma ray flux

Upper limit on gamma ray flux

Cosmic Ray flux measured \times Surviving event fraction of Hadronic EAS

=

Efficiency of the selection of Gamma ray EAS




Fig:
Amenomori
et al., EPJ
Web of
Conference
s 208, 03001
(2019)




Fig:
2107.065
41

Flux of gamma rays from DM decay

ray flux of direct production from DM decay

$$\frac{d\Phi_\gamma^G}{dE_\gamma} = \frac{1}{4\pi m_\chi \tau_\chi} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

$$\frac{d\phi_\gamma^{EG}}{dE_\gamma} = \frac{\Omega_{DM}\rho_{cr}}{4\pi m_\chi \tau_\chi} \int \frac{dz}{H(z)} \frac{dN_\gamma}{dE_\gamma} \Big|_{E'_\gamma=E_\gamma(1+z)} e^{-\tau_{\gamma\gamma}(E_\gamma, z)}$$

m = DM mass, τ = DM lifetime,

E, E_e = energy of the prompt photons and prompt electrons/positron

ρ = DM density profile, which we have taken as NFW profile

s = line-of-sight distance taken for our galaxy, b, l are Galactic latitude and longitude

τ = optical depth of photons due to CMB, SLL+IR and EBL



HDM Spectra

ray flux of Inverse compton production from DM decay

$$\frac{d\Phi_{IC\gamma}}{dE_\gamma d\Omega} = \frac{2}{E_\gamma} \frac{1}{4\pi m_\chi \tau_\chi} \int_{m_e}^{m_\chi/2} dE_s \frac{dN_e}{dE_e} (E_s) \int_{l.o.s.} ds (\rho(s, b, l)) \int_{m_e}^{E_s} dE \frac{\sum_i \mathcal{P}_{IC}^i(E_\gamma, E, s, b, l)}{b(E, s, b, l)} I(E, E_s, s, b, l),$$

$$\frac{d\Phi_{EG\gamma}}{dE_\gamma} (E_\gamma, z) = c \frac{1}{E_\gamma} \int_z^\infty dz' \frac{1}{H(z')(1+z')} \left(\frac{1+z}{1+z'} \right)^3 j_{EG\gamma}(E'_\gamma, z') e^{-\tau(E_\gamma, z, z')}.$$

$$j_{EG\gamma}^{IC}(E'_\gamma, z') = \frac{2}{\tau_\chi} \frac{\bar{\rho}(z')}{m_\chi} \int_{m_e}^{m_\chi/2} dE_e \frac{\mathcal{P}_{IC}^{CMB}(E'_\gamma, E_e, z')}{b_{IC}^{CMB}(E_e, z')} \int_{E_e}^{m_\chi/2} d\tilde{E}_e \frac{d\tilde{N}_e}{d\tilde{E}_e}$$

P_{IC} is ICS radiative power and b_{IC} is the energy loss of electrons/positrons due to ICS and Synchrotron radiation.

Flux of gamma rays from DM annihilation

ray flux of direct production from DM annihilation

$$\frac{d\Phi^G}{dE_\gamma} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN}{dE_\gamma} \int_0^\infty ds \rho^2(s, b, l) B_{sh}(s, b, l) e^{-\tau_{\gamma\gamma}(E_\gamma, s, b, l)}$$

$$\frac{d\phi_\gamma^{\text{EG}}}{dE_\gamma} = \frac{\langle\sigma v\rangle \Omega_{\text{DM}}^2 \rho_{\text{cr}}^2}{8\pi m_\chi^2} \int \frac{dz}{H(z)} \langle\delta^2(z)\rangle (1+z)^3 \left. \frac{dN_\gamma}{dE_\gamma} \right|_{E'_\gamma=E_\gamma(1+z)} e^{-\tau_{\gamma\gamma}(E_\gamma, z)}$$

ray flux of inverse Compton production from DM annihilation

$$\frac{d\Phi_{\text{IC}\gamma}}{dE_\gamma d\Omega} = \frac{2}{E_\gamma} \frac{\langle\sigma v\rangle}{4\pi m_\chi^2} \int_{m_e}^{m_\chi/2} dE_s \frac{dN_e}{dE_e}(E_s) \int_{\text{l.o.s.}} ds \frac{1}{2} B_{sh}(s, b, l) (\rho(s, b, l))^2 \int_{m_e}^{E_s} dE \frac{\sum_i \mathcal{P}_{\text{IC}}^i(E_\gamma, E, s, b, l)}{b(E, s, b, l)} I(E, E_s, s, b, l),$$

$$\frac{d\Phi_{\text{EG}\gamma}}{dE_\gamma}(E_\gamma, z) = c \frac{1}{E_\gamma} \int_z^\infty dz' \frac{1}{H(z')(1+z')} \left(\frac{1+z}{1+z'} \right)^3 j_{\text{EG}\gamma}(E'_\gamma, z') e^{-\tau(E_\gamma, z, z')}.$$

$$j_{\text{EG}\gamma}^{\text{IC}}(E'_\gamma, z') = 2 \langle\delta^2(z)\rangle \frac{1}{2} \langle\sigma v\rangle \left(\frac{\bar{\rho}(z')}{m_\chi} \right)^2 \int_{m_e}^{m_\chi/2} dE_e \frac{\mathcal{P}_{\text{IC}}^{\text{CMB}}(E'_\gamma, E_e, z')}{b_{\text{IC}}^{\text{CMB}}(E_e, z')} \int_{E_e}^{m_\chi/2} d\tilde{E}_e \frac{d\tilde{N}_e}{d\tilde{E}_e}$$

B_{sh} and $\langle\delta^2\rangle$ are Boost factor and Clumping factor due to dark matter substructure. Since the annihilation rate depends on the dark matter density squared (and $\langle\delta^2\rangle \geq \langle\Delta\rangle^2$), the presence of the subhalos will boost the gamma-ray signatures from dark matter annihilation.


Boost factor and Clumping factor

$$\langle \delta^2 \rangle = \left(\frac{1}{\Omega_m \rho_c} \right)^2 \int dM \frac{dn(M, z)}{dM} [1 + B_{\text{sh}}(M)] \times \int dV \rho_{\text{host}}^2(r | M)$$



Hiroshima, et al. 2018



Halo mass function



Inner and outer Galaxy constraints from LHAASO



Background for LHAASO diffuse gamma ray flux

