



Contribution ID: 105

Type: Oral

Tidal heating as a direct probe of Strange matter inside Neutron stars

The cores of neutron stars (NS) reach densities several times the nuclear saturation density and could contain strangeness containing exotic particles such as hyperons. During the binary inspiral, viscous processes inside the NS matter can damp out the tidal energy induced by the companion and convert this to thermal energy to heat up the star. In this work, we demonstrate that the bulk viscosity originating from the non-leptonic weak interactions involving hyperons is several orders of magnitude higher than the standard neutron matter shear viscosity in the relevant temperature range of $10^6 - 10^9$ K and for heavier mass NSs ($M \geq 1.6M_\odot$) that contain a significant fraction of hyperons in their core, the bulk viscosity can heat up the stars upto $0.1 - 1$ MeV before the final merger. We also show that this “tidal heating” process introduces a net phase shift in the order of $0.1 - 0.5$ rad in the gravitational wave (GW) signal that can potentially be detected using current and future generation GW detectors. Such a detection would be the first direct confirmation of the presence of hyperons inside the NS core, having a great significance for the study of dense matter under extreme condition.

Email

suprovo@iucaa.in

Affiliation

Inter-University Centre for Astronomy and Astrophysics (IUCAA)

Author: GHOSH, SUPROVO

Co-authors: PRADHAN, Bikram (Inter University Center For Astronomy and Astrophysics, Pune, India); Dr CHATTERJEE, Debarati (IUCAA)

Presenter: GHOSH, SUPROVO

Session Classification: Gravitational Waves

Track Classification: Gravitational Waves