



Contribution ID: 385

Type: **not specified**

Cosmic Millicharge Background and Reheating Probes

Tuesday 14 May 2024 15:15 (15 minutes)

We demonstrate that the searches for dark sector particles can provide probes of reheating scenarios, focusing on the cosmic millicharge background produced in the early universe. We discuss two types of millicharge particles (mCPs): either with, or without, an accompanying dark photon. These two types of mCPs have distinct theoretical motivations and cosmological signatures. We discuss constraints from the overproduction and mCP-baryon interactions of the mCP without an accompanying dark photon, with different reheating temperatures. We also consider the ΔN_{eff} constraints on the mCPs from kinetic mixing, varying the reheating temperature. The regions of interest in which the accelerator and other experiments can probe the reheating scenarios are identified for both scenarios. These probes can potentially allow us to set an upper bound on the reheating temperature down to ~ 10 MeV, much lower than the previously considered upper bound from inflationary cosmology at around $\sim 10^{16}$ GeV. In addition, we derive a new “distinguishability condition”, in which the two mCP scenarios may be differentiated by combining cosmological and theoretical considerations.

Mini Symposia (Invited Talks Only)

Author: GAN, Xucheng**Co-author:** TSAI, Yu-Dai (University of California, Irvine)**Presenter:** GAN, Xucheng**Session Classification:** Cosmology & Dark Energy**Track Classification:** Cosmology & Dark Energy