

DPF-PHENO 2024

Contribution ID: 390

Type: **not specified**

Exploring $0\nu\beta\beta$ decay and leptogenesis in an extended seesaw model

Wednesday 15 May 2024 14:30 (15 minutes)

We investigate the possibility of neutrinoless double beta decay ($0\nu\beta\beta$) and leptogenesis within a low-scale seesaw mechanism with additional sterile neutrinos. The general effective field theory (EFT) considerations suggest that if there are experimentally observable signatures in $0\nu\beta\beta$ decay and the lepton asymmetry generated by the right-handed neutrinos, the low-scale leptogenesis is likely to be unviable. However, in this work, we show that in the context of low-scale resonant leptogenesis, one can obtain the observed BAU and observable signatures in $0\nu\beta\beta$ decay in the presence of additional sterile neutrinos. In this framework, the neutrino masses are naturally suppressed by the extended seesaw parameter, μ , rather than introducing small Yukawa couplings in other leptogenesis scenarios. This can lead to both observable experimental signatures in $0\nu\beta\beta$ decay and charged lepton flavor violation (cLFV) as well as large washout effects. The resonant leptogenesis mechanism with light neutrino masses can overcome the latter, even in the presence of experimentally accessible $0\nu\beta\beta$ -decay and cLFV signatures. We have shown that the KamLAND-Zen experiment is sensitive to keV-MeV scale sterile neutrino masses, and future ton-scale experiments offer potential signals while maintaining viable leptogenesis.

Mini Symposia (Invited Talks Only)

Authors: Dr SENAPATI, Supriya (University of Massachusetts Amherst); LI, Gang (Sun Yat-sen University, Zhuhai); RAMSEY-MUSOLF, Michael (University of Massachusetts Amherst); URRUTIA QUIROGA, Sebastian (University of Washington, Seattle)

Presenter: Dr SENAPATI, Supriya (University of Massachusetts Amherst)

Session Classification: Neutrino Physics

Track Classification: Neutrino Physics