



Contribution ID: 398

Type: **not specified**

## Light dark matter, rare $B$ decays with missing energy in $L_\mu - L_\tau$ model

We investigate the phenomenology of light GeV-scale fermionic dark matter in  $U(1)_{L_\mu - L_\tau}$  gauge extension of the Standard Model. Heavy neutral fermions alongside with a  $S_1(\bar{3}, 1, 1/3)$  scalar leptoquark and an inert scalar doublet are added to address the flavor anomalies and light neutrino mass respectively. The light gauge boson associated with  $U(1)_{L_\mu - L_\tau}$  gauge group mediates dark to visible sector and helps to obtain the correct relic density. Aided with a colored scalar, we constrain the new model parameters by using the branching ratios of various  $b \rightarrow sll$  and  $b \rightarrow s\gamma$  decay processes as well as the lepton flavour non-universality observables  $R_{K^{(*)}}$  and then show the implication on the branching ratios of some rare semileptonic  $B \rightarrow (K^{(*)}, \phi) + \text{missing energy, processes}$ .

### Mini Symposia (Invited Talks Only)

**Author:** SAHOO, Suchismita

**Co-authors:** MOHANTA, Rukmani; SINGIRALA, Shivaramakrishna (School of physics, University of Hyderabad, Hyderabad 500046)

**Presenter:** SAHOO, Suchismita

**Session Classification:** Dark Matter

**Track Classification:** Dark Matter