



Contribution ID: 140

Type: Oral

## How bright can the brightest neutrino source be?

*Thursday 10 August 2017 16:30 (15 minutes)*

After the discovery of extraterrestrial high-energy neutrinos, the next major goal of neutrino telescopes will be identifying astrophysical objects that produce them. The flux of the brightest source  $F_{\max}$ , however, cannot be probed by studying the diffuse neutrino intensity. We aim at constraining  $F_{\max}$  by adopting a broken power-law flux distribution, a hypothesis supported by observed properties of any generic astrophysical sources. The first estimate of  $F_{\max}$  comes from the fact that we can only observe one universe, and hence, the expected number of sources above  $F_{\max}$  cannot be too small compared with one. For abundant source classes such as starburst galaxies, this one-source constraint yields a value of  $F_{\max}$  that is an order of magnitude lower than the current upper limits from point-source searches. Then we derive upper limits on  $F_{\max}$  assuming that the angular power spectrum is consistent with neutrino shot noise yet. We find that the limits obtained with ongoing muon neutrinos in IceCube can already be quite competitive, especially for rare but bright source populations such as blazars. The limits will improve nearly quadratically with exposure, and therefore be even more powerful for the next generation of neutrino telescopes.

**Author:** Dr ANDO, Shin'ichiro (GRAPPA, University of Amsterdam)

**Co-authors:** FEYEREISEN, Michael (University of Amsterdam); FORNASE, Mattia (GRAPPA Institute (University of Amsterdam))

**Presenter:** Dr ANDO, Shin'ichiro (GRAPPA, University of Amsterdam)

**Session Classification:** Neutrinos

**Track Classification:** Neutrinos (astrophysical, atmospheric)