



Contribution ID: 299

Type: Oral

## A hadronic origin of the high-energy gamma rays from LMC

*Tuesday 8 August 2017 14:45 (15 minutes)*

It has been suggested that high-energy gamma-ray emission ( $>100\text{MeV}$ ) of nearby star-forming galaxies may be produced predominantly by cosmic rays colliding with the interstellar medium through neutral pion decay. The pion-decay mechanism predicts a unique spectral signature in the gamma-ray spectrum, characterized by a fast rising spectrum and a spectral break below a few hundreds of MeV. We here report the evidence of a spectral break around 500 MeV in the disk emission of Large Magellanic Cloud (LMC), which is found in the analysis of the gamma-ray data extending down to 60 MeV observed by *Fermi*-Large Area Telescope. The break is well consistent with the pion-decay model for the gamma-ray emission, although leptonic models, such as the electron bremsstrahlung emission, cannot be ruled out completely.

**Authors:** Dr TANG, Qingwen (Nanchang University & CCAPP); Dr LIU, Ruoyu (Max-Planck-Institut für Kernphysik); Prof. WANG, Xiangyu (Nanjing University)

**Presenter:** Dr TANG, Qingwen (Nanchang University & CCAPP)

**Session Classification:** Gamma rays

**Track Classification:** Gamma rays