



Contribution ID: 162

Type: Talk

## Impact of High Energy Electron Neutrino (Anti-neutrino) Events on NOvA Oscillation Sensitivities

Wednesday 14 December 2022 12:15 (15 minutes)

NOvA is a two-detector, long-baseline neutrino oscillation experiment located at Fermilab, Batavia, IL, USA. It is primarily designed to constrain neutrino oscillation parameters using muon neutrino (anti-neutrino) disappearance data and electron neutrino (anti-neutrino) appearance data. NOvA detects neutrinos from Fermilab's Neutrinos at Main Injector (NuMI) beamline. The unoscillated muon neutrino and beam  $\nu_e$  events are observed by the NOvA Near Detector (ND), which is 100m underground and at a distance of 1km from the beam source. The Far Detector (FD), situated 809 km from the ND, is in Ash River, MN, USA, and observes  $\nu_\mu$  and  $\nu_e$  events after oscillations.

Traditionally, NOvA has used  $\nu_e$  events in the energy range  $1 < E_\nu < 4$  GeV for 3-flavor neutrino oscillation analyses to constrain the neutrino oscillation parameters. In this study, we looked at the impact of including high-energy neutrino events with energies up to 12 GeV in the analysis with the aim of constraining the beam electron neutrino/antineutrino background events. Event count predictions after adding these high-energy sample events and the latest three flavor oscillation results of the NOvA experiment are presented.

### Session

Neutrino Physics

**Author:** SINGH, Ishwar (University of Delhi)

**Co-authors:** Prof. CHOUDHARY, Brajesh (University of Delhi); Dr SINGH, Prabhjot (Queen Mary University of London)

**Presenter:** SINGH, Ishwar (University of Delhi)

**Session Classification:** WG7 - Neutrino Physics