

XXV DAE-BRNS High Energy Physics Symposium 2022



Contribution ID: 173

Type: **Talk**

Black Hole Normal Modes and Spectral Form Factor

Monday 12 December 2022 17:45 (15 minutes)

We consider a black hole with a stretched horizon as a toy model for a fuzzball microstate. The stretched horizon provides a cut-off, and therefore one can determine the normal (as opposed to quasi-normal) modes of a probe scalar in this geometry. For the BTZ black hole, we compute these as a function of the level n and the angular quantum number J . Though conventional level repulsion is absent in this system, we find that the Spectral Form Factor (SFF) shows clear evidence for a dip-ramp-plateau structure with a linear ramp of slope ~ 1 on a log-log plot, with or without ensemble averaging. We show that this is a robust feature of stretched horizons by repeating our calculations on 2d Rindler $\times S^1$ geometry. We also observe that this is not a generic feature of integrable systems, as illustrated by standard examples like integrable billiards and random 2-site coupled SYK model, among others.

Session

Formal Theory

Authors: KUMAR, A. Preetham (Indian Institute of Science); KUNDU, Arnab (SINP); KRISHNAN, Chethan (Indian Institute of Science); DAS, Suman (Saha Institute of Nuclear Physics)

Presenter: DAS, Suman (Saha Institute of Nuclear Physics)

Session Classification: WG3-Formal Theory