



Contribution ID: 305

Type: Poster

Secluded Dark Sector and Muon $(g - 2)$ in the Light of Fast Expanding Universe

Friday 16 December 2022 14:00 (1 hour)

The lack of information before Big Bang Nucleosynthesis (BBN) allow us to assume the presence of a new species ϕ whose energy density redshifts as $a^{-(4+n)}$ where $n > 0$ and a is the scale factor. In this non-standard cosmological setup, we have considered $U(1)_{L_\mu-L_\tau} \otimes U(1)_X$ gauge extension of the Standard Model (SM) and studied different phases of the cosmological evolution of a thermally decoupled dark sector such as leak-in, freeze-in, reannihilation, and late-time annihilation. This non-standard cosmological setup facilitates a larger portal coupling (ϵ) between the dark and the visible sectors even when the two sectors are not in thermal equilibrium. The dark sector couples with the μ and τ flavored leptons of the SM due to the tree level kinetic mixing between $U(1)_X$ and $U(1)_{L_\mu-L_\tau}$ gauge bosons. We show that in our scenario it is possible to reconcile the dark matter relic density and muon $(g - 2)$ anomaly. In particular, we show that for $3 \times 10^{-4} \leq \epsilon \leq 10^{-3}$, $30\text{MeV} \leq m_{Z'}$, $300\text{MeV} \leq m_\chi$, $n = 4$, and $1\text{TeV} \leq m_\chi \leq 10\text{TeV}$ relic density constraint of dark matter, constraint from muon $(g - 2)$ anomaly, and other cosmological, astrophysical constraints are satisfied.

Session

Astroparticle Physics and Cosmology

Authors: Mr GANGULY, Sougata (Indian Association for the Cultivation of Science); Prof. ROY, Sourov (Indian Association for the Cultivation of Science); TAPADAR, Ananya (Indian Association for the Cultivation of Science)

Presenter: TAPADAR, Ananya (Indian Association for the Cultivation of Science)

Session Classification: Poster - 4