9th Shivalik HEPCATS (High Energy Physics, Cosmology, Astronomy: Theory and Simulations)



Contribution ID: 11

Type: not specified

Axial charmonium production in $e^-e^+ \rightarrow \gamma^* \rightarrow H' + H''$ at $\sqrt{s} = 10.6$ GeV at B-factories

Saturday 27 January 2024 11:50 (20 minutes)

We have studied the production of ground and excited axial charmonia in processes, $e^-e^+ \rightarrow h_c(nP) + \eta_c(n'S)$, and $e^-e^+ \rightarrow \chi_{c1}(nP) + h_c(n'P)$ [1] for n, n' = 1, 2, through leading order (LO) tree-level diagrams $\sim O(\alpha_{em}\alpha_s)$, which proceed through exchange of a virtual photon and an internal gluon line connecting two quark lines in the triangle quark loop part of the diagram at center of mass energy, $\sqrt{s} = 10.6 GeV$. We employ the framework of 4×4 Bethe-Salpeter equation, and calculate their cross sections [1], which are compared with recent NRQCD and Relativistic quark model calculations. For both the above processes, we have drawn plots of total cross sections versus \sqrt{s} for n, n' = 1, 2, as well as plots of differential cross section versus $\cos\Theta$ at different center of mass energies. These studies might be of interest for future experiments at B-factories, since h_c, η_c , and χ_{c1}, h_c production might provide opportunities for observing h_c with higher statistics in future. Also our cross sections may provide guidance for future experiments at B-factories. Reference:

[1] M.Narang, S.Bhatnagar, Few-Body Syst. 64, 82 (2023).

Author: Ms NARANG, Monika (Chandigarh University)

Co-author: Prof. BHATNAGAR, Shashank (Chandigarh University)

Presenter: Ms NARANG, Monika (Chandigarh University)