XXVI DAE-BRNS High Energy Physics Symposium 2024



Contribution ID: 251

Type: Postar

Study of transverse polarization asymmetry $\Lambda_b \rightarrow nl^+l^-$ with non-universal Z' model

Baryonic decays which involve $b \rightarrow d$ are very sensitive to new physics effects. Recent experimental observations of $\Lambda_b^0 \to p K^- \mu^+ \mu^-$ decay motivate the theorist to study baryonic decay [1]. The $\Lambda_b \to n l^+ l^-$ decays are forbidden at the tree level in SM. It provides opportunities to test NP models like the leptoquark model [2], two-Higgs doublet model (2HDM) [3], non-universal Z' model [4] and fermion fourth generation model [5]. In the theoretical background, many attempts have been made in different approaches like light cone sum rule (LCSR), lattice quantum chromodynamics (LQCD), the 2HDM and the Bethe-Salpeter equation approach. In the Bethe-Salpeter equation approach the obtained branching ratios $Br(\Lambda_b \rightarrow nl^+l^-) \times 10^8$ are $6.79^{+8.6}_{-8.2}$ (for l = e), $4.08^{+5.44}_{-1.19}$ (for l =) and $(2.9)^{+3.7}_{-0.78}$ (for l =) [6]. In the context of LQCD and LCSR [7] branching ratios $Br(\Lambda_b \rightarrow nl^+l^-) \times 10^8$ are obtained as (3.19±0.32),(3.79±0.46) (for l = e), (3.15±0.29),(3.76±0.42) (for l =) and (1.42±0.32),(1.65±0.19) (for l =) respectively. Later on these results are updated in LCSR [8] $Br(\Lambda_b \rightarrow nl^+l^-) \times 10^8$ as (8±2) (for l = e), (7±2) (for l =) and (2±0.4) (for l =). In the relativistic quarkdiquark model of baryons, the branching ratios $Br(\Lambda_b \rightarrow nl^+l^-) \times 10^8$ are 3.81 (for l = e), 3.75 (for l =) and 1.21 (for l = 0 [9]. In this work, we intend to study $\Lambda_b \rightarrow nl^+l^-$ decays in non-universal Z' model. We will estimate the transverse polarization asymmetry for $\Lambda_b \rightarrow nl^+l^-$ decays. To determine new physics, we will use quark coupling from $B_d^0 - B_d^0$ mixing. We hope these results will help the experimental community to explore these kinds of decays at the LHCb/Belle II detector in the upcoming time.

Acknowledgement

R. Ray acknowledges DST, Govt. of India for providing the INSPIRE Fellowship (IF210427) during his research.

References

- 1. R. Aaij et al. (LHCb Collaboration), JHEP 05, 05 (2020).
- 2. M. Schmaltz and Y.M. Zhong, JHEP 01, 132 (2019).
- 3. H. B. Maïto, A. Falkowski, D. Fontes, J.C. Romao and J. P. Silva, Eur. Phys. J. C 77, 176 (2017).
- 4. P. Langacker, Rev. Mod. Phys. 81, 1199 (2009).
- 5. R. Mohanta and A. K. Giri, Phys. Rev. D 85, 014008 (2012).
- 6. L. L. Liu, C. Wang, X. Kang and X. H. Guo, Eur. Phys. J. C 80, 193 (2020).
- 7. K. Azizi, M. Bayar, Y. Sarac and H. Sundu, J. Phys. G: Nucl. Part. Phys. 37, 115007 (2010).
- 8. T. M. Aliev, T. Barakat and M. Savcı, Phys. Rev. D 98, 035033 (2018).
- 9. R. N. Faustov and V.O. Galkin, Mod. Phys. Lett. A 32, 1750125 (2017).

Field of contribution

Phenomenology

Author: RAY, raja (National Institute of Technology Durgapur)

Co-author: Dr SAHOO, Sukhdev (National Institute of Technology Durgapur)

Presenter: RAY, raja (National Institute of Technology Durgapur)

Track Classification: Beyond the standard model